

10/598,520

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:sssptal600txm

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

\* \* \* \* \* Welcome to STN International \* \* \* \* \*

NEWS 1 Web Page for STN Seminar Schedule - N. America  
NEWS 2 APR 02 CAS Registry Number Crossover Limits Increased to  
500,000 in Key STN Databases  
NEWS 3 APR 02 PATDPAFULL: Application and priority number formats  
enhanced  
NEWS 4 APR 02 DWPI: New display format ALLSTR available  
NEWS 5 APR 02 New Thesaurus Added to Derwent Databases for Smooth  
Sailing through U.S. Patent Codes  
NEWS 6 APR 02 EMBASE Adds Unique Records from MEDLINE, Expanding  
Coverage back to 1948  
NEWS 7 APR 07 50,000 World Traditional Medicine (WTM) Patents Now  
Available in Cplus  
NEWS 8 APR 07 MEDLINE Coverage Is Extended Back to 1947  
NEWS 9 JUN 16 WPI First View (File WPIFV) will no longer be  
available after July 30, 2010  
NEWS 10 JUN 18 DWPI: New coverage - French Granted Patents  
NEWS 11 JUN 18 CAS and FIZ Karlsruhe announce plans for a new  
STN platform  
NEWS 12 JUN 18 IPC codes have been added to the INSPEC backfile  
(1969-2009)  
NEWS 13 JUN 21 Removal of Pre-IPC 8 data fields streamline displays  
in CA/Cplus, CASREACT, and MARPAT  
NEWS 14 JUN 21 Access an additional 1.8 million records exclusively  
enhanced with 1.9 million CAS Registry Numbers --  
EMBASE Classic on STN  
NEWS 15 JUN 28 Introducing "CAS Chemistry Research Report": 40 Years  
of Biofuel Research Reveal China Now Atop U.S. in  
Patenting and Commercialization of Bioethanol  
NEWS 16 JUN 29 Enhanced Batch Search Options in DGENE, USGENE,  
and PCTGEN  
NEWS 17 JUL 19 Enhancement of citation information in INPADOC  
databases provides new, more efficient competitor  
analyses  
NEWS 18 JUL 26 CAS coverage of global patent authorities has  
expanded to 61 with the addition of Costa Rica  
NEWS 19 SEP 15 MEDLINE Cited References provide additional  
relevant records with no additional searching.  
NEWS 20 OCT 04 Removal of Pre-IPC 8 data fields streamlines  
displays in USPATFULL, USPAT2, and USPATOLD.  
NEWS 21 OCT 04 Precision of EMBASE searching enhanced with new  
chemical name field  
NEWS 22 OCT 06 Increase your retrieval consistency with new formats or  
for Taiwanese application numbers in CA/Cplus.  
NEWS 23 OCT 21 CA/Cplus kind code changes for Chinese patents  
increase consistency, save time  
NEWS 24 OCT 22 New version of STN Viewer preserves custom  
highlighting of terms when patent documents are  
saved in .rtf format  
NEWS 25 OCT 28 INPADOCDB/INPAFAMDB: Enhancements to the US national  
patent classification.  
NEWS 26 NOV 03 New format for Korean patent application numbers in  
CA/Cplus increases consistency, saves time.  
NEWS 27 NOV 04 Selected STN databases scheduled for removal on  
December 31, 2010

NEWS EXPRESS FEBRUARY 15 10 CURRENT WINDOWS VERSION IS V8.4.2,  
AND CURRENT DISCOVER FILE IS DATED 07 JULY 2010.

McIntosh




10/598,520

L2 15 SEA SSS SAM L1

=> d scan

L2 15 ANSWERS REGISTRY COPYRIGHT 2010 ACS on STN  
IN INDEX NAME NOT YET ASSIGNED  
MF C18 H22 N6 O5

Absolute stereochemistry.




\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):5

L2 15 ANSWERS REGISTRY COPYRIGHT 2010 ACS on STN  
IN Acetic acid, [(6-amino-9-beta-D-ribofuranosyl)-9H-purin-2-yl]oxy]- (9CI)  
MF C12 H15 N5 O7


Absolute stereochemistry.



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

L2 15 ANSWERS REGISTRY COPYRIGHT 2010 ACS on STN  
IN Adenosine, 2-[(3,6-dimethyl-6-heptenyl)oxy]-, (R)- (9CI)  
MF C19 H29 N5 O5

Absolute stereochemistry.




McIntosh

10/598,520

\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

L2 15 ANSWERS REGISTRY COPYRIGHT 2010 ACS on STN  
IN Adenosine, 2-[(2,2-diphenylethyl)amino]- (9CI)  
MF C24 H26 N6 O4


Absolute stereochemistry.



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*


L2 15 ANSWERS REGISTRY COPYRIGHT 2010 ACS on STN  
IN Cyclohexanepropanoic acid, 4-[2-[(6-amino-9-β-D-ribofuranosyl-9H-purin-2-yl)amino]ethyl]-  
MF C21 H32 N6 O6  
CI COM

Absolute stereochemistry.



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

L2 15 ANSWERS REGISTRY COPYRIGHT 2010 ACS on STN  
IN Adenosine, 2-[[2-(6,6-dimethylbicyclo[3.1.1]hept-2-en-2-yl)ethyl]amino]- (9CI)  
MF C21 H30 N6 O4



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

McIntosh

10/598,520

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):0

=>

Uploading C:\Program Files\Stnexp\Queries\10598520a.str

L3           STRUCTURE UPLOADED

=> s l3

SAMPLE SEARCH INITIATED 13:39:51 FILE 'REGISTRY'

SAMPLE SCREEN SEARCH COMPLETED -       1831 TO ITERATE

100.0% PROCESSED       1831 ITERATIONS

8 ANSWERS

SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS:   ONLINE   \*\*COMPLETE\*\*

BATCH   \*\*COMPLETE\*\*

PROJECTED ITERATIONS:       34053 TO       39187

PROJECTED ANSWERS:         8 TO       328

L4           8 SEA SSS SAM L3


=> d scan

L4   8 ANSWERS   REGISTRY   COPYRIGHT 2010 ACS on STN

IN   Adenosine, 2-[[4-(aminocarbonyl)phenyl]amino]- (9CI)

MF   C17 H19 N7 O5

Absolute stereochemistry.



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):0

=> s l3 full

FULL SEARCH INITIATED 13:40:03 FILE 'REGISTRY'

FULL SCREEN SEARCH COMPLETED -       36053 TO ITERATE

100.0% PROCESSED       36053 ITERATIONS

148 ANSWERS

SEARCH TIME: 00.00.01

L5           148 SEA SSS FUL L3

=> file caplus

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

197.91

198.13

FILE 'CAPLUS' ENTERED AT 13:40:10 ON 17 NOV 2010

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2010 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the

McIntosh

10/598,520

American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 17 Nov 2010 VOL 153 ISS 21  
FILE LAST UPDATED: 16 Nov 2010 (20101116/ED)  
REVISED CLASS FIELDS (/NCL) LAST RELOADED: Aug 2010  
USPTO MANUAL OF CLASSIFICATIONS THESAURUS ISSUE DATE: Aug 2010

CAPLUS now includes complete International Patent Classification (IPC) reclassification data for the third quarter of 2010.

CAS Information Use Policies apply and are available at:

<http://www.cas.org/legal/infopolicy.html>

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s 15

L6 195 L5

=> d bib abs hitstr 1-195 16

L6 ANSWER 1 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 2010:819643 CAPLUS  
DN 153:154787  
TI Combination of monosaccharides and adenosine for cosmetic uses  
IN Laboureaux, Julien; Simonnet, Jean-Thierry; Portes, Pascal  
PA L'Oreal, Fr.  
SO U.S. Pat. Appl. Publ., 17pp.  
CODEN: USXXCO  
DT Patent  
LA English  
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 20100168049	A1	20100701	US 2009-649367	20091230
	FR 2940611	A1	20100702	FR 2008-59151	20081230
	JP 2010155834	A	20100715	JP 2009-298271	20091228
	KR 2010080437	A	20100708	KR 2009-132924	20091229
	EP 2204154	A1	20100707	EP 2009-181014	20091230
	R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LI, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR, AL, BA, RS				
	CN 101856313	A	20101013	CN 2009-10265295	20091230
PRAI	FR 2008-59151	A	20081230		
	US 2009-144756P	P	20090115		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OS MARPAT 153:154787

AB The present invention relates to a composition, especially a cosmetic and/or dermatol. composition, containing, in a physiol. acceptable medium, a combination of a monosaccharide chosen from mannose, rhamnose and a mixture thereof, and of an addnl. compound chosen from adenosine, an analog thereof and a mixture thereof. Thus, a cosmetic formulation contained Hostacerin AMPS 1.00, cyclohexasiloxane 5.00, apricot kernel oil 7, Isononyl isononanoate 7, stearyl alc. 0.30, glyceryl stearate/PEG-100 stearate 0.70, Dimyristyl tartrate/cetearyl alc./C12-15-pareth-7/PPG-25 laureth-25 0.50, xanthan gum 0.20, mannose 2.5, rhamnose 2.5, adenosine 0.1, preservatives 0.5, and water qs to 100%.

IT 53296-10-9, 2-Phenylaminoadenosine

RL: COS (Cosmetic use); THU (Therapeutic use); BIOL (Biological study); USES (Uses)


(combination of monosaccharides and adenosine for cosmetic uses)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.

McIntosh



L6 ANSWER 2 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
 AN 2010:178356 CAPLUS  
 DN 152:255274  
 TI Administration by infusion for the treatment of ischemic effects  
 IN Weber, Uno Jakob; Gotfredsen, Jacob  
 PA Neurokey A/S, Den.  
 SO PCT Int. Appl., 160pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 FAN.CNT 9

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2010015260	A2	20100211	WO 2009-DK50196	20090807
	WO 2010015260	A3	20100617		
	W:	AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GE, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW			
	RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GE, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA			
	WO 2009071095	A2	20090611	WO 2008-DK50293	20081205
	WO 2009071095	A3	20090723		
	W:	AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GE, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW			
	RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GE, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA			
PRAI	DK 2008-1079	A	20080807		
	WO 2008-DK50293	A	20081205		
	DK 2007-1742	A	20071205		
	DK 2007-1743	A	20071205		
	DK 2008-716	A	20080523		
	DK 2008-1105	A	20080815		
	DK 2008-1337	A	20080926		

OS MARPAT 152:255274

AB The invention relates to the induction of hypothermia in humans, male and female, at any age, by use of a pharmaceutical composition of formula I, wherein R1 and R2 are chemical moieties or chemical bonds, wherein R1 is selected from the group of: C, S, N, O, optionally substituted with C, S, N, O, P, OH, 5 hydrogen, alkyl, alkenyl, alkynyl, any of which may or may not be branched or comprise substituents such as phosphate, cycloalkyl, etc., wherein R2 is selected from the group of: C, S, N, O, optionally substituted one or more times with C, S, N, O, P, OH, hydrogen, alkoxy, alkyl, alkenyl, alkynyl, Ph, di-Ph, benzyl, amine (NH), halogen, substituted lower alkyl, alkenyl, aryl, heterocycloalkyl, etc., to be administered parenterally by infusion or injection, comprising at least

one compound selected among vanilloid receptor agonists, capsaicinoids or capsaicinoid-like agonists reaching and binding to vanilloid receptors, and cannabinoids or cannabimimetic agonists reaching and binding to cannabinoid receptors, and adenosine receptor agonists, and neurotensin receptor agonists, and thyroxine derivs., and cytochrome c inhibitors, and oxygen tension reducers, thereby inducing hypothermia, thus benefiting patients suffering from illnesses characterized by tissue anoxia.

IT 53296-10-9, CV-1808


RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(induction of therapeutic hypothermia by pharmaceutical infusion of medications for prophylaxis, mono- and combination therapy of ischemia)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



L6 ANSWER 3 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 2009:1566750 CAPLUS

DN 152:67621

TI  $\beta$ -Adrenergic receptor agonists for the treatment of B-cell proliferative disorders

IN Rickles, Richard; Lee, Margaret S.

PA CombinatoRx, Inc., USA

SO PCT Int. Appl., 111 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2009151569	A2	20091217	WO 2009-US3449	20090608
	WO 2009151569	A3	20100225		
	W:	AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW			
	RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA			
	US 20100009934	A1	20100114	US 2009-480034	20090608
PRAI	US 2008-60064P	P	20080609		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT


AB The invention discloses a method for treating a B-cell proliferative disorder by administering to a patient a  $\beta$ -Adrenergic receptor (BAR) agonist, e.g., formulated for administration by a route other than inhalation (such as for oral or i.v. administration), in an amount effective to treat the B-cell proliferative disorder. The BAR agonist may be administered as a monotherapy or in combination with one or more other agents, e.g., a PDE inhibitor, an A2A receptor agonist, or an antiproliferative compound, in amts. that together are effective to treat the B-cell proliferative disorder. The invention further discloses pharmaceutical compns. and kits including a BAR agonist, alone or in combination with addnl. agents, for the treatment of a B-cell proliferative disorder.



10/598,520


IT 53296-10-9, CV 1808  
RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL  
(Biological study); USES (Uses)  
( $\beta$ -Adrenergic receptor agonists for treatment of B-cell  
proliferative disorders, and use with other agents)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



L6 ANSWER 4 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 2009:1496452 CAPLUS  
DN 153:34972  
TI Adenosine: roles of different receptor subtypes in mediating histamine  
release from human and rodent mast cells  
AU Yip, K. H.; Wong, L. L.; Lau, H. Y. A.  
CS Department of Pharmacology, Faculty of Medicine, Chinese University of  
Hong Kong, Hong Kong SAR, Peop. Rep. China  
SO Inflammation Research (2009), 58(Suppl. 1), S17-S19  
CODEN: INREFB; ISSN: 1023-3830  
PB Birkhaeuser Verlag  
DT Journal  
LA English  
AB The effects of adenosine and adenosine receptor agonists on basal and  
anti-IgE induced histamine release from rat (RPMC) were compared with  
human mast cells (HCMC). Adenosine and its analogs alone did not initiate  
histamine release from RPMC and HCMC. However, adenosine could modulate  
IgE-dependent mediator release from both mast cell types, but with totally  
opposite predominant actions. Adenosine ( $10^{-5}$ - $10^{-3}$  M) produced a  
dose-dependent potentiating effect on anti-IgE induced histamine release  
in RPMC but a predominantly inhibitory action in HCMC. When adenosine was  
added simultaneously with anti-IgE to RPMC, an inhibitory tendency was  
observed at concns. below  $10^{-5}$  M, while the potentiating effect observed at  
higher concns. remained. Contrastingly, when added to HCMC simultaneously  
with anti-IgE, adenosine produced only dose-dependent inhibition but  
slight potentiation between  $10^{-9}$  to  $10^{-7}$  M was observed before the strong  
inhibition above  $10^{-6}$  M when adenosine was incubated with HCMC 10 min  
before anti-IgE challenge.  
IT 53296-10-9, 2-Phenylamino-adenosine  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES  
(Uses)  
(adenosine receptor subtypes in mediating histamine release from human  
and rodent mast cells)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.




McIntosh

RE.CNT 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 5 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 2009:1368423 CAPLUS  
DN 152:51216  
TI Drug Effects Viewed from a Signal Transduction Network Perspective  
AU Fliri, Anton F.; Loging, William T.; Volkmann, Robert A.  
CS Pfizer Global Research and Development, Groton, CT, 06340, USA  
SO Journal of Medicinal Chemistry (2009), 52(24), 8038-8046  
CODEN: JMCMAR; ISSN: 0022-2623  
PB American Chemical Society  
DT Journal  
LA English  
AB Understanding how drugs affect cellular network structures and how resulting signals are translated into drug effects holds the key to the discovery of medicines. Herein we examine this cause-effect relationship by determining protein network structures associated with the generation of specific in vivo drug-effect patterns. Medicines having similar in vivo pharmacol. have been identified by a comparison of drug-effect profiles of 1320 medicines. Protein network positions reached by these medicines were ascertained by examining the coinvestigation frequency of these medicines and 1179 protein network constituents in millions of scientific investigations. Interestingly, medicine assocns. obtained by comparing by drug-effect profiles mirror those obtained by comparing drug-protein coinvestigation frequency profiles, demonstrating that these drug-protein reachability profiles are relevant to in vivo pharmacol. By using protein assocns. obtained in these investigations and independent, curated protein interaction information, drug-mediated protein network topol. models can be constructed. These protein network topol. models reveal that drugs having similar pharmacol. profiles reach similar discrete positions in cellular protein network systems and provide a network view of medicine cause-effect relationships.  
IT 53296-10-9, 2-Phenylaminoadenosine  
RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(drug effects viewed from a signal transduction network perspective)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)  
RE.CNT 36 THERE ARE 36 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 6 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 2009:1180279 CAPLUS  
DN 152:6936  
TI CkI $\alpha$ / $\delta$ -dependent phosphorylation is a temperature-insensitive, period-determining process in the mammalian circadian clock  
AU Isojima, Yasushi; Nakajima, Masato; Ukai, Hideki; Fujishima, Hiroshi; Yamada, Rikuhiko G.; Masumoto, Koh-Hei; Kiuchi, Reiko; Ishida, Mayumi; Ukai-Tadenuma, Maki; Minami, Yoichi; Kito, Ryotaku; Nakao, Kazuki; Kishimoto, Wataru; Yoo, Seung-Hee; Shimomura, Kazuhiro; Takao, Toshifumi; Takano, Atsuko; Kojima, Toshio; Nagai, Katsuya; Sakaki, Yoshiyuki; Takahashi, Joseph S.; Ueda, Hiroki R.  
CS Comparative Systems Biology Team, Genomic Science Center, RIKEN, 1-7-22, Suehiro-cho, Tsurumi, Yokohama, 230-0045, Japan  
SO Proceedings of the National Academy of Sciences of the United States of

America (2009), 106(37), 15744-15749, S15744/1-S15744/74  
 CODEN: PNASA6; ISSN: 0027-8424

PB National Academy of Sciences

DT Journal

LA English

AB A striking feature of the circadian clock is its flexible yet robust response to various environmental conditions. To analyze the biochem. processes underlying this flexible-yet-robust characteristic, we examined the effects of 1260 pharmacol. active compds. in mouse and human clock cell lines. Compds. that markedly (>10 s.d.) lengthened the period in both cell lines, also lengthened it in central clock tissues and peripheral clock cells. Most compds. inhibited casein kinase 1 $\alpha$  (CK1 $\alpha$ ) or CK1 $\delta$  phosphorylation of the PER2 protein. Manipulation of CK1 $\alpha$ / $\delta$ -dependent phosphorylation by these compds. lengthened the period of the mammalian clock from circadian (24 h) to circadian (48 h), revealing its high sensitivity to chemical perturbation. The degradation rate of PER2, which is regulated by CK1 $\alpha$ / $\delta$ -dependent phosphorylation, was temperature-insensitive in living clock cells, yet sensitive to chemical perturbations. This temperature-insensitivity was preserved in the CK1 $\alpha$ / $\delta$ -dependent phosphorylation of a synthetic peptide in vitro. Thus, CK1 $\alpha$ / $\delta$ -dependent phosphorylation is likely a temperature-insensitive period-determining process in the mammalian circadian clock.

IT 53296-10-9, CV-1808


RL: BSU (Biological study, unclassified); PAC (Pharmacological activity);  
 BIOL (Biological study)

(CK1 $\alpha$ / $\delta$ -dependent phosphorylation is temperature-insensitive,  
 period-determining process in mammalian circadian clock)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 11 THERE ARE 11 CAPLUS RECORDS THAT CITE THIS RECORD (11 CITINGS)

RE.CNT 93 THERE ARE 93 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 7 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 2009:1173082 CAPLUS

DN 151:542900

TI Structure-based discovery of low molecular weight compounds that stimulate neurite outgrowth and substitute for nerve growth factor

AU Williams, Britney; Dwyer, Donard S.

CS Departments of Psychiatry, and Pharmacology, Toxicology and Neuroscience,  
 Louisiana State University Health Sciences Center, Shreveport, LA, USA

SO Journal of Neurochemistry (2009), 110(6), 1876-1884

CODEN: JONRA9; ISSN: 0022-3042

PB Wiley-Blackwell

DT Journal

LA English

AB Olanzapine, an atypical antipsychotic drug, was previously shown to protect neuronal cells against nutrient deprivation and to enhance neurite outgrowth. In an effort to identify small mols. with greater potency, the structure of olanzapine was used as a template to search com. available chemical inventories for compds. with similar features. These compds. were evaluated for their ability to protect cells against glutamine deprivation and low-serum conditions. Pos. compds., 'hits' from initial screening, were then tested for stimulation of neurite outgrowth, alone and in combination with suboptimum concns. of nerve growth factor (NGF). Numerous neuroprotective compds. (mw < 550 Da) were identified that significantly stimulated neurite outgrowth in PC12 cells. These included

4', 6'-diamidino-2-phenylindole, a nuclear stain; staurosporine, an antibiotic and kinase inhibitor; and 2-phenylamino-adenosine, an adenosine analog. The small mols. were comparable with NGF, and in fact, replaced NGF in outgrowth assays. Pharmacophore anal. of the hits led to the design and synthesis of an active compound, LSU-D84, which represented an initial lead for drug discovery efforts.


IT 53296-10-9, LSU 165

RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(structure-based discovery of low mol. weight compds. that stimulate neurite outgrowth and substitute for nerve growth factor)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



RE.CNT 33 THERE ARE 33 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 8 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 2009:740234 CAPLUS

DN 151:70285

TI Compositions and methods coactivating both A1 and A2A adenosine receptors for the treatment and prevention of cardiovascular diseases

IN Feldman, Arthur; Chan, Tung

PA Thomas Jefferson University, USA

SO PCT Int. Appl., 127pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2009076580	A2	20090618	WO 2008-US86528	20081212
WO 2009076580	A3	20090820		
W:	AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW			
RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA			
US 20100272711	A1	20101028	US 2010-747147	20100706
PRAI US 2007-13057P	P	20071212		
WO 2008-US86528	W	20081212		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB The present invention is directed to a pharmaceutical composition, and methods of use thereof, comprising at least one agent which target multiple adenosine receptors (AR) simultaneously in a stoichiometric relationship (i.e. each AR receptor is targeted to an equal extent). Aspects of the present invention relate to pharmaceutical compns., and uses thereof, comprising at least one agent which co-activates an A1-adenosine receptor (A1-AR) and an A2A-adenosine receptor (A2A-AR) or a combination of at least one agent which activates an A1-AR and at least one agent which activates an A2A-AR, where both the A1-AR and A2A-AR are activated in a stoichiometric relationship such that the level of biol. activation of

A1-AR is approx. the same level of biol. activation of A2A-AR. Other aspects of the present invention relate to methods for the therapeutic and prophylactic treatment of cardiac dysfunction in a subject having or at risk of having a cardiac dysfunction, for example, but not limited to, for the treatment of a subject with myocardial infarction, such as acute myocardial infarction, coronary ischemia or congestive heart failure and other cardiac dysfunctions. Long term or chronic administration of agonists which activate only the A1-AR or alternatively only the A2A-AR results in deleterious effects on cardiac function. If both the A1-AR and the A2A-AR are co-activated substantially simultaneously, the cardiac function was unexpectedly not compromised. Thus, use of at least one agent which co-activates both the A1-AR and the A2A-AR, or a combination of at least one or more agents which activates the A1-AR and at least one or more agents which activate the A2A-AR is useful to mediate cardioprotective effect.


IT 53296-10-9, CV1808

RL: BSU (Biological study, unclassified); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(or analogs or derivs. or salts thereof, as agent activating adenosine receptor A1; compns. and methods coactivating both A1 and A2A adenosine receptors for treatment and prevention of cardiovascular diseases)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

L6 ANSWER 9 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 2009:710118 CAPLUS

DN 151:49344

TI Combination of medical and physical cooling treatment of ischemic effects

IN Gotfredsen, Jacob; Weber, Uno Jakob

PA Neurokey A/S, Den.

SO PCT Int. Appl., 142pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 9

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2009071096	A2	20090611	WO 2008-DK50294	20081205
	WO 2009071096	A3	20100107		
	W:	AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW			
	RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA			
PRAI	DK 2007-1742	A	20071205		
	DK 2007-1744	A	20071205		
	DK 2008-1104	A	20080815		
	DK 2008-1105	A	20080815		

OS MARPAT 151:49344

AB The present invention relates to the induction of hypothermia in humans in

a predictable and dose responsive fashion by use of combination of phys./mech. hypothermia therapy and a pharmaceutical composition comprising at least one compound selected among a (1) vanilloid receptor agonists, capsaicinoids or capsaicinoid-like agonists reaching and binding to vanilloid receptors, and (2) cannabinoids or cannabimimetic agonists reaching and binding to cannabinoid receptors, and (3) adenosine receptor agonists, and (4) neurotensin receptor agonists, and (5) thyroxine derivs., and (6) cytochrome c oxidase inhibitors and (7) oxygen tension reducers thereby inducing hypothermia, thus benefiting patients suffering from illnesses characterized by tissue anoxia.

IT 53296-10-9, CV-1808


RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(as adenosine receptor agonist; combination of medical and phys. cooling treatment of ischemic effects)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



L6 ANSWER 10 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 2009:703623 CAPLUS

DN 151:49342

TI Combination treatment of ischemic effects

IN Gotfredsen, Jacob; Weber, Uno Jakob

PA Neurokey A/S, Den.

SO PCT Int. Appl., 131pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 9

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2009071094	A2	20090611	WO 2008-DK50292	20081205
	WO 2009071094	A3	20090806		

W: AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW

RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA

PRAI	DK 2007-1742	A	20071205
	DK 2008-1079	A	20080807
	DK 2008-1105	A	20080815

OS MARPAT 151:49342

AB The present invention relates to the induction of hypothermia in humans, male and female, at any age, in a predictable and dose responsive fashion by use of a pharmaceutical composition comprising a combination of two or more compds. selected among (1) vanilloid receptor agonists, capsaicinoids or capsaicinoid-like agonists reaching and binding to vanilloid receptors, and (2) cannabinoids or cannabimimetic agonists reaching and binding to cannabinoid receptors, and (3) adenosine receptor agonists, and (4) neurotensin receptor agonists, and (5) thyroxine derivs., and (6) cytochrome c inhibitors, and (7) oxygen tension reducers, with the proviso

10/598,520

that if the first compound is (1) then the second is not (2), thereby inducing hypothermia, thus benefiting patients suffering from illnesses characterized by tissue anoxia.

IT 53296-10-9, CV-1808


RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(as adenosine receptor agonist; combination treatment of ischemic effects using hypothermia inducing agents)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



L6 ANSWER 11 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 2009:343325 CAPLUS

DN 151:212164

TI Activation of adenosine A2A receptor impairs memory acquisition but not consolidation or retrieval phases

AU Kim, Dong Hyun; Ryu, Jong Hoon

CS Department of Life and Nanopharmaceutical Sciences, Kyung Hee East-West Pharmaceutical Research Institute, College of Pharmacy, Kyung Hee University, Hoeki-dong, Dongdaemoon-Ku, Seoul, 130-701, S. Korea

SO Biomolecules & Therapeutics (2008), 16(4), 320-327

CODEN: BTIHA3; ISSN: 1976-9148

PB Korean Society of Applied Pharmacology

DT Journal

LA English

AB Several lines of evidence indicate that adenosine A2A agonist disrupts spatial working memory. However, it is unclear which stages of learning and memory are affected by the stimulation of adenosine A2A receptor. To clarify these points, we employed CV-1808 as adenosine A2A agonist and investigated its effects on acquisition, consolidation, and retrieval phases of learning and memory using passive avoidance and the Morris water maze tasks. During the acquisition phase, CV-1808 (2-phenylaminoadenosine, 1 and 2 mg/kg, i.p.) decreased the latency time in passive avoidance task and the mean savings in the Morris water maze task, resp. During the consolidation and retrieval phase tests, CV-1808 did not exhibited any effects on latency time in passive avoidance task and the mean savings in the Morris water maze task. These results suggest that CV-1808 as an adenosine A2A agonist impairs memory acquisition but not consolidation or retrieval.

IT 53296-10-9, CV-1808


RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(activation of adenosine A2A receptor by CV-1808 impaired acquisition but not consolidation or retrieval phase of memory and learning in mouse)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)  
 RE.CNT 60 THERE ARE 60 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 12 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
 AN 2009:86451 CAPLUS  
 DN 150:160095  
 TI Use of adenosine A2A receptor agonists and phosphodiesterase (PDE)  
 inhibitors for the treatment of B-cell proliferative disorders, and  
 combinations with other agents  
 IN Rickles, Richard; Lee, Margaret S.  
 PA CombinatoRx, Incorporated, USA  
 SO PCT Int. Appl., 70 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI WO 2009011893	A2	20090122	WO 2008-US8758	20080717
WO 2009011893	A3	20090319		
W:	AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW			
RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA			
AU 2008276451	A1	20090122	AU 2008-276451	20080717
CA 2694983	A1	20090122	CA 2008-2694983	20080717
US 20090053168	A1	20090226	US 2008-175219	20080717
EP 2178369	A2	20100428	EP 2008-780231	20080717
R:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GE, GR, HR, HU, IE, IS, IT, LI, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR, AL, BA, MK, RS			
PRAI US 2007-950307P	P	20070717		
US 2007-965587P	P	20070821		
WO 2008-US8758	W	20080717		

# ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT


AB The invention provides compns. and methods for the treatment of B-cell proliferative disorders that employ an A2A receptor agonist or one or more PDE inhibitors. The methods and compns. may further include an antiproliferative compound

IT 53296-10-9, CV 1808  
 RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (adenosine A2A receptor agonists and phosphodiesterase inhibitors for treatment of B-cell proliferative disorders, and combinations with other agents)

RN 53296-10-9 CAPLUS  
 CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.





OSC.G 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)

L6 ANSWER 13 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
 AN 2009:83374 CAPLUS  
 DN 150:160094  
 TI Combinations for the treatment of B-cell proliferative disorders  
 IN Rickles, Richard; Pierce, Laura; Lee, Margaret S.  
 PA Combinatorx, Incorporated, USA  
 SO PCT Int. Appl., 79pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI WO 2009011897	A1	20090122	WO 2008-US8764	20080717
W:	AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW			
RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
AU 2008276455	A1	20090122	AU 2008-276455	20080717
CA 2694987	A1	20090122	CA 2008-2694987	20080717
US 20090047243	A1	20090219	US 2008-175121	20080717
EP 2178370	A1	20100428	EP 2008-780237	20080717
R:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LI, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR, AL, BA, MK, RS			
PRAI US 2007-959877P	P	20070717		
US 2007-965595P	P	20070821		
WO 2008-US8764	W	20080717		

# ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT


AB The invention features compns. and methods employing combinations of an A2A receptor agonist and a PDE (phosphodiesterase) inhibitor for the treatment of a B-cell proliferative disorder, e g, multiple myeloma. In at least one embodiment, the compns. of the invention comprise a PDE inhibitor active against at least two of PDE 2, 3,4, and 7. In at least one embodiment, the compns. of the invention comprises further administering an antiproliferative compound

IT 53296-10-9, CV 1808  
 RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (combinations for treatment of B-cell proliferative disorders using PDE inhibitors and A2A receptor agonists and antiproliferative compds.)

RN 53296-10-9 CAPLUS  
 CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.


10/598,520



OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)  
RE.CNT 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 14 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 2008:1383562 CAPLUS  
DN 149:555078  
TI The Stille reaction  
AU Farina, Vittorio; Krishnamurthy, Venkat; Scott, William J.  
CS Boehringer Ingelheim Pharmaceuticals, Ridgefield, CT, USA  
SO Organic Reactions (Hoboken, NJ, United States) (1997), 50, No pp. given  
CODEN: ORHNBA  
URL: <http://www3.interscience.wiley.com/cgi-bin/mrwhome/107610747/HOME>  
PB John Wiley & Sons, Inc.  
DT Journal; General Review; (online computer file)  
LA English  
OS CASREACT 149:555078  
AB A review of the article The Stille reaction.  
IT 79936-11-1P  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(The Stille Reaction)  
RN 79936-11-1 CAPLUS  
CN Adenosine, 2-cyano- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




L6 ANSWER 15 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 2008:991314 CAPLUS  
DN 149:200258  
TI Cyanotributylstannane  
AU Tanaka, Masato; Sakakura, Toshiyasu  
CS Japan  
SO e-EROS Encyclopedia of Reagents for Organic Synthesis (2001), No pp. given  
Publisher: John Wiley & Sons, Ltd., Chichester, UK.  
CODEN: 69KUHI  
URL: <http://www3.interscience.wiley.com/cgi-bin/mrwhome/104554785/HOME>  
DT Conference; General Review; (online computer file)  
LA English  
OS CASREACT 149:200258  
AB A review of the article Cyanotributylstannane.  
IT 79936-11-1P  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(Cyanotributylstannane)  
RN 79936-11-1 CAPLUS  
CN Adenosine, 2-cyano- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

McIntosh

10/598,520



L6 ANSWER 16 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 2008:493012 CAPLUS  
DN 148:509885  
TI Compositions and methods for treating neurological disorders or damage  
IN Diamandis, Phedias; Tyers, Mike; Dirks, Peter B.  
PA Can.  
SO Can. Pat. Appl., 3pp.  
CODEN: CPXXEB  
DT Patent  
LA English  
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	CA 2606658	A1	20080413	CA 2007-2606658	20071012
	US 20090076019	A1	20090319	US 2007-871562	20071012
PRAI	US 2006-851615P	P	20061013		


ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB The invention relates to a clonogenic neurosphere assay to carry out high throughput screens (HTS) to identify potent and/or selective modulators of proliferation, differentiation and/or renewal of neural precursor cells, neural progenitor cells and/or self-renewing and multipotent neural stem cells (NSCs). The invention also relates to compns. comprising the identified modulators and methods of using the modulators and compns., in particular to treat neurol. disorders (e.g. brain or CNS cancer) or damage.

IT 53296-10-9, 2-Phenylaminoadenosine  
RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(screening for compns. and methods for treating neurol. disorders or damage with modulators of neural stem cells)

RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

L6 ANSWER 17 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 2008:11519 CAPLUS  
DN 148:100840  
TI Preparation of nucleosides as therapeutic compounds for the treatment of pain and inflammation  
IN Higginbottom, Michael; Savory, Edward Daniel; Brown, Giles Albert; Horgan, Viet-Anh Anne; Chapman, Emma Jane  
PA Biovitrum AB, Swed.  
SO PCT Int. Appl., 70 pp.  
CODEN: PIXXD2  
DT Patent

McIntosh

10/598,520


LA English  
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2008000745	A2	20080103	WO 2007-EP56378	20070626
	WO 2008000745	A3	20080417		
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW			
	RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, CG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA			
	AU 2007263728	A1	20080103	AU 2007-263728	20070626
	CA 2654375	A1	20080103	CA 2007-2654375	20070626
	US 20080076776	A1	20080327	US 2007-823377	20070626
	US 7807685	B2	20101005		
	EP 2081946	A2	20090729	EP 2007-786850	20070626
	R:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR, AL, BA, HR, MK, RS			
	JP 2009541438	T	20091126	JP 2009-517175	20070626
	IN 2008KN04769	A	20090313	IN 2008-KN4769	20081125
	CN 101479289	A	20090708	CN 2007-80023745	20081224
PRAI	SE 2006-1398	A	20060627		
	US 2006-837146P	P	20060811		
	WO 2007-EP56378	W	20070626		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OS CASREACT 148:100840; MARPAT 148:100840

GI



I

AB Nucleosides I, when X = Y = Z = OH, R<sub>1</sub> is OCH<sub>2</sub>CF<sub>2</sub>CF<sub>3</sub>, phenoxy (substituted with 3-(4- trifluoromethylphenyl), 3,4-dichloro, (3-trifluoromethyl,4-fluoro), (3-trifluoromethyl,4- chloro), (3-chloro, 4-cyano), or 3,5-bis(trifluoromethyl)), 1-piperazinyl(4-(3,4- dichlorophenyl)), Ph (substituted with 3,4-dichloro, 3,5-difluoro, 3,5- bis(trifluoromethyl) or 3,4,5-trifluoro) or 2-benzofuranyl; or when X = Y = OH and Z = OMe, R<sub>1</sub> is OCH<sub>3</sub>, OCH<sub>2</sub>CHF<sub>2</sub>, OCH<sub>2</sub>cyclopentyl, O-(2,5- difluorophenyl) or (S)-sec-butylamino; or when X = H and Y = Z = OH, R<sub>1</sub> is n-hexylamino or cyclopentylamino; or when (IV) X = Z = OH and Y = H, R<sub>1</sub> is cyclopentylamino; were prepared for the treatment of pain and inflammation. Thus, nucleoside I (X = Y = Z = OH, R<sub>1</sub> = CH<sub>2</sub>CF<sub>2</sub>CF<sub>3</sub>) was prepared and tested for the treatment of pain and inflammation.

IT 1000003-47-3P 1000003-48-4P 1000003-49-5P  
1000003-50-8P 1000003-51-9P 1000003-52-0P  
1000003-53-1P 1000003-57-5P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of nucleosides as therapeutic compds. for treatment of pain and inflammation)


McIntosh

10/598,520

RN 1000003-47-3 CAPLUS

CN Adenosine, 2-(2,2,3,3,3-pentafluoropropoxy)- (CA INDEX NAME)


Absolute stereochemistry.



RN 1000003-48-4 CAPLUS

CN Adenosine, 2-[[4'-(trifluoromethyl)[1,1'-biphenyl]-3-yl]oxy]- (CA INDEX NAME)


Absolute stereochemistry.



RN 1000003-49-5 CAPLUS

CN Adenosine, 2-(3,4-dichlorophenoxy)- (CA INDEX NAME)


Absolute stereochemistry.



RN 1000003-50-8 CAPLUS

CN Adenosine, 2-[4-fluoro-3-(trifluoromethyl)phenoxy]- (CA INDEX NAME)

Absolute stereochemistry.




McIntosh

10/598,520


RN 1000003-51-9 CAPLUS  
CN Adenosine, 2-[4-chloro-3-(trifluoromethyl)phenoxy]- (CA INDEX NAME)

Absolute stereochemistry.




RN 1000003-52-0 CAPLUS  
CN Adenosine, 2-(3-chloro-4-cyanophenoxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 1000003-53-1 CAPLUS  
CN Adenosine, 2-[3,5-bis(trifluoromethyl)phenoxy]- (CA INDEX NAME)

Absolute stereochemistry.



RN 1000003-57-5 CAPLUS  
CN Adenosine, 2-[3,5-bis(trifluoromethyl)phenyl]- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

L6 ANSWER 18 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

McIntosh

10/598,520

AN 2008:9613 CAPLUS  
DN 148:106191  
TI 2',3'-Methylidene acetal adenosine prodrugs of improved oral absorption  
and their use as therapeutic analgesic or antiinflammatory compounds  
IN Savory, Edward Daniel  
PA Biovitrum AB, Swed.  
SO PCT Int. Appl., 64pp.  
CODEN: PIXXD2

DT Patent  
LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2008000743	A2	20080103	WO 2007-EP56375	20070626
	WO 2008000743	A3	20080221		
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW			
	RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA			
	AU 2007263726	A1	20080103	AU 2007-263726	20070626
	CA 2657973	A1	20080103	CA 2007-2657973	20070626
	US 20080027081	A1	20080131	US 2007-823335	20070626
	EP 2066685	A2	20090610	EP 2007-765638	20070626
	R:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR, AL, BA, HR, MK, RS			
	JP 2009541436	T	20091126	JP 2009-517173	20070626
	IN 2008KN04767	A	20090313	IN 2008-KN4767	20081125
	CN 101479290	A	20090708	CN 2007-80024647	20081229
PRAI	SE 2006-1396	A	20060627		
	US 2006-837308P	P	20060811		
	WO 2007-EP56375	W	20070626		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OS CASREACT 148:106191; MARPAT 148:106191

AB The invention relates to a method of improving oral drug absorption of adenosine analogs by the use of 2',3'-methylidene acetal adenosine pro-drugs and to the use of these pro-drugs as medicaments. The invention further relates to compds. that are prodrugs of adenosine receptor agonists, and to their use as therapeutic compds., in particular as analgesic or anti-inflammatory compds., or as disease modifying antirheumatic drugs (DMARDs), and to methods of preventing, treating or ameliorating pain or inflammation using these compds. Thus, for a range of five 2-substituted adenosines of the current invention, the oral bioavailability in rats was found to increase on average from 19% to 53% and the oral half-life from 1.3 h to 3.2 h by employing a 2',3'-methylidene acetal prodrug strategy.

IT 864061-82-5 864061-84-7 1000003-48-4  
1000003-53-1


RL: BCP (Biochemical process); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)  
(2',3'-Methylidene acetal adenosine prodrugs of improved oral absorption and their use as therapeutic analgesic or antiinflammatory compds.)

RN 864061-82-5 CAPLUS

CN Adenosine, 2-(2,2-difluoroethoxy)- (CA INDEX NAME)


Absolute stereochemistry.

10/598,520




RN 864061-84-7 CAPLUS  
CN Adenosine, 2-(2,5-difluorophenoxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 1000003-48-4 CAPLUS  
CN Adenosine, 2-[[4'-(trifluoromethyl)[1,1'-biphenyl]-3-yl]oxy]- (CA INDEX NAME)

Absolute stereochemistry.



RN 1000003-53-1 CAPLUS  
CN Adenosine, 2-[3,5-bis(trifluoromethyl)phenoxy]- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)

L6 ANSWER 19 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 2007:1320679 CAPLUS  
DN 148:135926

McIntosh



TI Effect of adenosine agonists on the proliferation and differentiation of chick embryo fibroblasts in three dimensional reconstituted tissue constructs

AU Malihi, Golrokh; Elson, Elliot; Mascarenhas, Francesca

CS Department of Biochemistry/Molecular Biophysics, School of medicine, Washington University, St. Louis, MO, 63110-8231, USA

SO Iranian Journal of Pharmacology & Therapeutics (2006), 5(2), 151-157  
CODEN: IJPTDG; ISSN: 1735-2657  
URL: <http://ijpt.iuums.ac.ir/index.php/ijpt/article/view/060502151/237>

PB Razi Institute for Drug Research, Iran University of Medical Sciences and Health Services

DT Journal; (online computer file)

LA English


AB Previous studies indicate that organ fibroblasts play an important role in wound healing, collagen production, remodeling processes and pathogenesis of progressive heart, lung, renal and hepatic fibrotic diseases. Several studies suggest a possible inhibitory role for adenosine in the regulation of fibroblast proliferation. The effect of adenosine A2 agonists on proliferation and differentiation of chick embryo skin/muscle fibroblasts was studied in collagen-based 3-dimensional tissue constructs and also in plated monolayer cells. Materials and Methods: Chick embryo primary fibroblasts were plated in sep. groups and were synchronized by growth arrest before stimulation by different doses of adenosine, and A2 receptor agonists, CV1808, NECA and an A2 receptor antagonist, CGS15943, and control, in the presence of serum or serum free medium. The cell counts for each treatment of monolayer fibroblasts were compared to determine fibroblast proliferation. Western blot anal., immunostaining and myofibroblast size measurements were conducted to measure the effect of adenosine on the fibroblast differentiation into myofibroblasts. Cell proliferation was also gauged with DNA assays in the 3-D constructs. Results: Adenosine agonists at low doses significantly reduced fibroblast proliferation in monolayer and 3-D cell culture in the presence of 5% Fetal Calf Serum (FCS) demonstrating a potential antifibrotic activity possibly by activation of the A2B receptor. Western blot anal. and immunostaining of cells revealed no significant inhibition of the expression of  $\alpha$ -smooth muscle actin on a per cell basis by adenosine agonists. Cell size measurements indicated increased nos. of smaller fibroblasts suggesting that adenosine may inhibit the conversion of fibroblasts to myofibroblasts. Conclusion: This study suggests that agents that increase tissue cAMP levels may be of beneficial therapeutic value in organ tissue fibrosis.

IT 53296-10-9, CV1808  
RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(adenosine-5'-N-ethylcarboxamide or CV1808 was associated with decrease in proliferation, differentiation and size in chick embryo fibroblast and can be useful in treatment of fibrosis)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



RE.CNT 28 THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 20 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 2007:873285 CAPLUS

DN 147:242695

TI Compounds useful as agonists of a2a adenosine receptors, cosmetic skin whitening compositions with a2a agonists and a method for using the same

IN Nip, John Chun-Sing; Bosko, Carol Annette; Rosa, Jose Guillermo;

10/598,520

Harichian, Bijan; Santana, Isabel Cristina  
PA Unilever PLC, USA  
SO U.S. Pat. Appl. Publ., 8pp.  
CODEN: USXXCO  
DT Patent  
LA English  
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 20070183995	A1	20070809	US 2006-350658	20060209
	AU 2007214068	A1	20070816	AU 2007-214068	20070125
	WO 2007090553	A2	20070816	WO 2007-EP847	20070125
	WO 2007090553	A3	20071101		
	W: AE, AG, AI, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW				
	RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA				
	ZA 2008006447	A	20091230	ZA 2008-6447	20070125
	AR 59370	A1	20080326	AR 2007-100527	20070208
	IN 2008MN01691	A	20081226	IN 2008-MN1691	20080807
	CN 101378725	A	20090304	CN 2007-80004724	20080807
	MX 2008010208	A	20081031	MX 2008-10208	20080808
	KR 2008108418	A	20081215	KR 2008-7019553	20080808
PRAI	US 2006-350658	A	20060209		
	WO 2007-EP847	W	20070125		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OS MARPAT 147:242695

AB Comps. useful as agonists of A2A adenosine receptors are described. Also described is a cosmetically acceptable composition having an agonists of A2A adenosine receptors where the composition is suitable to apply to human skin to reduce the effects of melanin, resulting in skin whitening. Thus, solns. of the A2A adenosine receptor agonists 2-para(2-carboxyethyl)phenethylamino-5'-N-Et carboxamido adenosine and phenylaminoadenosinehaving, of a final concentration of 3  $\mu$ M were prepared from a 10 mM DMSO stock solution and dosed on human skin equivalent (Melanoderm from Mattek). The colorimetric results showed that compns. with an agonist of A2A adenosine receptors could result in skin lightening.

IT 53296-10-9


RL: BSU (Biological study, unclassified); COS (Cosmetic use); BIOL (Biological study); USES (Uses)

(comps. useful as agonists of a2a adenosine receptors, cosmetic skin whitening compns. with a2a agonists and a method for using the same)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

L6 ANSWER 21 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 2007:870078 CAPLUS

DN 147:340476

TI Determination of adenosine effects and adenosine receptors in murine

McIntosh

corpus cavernosum

AU Tostes, Rita C.; Giachini, Fernanda R. C.; Carneiro, Fernando S.; Leite, Romulo; Inscho, Edward W.; Webb, R. Clinton

CS Department of Pharmacology, Institute of Biomedical Sciences, University of Sao Paulo, Sao Paulo, Brazil

SO Journal of Pharmacology and Experimental Therapeutics (2007), 322(2), 678-685  
CODEN: JPETAB; ISSN: 0022-3565

PB American Society for Pharmacology and Experimental Therapeutics

DT Journal

LA English


AB This study tested the hypothesis that adenosine, in murine corpora cavernosa, produces direct relaxation of smooth muscle cells and inhibition of contractile responses mediated by sympathetic nerve stimulation. Penes were excised from anesthetized male C57BL/6 mice, dissected, and cavernosal strips were mounted to record isometric force. Adenosine, 2-chloro-adenosine (stable analog of adenosine), and 2-phenylaminoadenosine (CV1808) (A2A/A2B agonist) produced concentration-dependent relaxations of phenylephrine-contracted tissues. Relaxation to 2-chloroadenosine was inhibited, in a concentration-dependent manner, by 2-(2-furanyl)-7-(2-phenylethyl)-7H-pyrazolo[4,3-e][1,2,4]triazolo [1,5-c]pyrimidin-5-amine (SCH58261; A2A antagonist; 10-9-10-6 M) and N-(4-acetylphenyl)-2-[4-(2,3,6,7-tetrahydro-2,6-dioxo-1,3-dipropyl-1H-purin-8-yl)phenoxy]acetamida (MRS1706; A2B antagonist; 10-8-10-6 M). The combination of both antagonists abrogated 2-chloroadenosine-induced relaxation. Elec. field stimulation (EFS; 1-32 Hz) of adrenergic nerves produced frequency-dependent contractions that were inhibited by compds. that increase adenosine levels, such as 5'-iodotubercidin (adenosine kinase inhibitor), erythro-9-(2-hydroxy-3-nonyl)adenine (adenosine deaminase inhibitor), and dipyridamole (inhibitor of adenosine transport). The adenosine A1 receptor agonist N6-cyclopentyladenosine (C8031) right-shifted contractile responses to EFS, with a significant inhibitory effect at 10-6 M. Blockade of adenosine A1 receptors with 8-cyclopentyl-1,3-dipropylaxanthine (C101) (10-7 M) enhanced contractile responses to EFS and eliminated the inhibitory effects of 5'-iodotubercidin. Dipyridamole and 5'-iodotubercidin had no effect on adenosine-mediated relaxation. In summary, adenosine directly relaxes cavernosal smooth muscle cells, by the activation of A2A/A2B receptor subtypes. In addition, adenosine neg. modulates sympathetic neurotransmission, by A1 receptor subtype activation, in murine corpora cavernosa. Adenosine may subserve dual roles in modulating the physiol. mechanisms of erection in mice.

IT 53296-10-9, 2-Phenylaminoadenosine  
RL: BSU (Biological study, unclassified); BIOL (Biological study) (effects of adenosine and adenosine receptors in murine corpus cavernosum)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 9 THERE ARE 9 CAPLUS RECORDS THAT CITE THIS RECORD (10 CITINGS)

RE.CNT 39 THERE ARE 39 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 22 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 2007:859948 CAPLUS

DN 148:45665

TI Effect of adenosine agonists on the proliferation and differentiation of chick embryo fibroblasts in three dimensional reconstituted tissue

constructs

AU Malihi, Golrokh; Elson, Elliot; Mascarenhas, Francesca

CS Department of Biochemistry/Molecular Biophysics, School of Medicine, Washington University in St. Louis, St. Louis, MO, USA

SO Iranian Journal of Pharmacology & Therapeutics (2006), 5(2), 151-157

CODEN: IJPTDG; ISSN: 1735-2657

URL: <http://ijpt.iums.ac.ir/index.php/ijpt/article/view/060502151/237>

PB Razi Institute for Drug Research, Iran University of Medical Sciences and Health Services

DT Journal; (online computer file)

LA English

AB Previous studies indicate that organ fibroblasts play an important role in wound healing, collagen production, remodeling processes and pathogenesis of progressive heart, lung, renal and hepatic fibrotic diseases. Several studies suggest a possible inhibitory role for adenosine in the regulation of fibroblast proliferation. The effect of adenosine A2 agonists on proliferation and differentiation of chick embryo skin/muscle fibroblasts was studied in collagen-based 3-dimensional tissue constructs and also in plated monolayer cells. Materials and Methods: Chick embryo primary fibroblasts were plated in sep. groups and were synchronized by growth arrest before stimulation by different doses of adenosine, and A2 receptor agonists, CV1808, NECA and an A2 receptor antagonist, CGS15943, and control, in the presence of serum or serum free medium. The cell counts for each treatment of monolayer fibroblasts were compared to determine fibroblast proliferation. Western blot anal., immunostaining and myofibroblast size measurements were conducted to measure the effect of adenosine on the fibroblast differentiation into myofibroblasts. Cell proliferation was also gauged with DNA assays in the 3-D constructs. Results: Adenosine agonists at low doses significantly reduced fibroblast proliferation in monolayer and 3-D cell culture in the presence of 5% Fetal Calf Serum (FCS) demonstrating a potential antifibrotic activity possibly by activation of the A2B receptor. Western blot anal. and immunostaining of cells revealed no significant inhibition of the expression of  $\alpha$ -smooth muscle actin on a per cell basis by adenosine agonists. Cell size measurements indicated increased nos. of smaller fibroblasts suggesting that adenosine may inhibit the conversion of fibroblasts to myofibroblasts. Conclusion: This study suggests that agents that increase tissue cAMP levels may be of beneficial therapeutic value in organ tissue fibrosis.

IT 53296-10-9, CV1808


RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(adenosine A2 agonists like CV1808 increased cAMP production and inhibited proliferation of chick embryo fibroblast in 3-dimensional reconstituted tissue constructs)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



RE.CNT 28      THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 23 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 2007:705774 CAPLUS  
DN 147:110249  
TI Agents for treating neurodegenerative diseases  
IN Stockwell, Brent R.; Hoffstrom, Benjamin; Varma, Hemant  
PA USA  
SO U.S. Pat. Appl. Publ., 117pp., Cont.-in-part of US Ser. No. 498,110.  
CODEN: USXXCO

McIntosh

10/598,520

DT Patent  
LA English  
FAN.CNT 5

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 20070149543	A1	20070628	US 2006-612286	20061218
	US 20050032124	A1	20050210	US 2004-767591	20040129
	US 20070027164	A1	20070201	US 2006-349653	20060207
	US 20070078144	A1	20070405	US 2006-498110	20060802
PRAI	US 2003-443728P	P	20030129		
	US 2003-457401P	P	20030325		
	US 2003-467290P	P	20030502		
	US 2003-482688P	P	20030625		
	US 2003-496209P	P	20030819		
	US 2004-767591	B2	20040129		
	US 2004-837360	A2	20040430		
	US 2006-349653	A2	20060207		
	US 2006-498110	A2	20060802		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OS MARPAT 147:110249


AB The present invention relates to compds. effective in preventing neuronal cell death, which may be used in the treatment of neurodegenerative diseases. It is based, at least in part, on the discovery that particular compds. were effective in preventing neuronal death in model systems of Huntington's Disease.

IT 53296-10-9, 2-Phenylaminoadenosine  
RL: DMA (Drug mechanism of action); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(agents for treating neurodegenerative diseases)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



L6 ANSWER 24 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 2007:643780 CAPLUS

DN 147:78751

TI Cosmetic composition containing a nonphosphate compound based on adenosine

IN Rolland, Anne; Catroux, Philippe

PA L'Oreal, Fr.

SO Fr. Demande, 20pp.

CODEN: FRXXBL

DT Patent

LA French

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	FR 2894466	A1	20070615	FR 2005-53770	20051208
PRAI	FR 2005-53770		20051208		

AB A cosmetic composition contains a nonphosphate compound based on adenosine and at least a vehicle in a quantity higher than 3% in weight, compared to the total weight of the composition. The cosmetic is used for care of skin, more particularly wrinkled skin of the face. A cream contained adenosine 0.04, stearic acid 3.0, a mixture of glyceryl monostearate and polyethylene glycol stearate 2.5, PEG stearate 1.0, cyclopentasiloxane 10, silica 3.5, vegetable oils 7.0, synthetic oils 6.0, preservatives 1.2, siliogum 0.2, polyoxyethylene polydimethylsiloxane 1.0, Simugel-600 1.7, stearyl alc. 1, and water q.s. 100%.


IT 53296-10-9, 2-Phenylaminoadenosine  
RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)

McIntosh

10/598,520

(cosmetic composition containing nonphosphate compound based on adenosine)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)  
RE.CNT 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 25 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 2007:438825 CAPLUS  
DN 146:427844  
TI Cosmetic composition containing a non-phosphate compound based on  
adenosine and a polymer  
IN Catroux, Philippe; Rolland, Anne  
PA L'Oreal, Fr.  
SO PCT Int. Appl., 25pp.  
CODEN: PIXXD2  
DT Patent  
LA French  
FAN.CNT 1


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2007042679	A2	20070419	WO 2006-FR2302	20061012
	WO 2007042679	A3	20070607		
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW			
	RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA			
	FR 2892017	A1	20070420	FR 2005-53131	20051014
PRAI	FR 2005-53131	A	20051014		
	US 2005-731261P	P	20051031		

AB The invention concerns a cosmetic composition comprising in a physiologically acceptable medium: at least one non-phosphate compound based on adenosine and at least a polymer, said polymer being different from a copolymer comprising units derived from styrene and units derived from (meth)acrylate. The invention also concerns a cosmetic method for skin care, more particularly facial skin, in particular wrinkled skin which consists in applying a composition on said skin. A lotion contained Hostacerin AMPS 2.00, preservatives 0.85, adenosine 0.50, Hybridur 875 17.00, and water q.s. 100%.

IT 53296-10-9, 2-Phenylaminoadenosine  
RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)  
(cosmetic composition containing non-phosphate compound based on adenosine and polymer)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.

McIntosh



OSC.G 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD (3 CITINGS)

L6 ANSWER 26 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
 AN 2007:384430 CAPLUS  
 DN 146:372825  
 TI Agents for treating neurodegenerative diseases  
 IN Stockwell, Brent R.; Hoffstrom, Benjamin; Varma, Hemant  
 PA Columbia University, USA  
 SO U.S. Pat. Appl. Publ., 99 pp., Cont.-in-part of U.S. Ser. No. 349,653.  
 CODEN: USXXCO  
 DT Patent  
 LA English  
 FAN.CNT 5

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 20070078144	A1	20070405	US 2006-498110	20060802
	US 20050032124	A1	20050210	US 2004-767591	20040129
	US 20070027164	A1	20070201	US 2006-349653	20060207
	US 20070149543	A1	20070628	US 2006-612286	20061218
	WO 2008016659	A2	20080207	WO 2007-US17221	20070802
	WO 2008016659	A3	20081113		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW				
RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA				
PRAI	US 2003-443728P	P	20030129		
	US 2003-457401P	P	20030325		
	US 2003-467290P	P	20030502		
	US 2003-482688P	P	20030625		
	US 2003-496209P	P	20030819		
	US 2004-767591	B2	20040129		
	US 2004-837360	A2	20040430		
	US 2006-349653	A2	20060207		
	US 2006-498110	A2	20060802		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OS MARPAT 146:372825

AB The present invention relates to compds. effective in preventing neuronal cell death, which may be used in the treatment of neurodegenerative diseases. It is based, at least in part, on the discovery that particular compds. were effective in preventing neuronal death in model systems of Huntington's Disease.


IT 53296-10-9, 2-Phenylaminoadenosine

RL: DMA (Drug mechanism of action); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (agents for treating neurodegenerative diseases)

RN 53296-10-9 CAPLUS


CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



L6 ANSWER 27 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
 AN 2007:328553 CAPLUS  
 DN 146:501286  
 TI Structure-Activity Relationships of 2,N6,5'-Substituted Adenosine  
 Derivatives with Potent Activity at the A2B Adenosine Receptor  
 AU Adachi, Hayamitsu; Palaniappan, Krishnan K.; Ivanov, Andrei A.; Bergman,  
 Nathaniel; Gao, Zhan-Guo; Jacobson, Kenneth A.  
 CS Molecular Recognition Section, Laboratory of Bioorganic Chemistry,  
 National Institute of Diabetes and Digestive and Kidney Diseases, National  
 Institutes of Health, Bethesda, MD, 20892, USA  
 SO Journal of Medicinal Chemistry (2007), 50(8), 1810-1827  
 CODEN: JMCMAR; ISSN: 0022-2623  
 PB American Chemical Society  
 DT Journal  
 LA English  
 OS CASREACT 146:501286  
 AB 2, N6, and 5'-substituted adenosine derivs. were synthesized via  
 alkylation of 2-oxypurine nucleosides leading to 2-arylalkyl-ether derivs.  
 2-(3-(Indolyl)ethyl-oxy)adenosine (I) was examined in both binding and cAMP  
 assays and found to be a potent agonist of the human A2BAR.  
 Simplification, altered connectivity, and mimicking of the indole ring of  
 I failed to maintain A2BAR potency. Introduction of N6-Et or N6-guanidino  
 substitution, shown to favor A2BAR potency, failed to enhance potency in  
 the 2-(3-(indolyl)ethyloxy)adenosine series. Indole 5'''- or 6'''-halo  
 substitution was favored at the A2BAR, but a 5'-N-ethylcarboxamide did  
 not further enhance potency. 2-(3'''-(6'''-Bromoindolyl)ethyloxy)adenosine  
 (II) displayed an A2BAR EC50 value of 128 nM, i.e., more potent than the  
 parent I (299 nM) and similar to 5'-N-ethylcarboxamidoadenosine (140 nM).  
 Compound II was a full agonist at A2B and A2AARs and a low efficacy partial  
 agonist at A1 and A3ARs. Thus, we have identified and optimized  
 2-(2-arylethyl)oxo moieties in AR agonists that enhance A2BAR potency and  
 selectivity.  
 IT 131865-81-1P 145747-87-1P  
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); BIOL  
 (Biological study); PREP (Preparation)  
 (structure-activity relationships of 2,N6,5'-substituted adenosine  
 derivs. with potent activity at A2B adenosine receptor)  
 RN 131865-81-1 CAPLUS  
 CN Adenosine, 2-(4-phenylbutoxy)- (CA INDEX NAME)


Absolute stereochemistry.



RN 145747-87-1 CAPLUS  
 CN Adenosine, 2-[(3S)-3-phenylbutoxy]- (CA INDEX NAME)

Absolute stereochemistry.






OSC.G 15 THERE ARE 15 CAPLUS RECORDS THAT CITE THIS RECORD (15 CITINGS)  
 RE.CNT 46 THERE ARE 46 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 28 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
 AN 2007:245615 CAPLUS  
 DN 146:474750  
 TI Three-Dimensional Quantitative Structure-Activity Relationship of  
 Nucleosides Acting at the A3 Adenosine Receptor: Analysis of Binding and  
 Relative Efficacy  
 AU Kim, Soo-Kyung; Jacobson, Kenneth A.  
 CS Molecular Recognition Section Laboratory of Bioorganic Chemistry, National  
 Institute of Diabetes and Digestive and Kidney Diseases (NIDDK), National  
 Institutes of Health (NIH), Bethesda, MD, 20892, USA  
 SO Journal of Chemical Information and Modeling (2007), 47(3), 1225-1233  
 CODEN: JCISD8; ISSN: 1549-9596  
 PB American Chemical Society  
 DT Journal  
 LA English  
 AB The binding affinity and relative maximal efficacy of human A3 adenosine  
 receptor (AR) agonists were each subjected to ligand-based  
 three-dimensional quant. structure-activity relation anal. Comparative  
 mol. field anal. (CoMFA) and comparative mol. similarity indexes anal.  
 (CoMSIA) used as training sets a series of 91 structurally diverse  
 adenosine analogs with modifications at the N6 and C2 positions of the  
 adenine ring and at the 3', 4', and 5' positions of the ribose moiety.  
 The CoMFA and CoMSIA models yielded significant cross-validated q2 values  
 of 0.53 (r2 = 0.92) and 0.59 (r2 = 0.92), resp., and were further  
 validated by an external test set (25 adenosine derivs.), resulting in the  
 best predictive r2 values of 0.84 and 0.70 in each model. Both the CoMFA  
 and the CoMSIA maps for steric or hydrophobic, electrostatic, and  
 hydrogen-bonding interactions well reflected the nature of the putative  
 binding site previously obtained by mol. docking. A conformationally  
 restricted bulky group at the N6 or C2 position of the adenine ring and a  
 hydrophilic and/or H-bonding group at the 5' position were predicted to  
 increase A3AR binding affinity. A small hydrophobic group at N6 promotes  
 receptor activation. A hydrophilic and hydrogen-bonding moiety at the 5'  
 position appears to contribute to the receptor activation process, associated  
 with the conformational change of transmembrane domains 5, 6, and 7. The  
 3D-CoMFA/CoMSIA model correlates well with previous receptor-docking  
 results, current data of A3AR agonists, and the successful conversion of  
 the A3AR agonist into antagonists by substitution (at N6) or  
 conformational constraint (at 5'-N-methyluronamide).  
 IT 50257-82-4 50257-85-7 50257-95-9,  
 2-Hexyloxyadenosine 131865-78-6 131865-81-1  
 131865-82-2 131933-15-8  
 RL: PAC (Pharmacological activity); PRP (Properties); BIOL (Biological  
 study)  
 (QSAR of nucleosides acting at A3 adenosine receptor)  
 RN 50257-82-4 CAPLUS  
 CN Adenosine, 2-phenoxy- (CA INDEX NAME)


Absolute stereochemistry.

10/598,520




RN 50257-85-7 CAPLUS  
CN Adenosine, 2-(phenyloxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 50257-95-9 CAPLUS  
CN Adenosine, 2-(hexyloxy)- (CA INDEX NAME)

Absolute stereochemistry.



RN 131865-78-6 CAPLUS  
CN Adenosine, 2-(phenylmethoxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 131865-81-1 CAPLUS  
CN Adenosine, 2-(4-phenylbutoxy)- (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 131865-82-2 CAPLUS  
CN Adenosine, 2-[(5-phenylpentyl)oxy]- (CA INDEX NAME)

Absolute stereochemistry.



RN 131933-15-8 CAPLUS  
CN Adenosine, 2-(3-methylbutoxy)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 9 THERE ARE 9 CAPLUS RECORDS THAT CITE THIS RECORD (9 CITINGS)  
RE.CNT 39 THERE ARE 39 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 29 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 2007:119050 CAPLUS  
DN 146:198709  
TI Neuroprotective agents for treating neurodegenerative diseases  
IN Stockwell, Brent R.; Hoffstrom, Benjamin; Varma, Hemant  
PA Columbia University, USA  
SO U.S. Pat. Appl. Publ., 83 pp., Cont.-in-part of U.S. Ser. No. 837,360.  
CODEN: USXXCO  
DT Patent  
LA English  
FAN.CNT 5

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 20070027164	A1	20070201	US 2006-349653	20060207
	US 20050032124	A1	20050210	US 2004-767591	20040129
	US 20070078144	A1	20070405	US 2006-498110	20060802
	US 20070149543	A1	20070628	US 2006-612286	20061218
PRAI	US 2003-443728P	P	20030129		
	US 2003-457401P	P	20030325		
	US 2003-467290P	P	20030502		
	US 2003-482688P	P	20030625		
	US 2003-496209P	P	20030819		
	US 2004-767591	A2	20040129		

McIntosh

US 2004-837360 A2 20040430  
 US 2006-349653 A2 20060207  
 US 2006-498110 A2 20060802

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OS MARPAT 146:198709

AB The invention relates to compds. effective in preventing neuronal cell death, which may be used in the treatment of neurodegenerative diseases. It is based, at least in part, on the discovery that particular chemotherapeutic compds. were effective in preventing neuronal death in model systems of Huntington's Disease.


IT 53296-10-9, 2-Phenylaminoadenosine

RL: DMA (Drug mechanism of action); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (agents for treating neurodegenerative diseases)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



L6 ANSWER 30 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 2006:854003 CAPLUS

DN 146:221613

TI Cloning and pharmacological characterization of the equine adenosine A2A receptor: a potential therapeutic target for the treatment of equine endotoxemia

AU Brandon, C. I.; Vandenplas, M.; Dookwah, H.; Linden, J.; Murray, T. F.  
 CS Departments of Physiology and Pharmacology, College of Veterinary Medicine, University of Georgia, Athens, GA, USA

SO Journal of Veterinary Pharmacology and Therapeutics (2006), 29(4), 243-253  
 CODEN: JVPTD9; ISSN: 0140-7783

PB Blackwell Publishing Ltd.

DT Journal

LA English

AB The aim of the current study was to clone the equine adenosine A2A receptor gene and to establish a heterologous expression system to ascertain its pharmacol. profile via radioligand binding and functional assays. An eA2A-R expression construct was generated by ligation of the eA2A cDNA into the pcDNA3.1 expression vector, and stably transfected into human embryonic kidney cells (HEK). Binding assays identified those clones expressing the eA2A-R, and equilibrium saturation isotherm expts. were utilized to determine dissociation consts. (KD), and receptor densities (Bmax) of selected clones. Equilibrium competition binding revealed a rank order of agonist potency of ATL > CV-1808 > NECA > 2-CADO > CGS21680, and a rank order of antagonist potency as ZM241385 > 8-phenyltheophylline > p-sulfophenyltheophylline > caffeine. Furthermore, adenylate cyclase assays using selective A2A-R agonists revealed that the eA2A-R functionally coupled to Gas as indicated by an increase in intracellular [3H]cAMP upon receptor activation. Finally, NF-κB reporter gene assays revealed a CGS21680 concentration-dependent inhibition of NF-κB activity. These results indicate that the heterologously expressed eA2A-R has a pharmacol. profile similar to that of other mammalian A2A receptors and thus can be utilized for further characterization of the eA2A-R to ascertain whether it can serve as a suitable pharmacol. target for equine inflammatory disease.

IT 53296-10-9, CV-1808


RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (binding potentials of equine adenosine A2A receptor were determined by using adenosine A2A receptor agonist CV-1808 in human embryonic kidney cells)

RN 53296-10-9 CAPLUS

10/598,520

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 7 THERE ARE 7 CAPLUS RECORDS THAT CITE THIS RECORD (7 CITINGS)  
RE.CNT 21 THERE ARE 21 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 31 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 2006:756818 CAPLUS

DN 145:203066

TI Functional coupling of the G $\alpha$ olf variant XLG $\alpha$ olf with the human adenosine A2A receptor

AU Ravyn, Vipaj; Bostwick, J. Robert

CS Lead Discovery, AstraZeneca Pharmaceuticals, Wilmington, DE, USA

SO Journal of Receptors and Signal Transduction (2006), 26(4), 241-258

CODEN: JRSTCT

PB Taylor & Francis, Inc.

DT Journal

LA English

AB A recently identified novel G $\alpha$ olf variant, XLG $\alpha$ olf, is shown to functionally couple to the human adenosine A2A receptor (A2AR). In Sf9 cells expressing A2AR,  $\beta$ 1, and  $\gamma$ 2, co-expression of XLG $\alpha$ olf increased NECA-induced [35S]GTP $\gamma$ S binding from approx. 130% to 300% of basal levels. Pharmacol. characteristics of A2AR ligands on these cells were evaluated by using [3H]ZM241385- and [35S]GTP $\gamma$ S-binding assays. The rank order of the equilibrium binding consts. (Kd or Ki) of adenosine receptor ligands were [3H]ZM241385  $\approx$  CGS15943 < MRS1220 < CV1808  $\approx$  NECA < CGS21680  $\approx$  adenosine < IBMECA < HEMADO  $\approx$  CPA  $\approx$  CCPA. The rank order of EC50 values for agonists were CV1808  $\approx$  NECA < adenosine  $\approx$  CGS26180 < IBMECA < HEMADO  $\approx$  CPA  $\approx$  CCPA. This pharmacol. is consistent with the literature for A2AR and suggests that Sf9 cells co-expressing A2AR,  $\beta$ 1,  $\gamma$ 2, and XLG $\alpha$ olf could serve as a heterologous expression system for A2AR drug screening.

IT 53296-10-9, CV1808


RL: BSU (Biological study, unclassified); PKT (Pharmacokinetics); BIOL (Biological study)

(functional coupling of G $\alpha$ olf variant XLG $\alpha$ olf with the human adenosine A2A receptor)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

RE.CNT 42 THERE ARE 42 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

McIntosh


L6 ANSWER 32 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
 AN 2006:365436 CAPLUS  
 DN 144:412837  
 TI Preparation of substituted adenine nucleosides as antibacterial agents  
 IN Caverro-Tomas, Marta; Gowravaram, Madhu; Huynh, Hoan; Ni, Haihong; Stokes, Suzanne  
 PA Astrazeneca AB, Swed.; Astrazeneca Uk Limited  
 SO PCT Int. Appl., 180 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2006040558	A1	20060420	WO 2005-GB3934	20051013
	W:				
	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW:				
	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	EP 1805196	A1	20070711	EP 2005-792667	20051013
	R:				
	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR				
	CN 101072787	A	20071114	CN 2005-80042160	20051013
	JP 2008516938	T	20080522	JP 2007-536258	20051013
	AR 51393	A1	20070110	AR 2005-104326	20051014
	US 20090048203	A1	20090219	US 2007-577278	20070413
	IN 2007DN03360	A	20070831	IN 2007-DN3360	20070504
PRAI	US 2004-619218P	P	20041015		
	US 2005-663459P	P	20050318		
	US 2005-699615P	P	20050715		
	WO 2005-GB3934	W	20051013		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OS CASREACT 144:412837; MARPAT 144:412837

GI




AB Adenine nucleosides I, wherein X is O, CH<sub>2</sub>; Y is O, S, CO, CH<sub>2</sub>, CH=CH, SO, SO<sub>2</sub>; Y and R taken together form heterocycle; R is alkyl, alkenyl, alkynyl, carbocycle, sulfonyl, acyl, heterocycle; R<sub>1</sub>-R<sub>3</sub> are independently H, OH, CN, N<sub>3</sub>, alkyl, carbocycle, halogen, acyl, O-acyl, sulfonyl, oxime, alkenyl, alkynyl, heterocycle, alkoxy, substituted amine; were prepared and their use in the treatment of bacterial infections is reported. Thus, 9-[3-bromo-3,5-dideoxy-5-fluoro-2-O-[(isopropylamino)carbonyl]-β-D-xylofuranosyl]-2-(cyclopentyl-oxy)-9H-purin-6-amine was prepared and tested in vitro as antibacterial agent. A method for inhibition of bacterial DNA ligase in a warm-blooded animal, such as a human, in need of such treatment which comprises administering to human an effective amount of title compds. The compds. described have a measured IC<sub>50</sub> of 0.5-1.8 μM range in vitro against at least one isoenzyme (*S. pneumoniae*, *S. aureus*, *H. influenzae*, *E. coli*, or *M. pneumoniae*) of < 400 μM or the compds.

10/598,520

inhibited the ligation reaction by >20 % at the limit of their solubility in the assay medium. A formulation intended for oral administration to humans will generally contain, for example, from 0.5 mg to 2 g of active agent compounded with an appropriate and convenient amount of excipient which may vary from about 5 to about 98 percent by weight of the total composition Dosage unit forms will generally contain about 1 mg to about 500 mg of an active ingredient.


IT 131865-78-6P 756818-76-5P 883729-40-6P  
883729-41-7P 883731-37-1P 883731-38-2P  
RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU  
(Therapeutic use); BIOL (Biological study); PREP (Preparation); USES  
(Uses)  
(preparation of substituted adenine nucleosides as antibacterial agents)  
RN 131865-78-6 CAPLUS  
CN Adenosine, 2-(phenylmethoxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 756818-76-5 CAPLUS  
CN Adenosine, 2-(cyclopentylmethoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 883729-40-6 CAPLUS  
CN Adenosine, 2-[[1-(trifluoromethyl)cyclobutyl]methoxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 883729-41-7 CAPLUS  
CN Adenosine, 2-[(2,3,3-trifluorocyclobutyl)methoxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 883731-37-1 CAPLUS  
CN Adenosine, 2-(cyclobutylmethoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 883731-38-2 CAPLUS  
CN Adenosine, 2-(2,2,2-trifluoroethoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 6 THERE ARE 6 CAPLUS RECORDS THAT CITE THIS RECORD (6 CITINGS)  
RE.CNT 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 33 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 2005:1004549 CAPLUS  
DN 143:286636  
TI Preparation of nucleosides as adenosine receptors and used for the  
treatment of pain and inflammation  
IN Pritchard, Martyn; Ouzman, Jacqueline; Savory, Edward; Brown, Giles  
PA Cambridge Biotechnology Limited, UK  
SO PCT Int. Appl., 81 pp.  
CODEN: PIXXD2  
DT Patent  
LA English  
FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2005084653	A2	20050915	WO 2005-GB800	20050304
	WO 2005084653	A3	20060518		
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,			

McIntosh




LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI,  
 NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM,  
 SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW  
 RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM,  
 AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK,  
 EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT,  
 RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML,  
 MR, NE, SN, TD, TG  
 WO 2004079329 A2 20040916 WO 2004-GB902 20040305  
 WO 2004079329 A3 20041209  
 W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH,  
 CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD,  
 GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
 LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI  
 RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE,  
 BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU,  
 MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA,  
 GN, GQ, GW, ML, MR, NE, SN, TD, TG  
 AU 2005218997 A1 20050915 AU 2005-218997 20050304  
 CA 2557285 A1 20050915 CA 2005-2557285 20050304  
 EP 1749016 A2 20070207 EP 2005-717878 20050304  
 R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE,  
 IS, IT, LI, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR  
 CN 1946732 A 20070411 CN 2005-80007119 20050304  
 BR 2005008488 A 20070731 BR 2005-8488 20050304  
 JP 2007526291 T 20070913 JP 2007-501345 20050304  
 NZ 549235 A 20100129 NZ 2005-549235 20050304  
 MX 2006010075 A 20070410 MX 2006-10075 20060904  
 NO 2006004365 A 20061122 NO 2006-4365 20060926  
 KR 2007004792 A 20070109 KR 2006-7020304 20060929  
 IN 2006CN03674 A 20070706 IN 2006-CN3674 20061005  
 US 20080221060 A1 20080911 US 2007-598520 20071207  
 PRAI GB 2004-5009 A 20040305  
 GB 2004-5012 A 20040305  
 WO 2004-GB902 A 20040305  
 GB 2004-12261 A 20040602  
 GB 2004-12262 A 20040602  
 GB 2004-13627 A 20040618  
 GB 2004-19718 A 20040906  
 GB 2004-20063 A 20040909  
 GB 2004-20615 A 20040916  
 GB 2003-5153 A 20030307  
 WO 2005-GB800 W 20050304

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OS CASREACT 143:286636; MARPAT 143:286636

GI



I

AB Nucleosides I, wherein X is H, OH; R is H, Me; R1 is H, alkoxy,  
 OCH2-cyclopropyl, OCH2-cyclopentyl, phenoxy, OCH2CH2OH, OCH2CH2F2,  
 (5-indanyl)oxy, alkylamino, cyclo-alkylamino, exo-norbornane, amino,  
 phenylamino; R2 is NH2, CH2OH, NMe2, methylamino, isoamyl; R3 is CH2OH,  
 amide, CH2NHCOPr-n, CH2NHCONHET; were prepared and used for the treatment of  
 pain and inflammation. Title nucleosides were prepared and used the  
 treatment of pain associated with cancer, pancreatic pain, pain associated with

HIV infection, chronic neuropathic pain, lower back pain, failed back surgery pain, back pain, post-operative pain, post phys. trauma pain, cardiac pain, chest pain, joint pain, neck pain, bowel pain, phantom limb pain, obstetric pain, acute herpes zoster pain, acute pancreatitis breakthrough pain, or for the prevention, treatment, or amelioration of neuropathic or other pain caused by, or associated with diabetic neuropathy, poly-neuropathy, fibromyalgia, myo-fascial pain syndrome, osteoarthritis, post herpetic neuralgia, rheumatoid arthritis, sciatica/lumbar radiculopathy, spinal stenosis, trigeminal neuralgia, renal colic, dysmenorrhea/endometriosis. Thus, I (R = H, R1 = OMe, R2 = NH1, R3 = CH2OH) was prepared and tested for the treatment of pain and inflammation.

IT 50257-82-4P 50257-84-6P 50257-89-1P  
 50257-95-9P 53296-10-9P 53296-19-8P  
 70255-72-0P 71231-79-3P 79936-11-1P  
 756818-70-9P 756818-71-0P 756818-72-1P  
 756818-73-2P 756818-76-5P 756818-78-7P  
 864061-82-5P 864061-83-6P 864061-84-7P  
 864061-85-8P 864061-86-9P 864061-87-0P  
 864061-88-1P 864061-89-2P 864061-90-5P  
 864061-91-6P 864061-94-9P 864061-95-0P


RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of nucleosides as adenosine receptors and used for the treatment of pain and inflammation)

RN 50257-82-4 CAPLUS

CN Adenosine, 2-phenoxy- (CA INDEX NAME)


Absolute stereochemistry.



RN 50257-84-6 CAPLUS

CN Adenosine, 2-butoxy- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 50257-89-1 CAPLUS

CN Adenosine, 2-(2-hydroxyethoxy)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

10/598,520




RN 50257-95-9 CAPLUS  
CN Adenosine, 2-(hexyloxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



RN 53296-19-8 CAPLUS  
CN Adenosine, 2-[(4-methoxyphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 70255-72-0 CAPLUS  
CN Adenosine, 2-(aminocarbonyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 71231-79-3 CAPLUS  
CN Adenosine, 2-[(4-fluorophenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 79936-11-1 CAPLUS  
CN Adenosine, 2-cyano- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 756818-70-9 CAPLUS  
CN Adenosine, 2-(4-cyanophenoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 756818-71-0 CAPLUS  
CN Adenosine, 2-([1,1'-biphenyl]-3-yloxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 756818-72-1 CAPLUS  
CN Adenosine, 2-[(2,3-dihydro-1H-inden-5-yl)oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 756818-73-2 CAPLUS  
CN Adenosine, 2-[3-(1-methylethyl)phenoxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 756818-76-5 CAPLUS  
CN Adenosine, 2-(cyclopentylmethoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 756818-78-7 CAPLUS  
CN Adenosine, 2-heptyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 864061-82-5 CAPLUS  
CN Adenosine, 2-(2,2-difluoroethoxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 864061-83-6 CAPLUS  
CN Adenosine, 2-(cyclopropylmethoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 864061-84-7 CAPLUS  
CN Adenosine, 2-(2,5-difluorophenoxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 864061-85-8 CAPLUS  
CN Adenosine, 2-(2,4-difluorophenoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 864061-86-9 CAPLUS  
CN Adenosine, 2-(3,4-difluorophenoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 864061-87-0 CAPLUS  
CN Adenosine, 2-(2,3,5-trifluorophenoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 864061-88-1 CAPLUS  
CN Adenosine, 2-(4-fluoro-3-methylphenoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 864061-89-2 CAPLUS  
CN Adenosine, 2-(2-methylphenoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 864061-90-5 CAPLUS  
CN Adenosine, 2-(3-bromophenoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 864061-91-6 CAPLUS  
CN Adenosine, 2-(4-methylphenoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 864061-94-9 CAPLUS  
CN Adenosine, 2-[(2,2,3,3-tetrafluorocyclobutyl)oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 864061-95-0 CAPLUS  
CN Adenosine, 2-(3,5-dimethylphenyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

McIntosh




10/598,520



OSC.G 4 THERE ARE 4 CAPLUS RECORDS THAT CITE THIS RECORD (4 CITINGS)

L6 ANSWER 34 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 2005:44237 CAPLUS  
DN 142:290603  
TI A radial distribution function approach to predict A2B agonist effect of adenosine analogues  
AU Gonzalez, Maykel Perez; Teran, Carmen; Fall, Yagamare; Teijeira, Marta; Besada, Pedro  
CS Unit of Services, Department of Drug Design, Experimental Sugar Cane Station 'Villa Clara-Cienfuegos', Ranchuelo, Cuba  
SO Bioorganic & Medicinal Chemistry (2005), 13(3), 601-608  
CODEN: BMECEP; ISSN: 0968-0896  
PB Elsevier Ltd.  
DT Journal  
LA English  
AB The radial distribution function (RDF) approach has been applied to the study of the A2B agonist effect of a set of 89 adenosine analogs reported with this activity. A model able to describe more than 70% of the variance in the exptl. activity was developed with the use of the mentioned approach. In contrast, none of the eleven different approaches including the use of Constitutional, Topol., Mol. walk count, BCUT, Galvez topol. charge indexes, 2D autocorrelations, Randic mol. profiles, Geometrical, 3D Morse, WHIM and GETAWAY descriptors was able to explain more than 47% of the variance in the mentioned property with the same number of descriptors.  
IT 53296-10-9  
RL: PAC (Pharmacological activity); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(radial distribution function approach to predict A2B agonist effect of adenosine analogs)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 21 THERE ARE 21 CAPLUS RECORDS THAT CITE THIS RECORD (21 CITINGS)  
RE.CNT 33 THERE ARE 33 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 35 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 2005:34766 CAPLUS  
DN 142:127629  
TI Compositions and methods for use of a protease inhibitor and adenosine for preventing organ ischemia and reperfusion injury  
IN Vinten-Johansen, Jakob  
PA Emory University, USA

McIntosh

10/598,520

SO PCT Int. Appl., 82 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2005003150	A2	20050113	WO 2004-US21387	20040702
	WO 2005003150	A3	20051013		
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
	RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
	CA 2531062	A1	20050113	CA 2004-2531062	20040702
	EP 1638579	A2	20060329	EP 2004-756603	20040702
	R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK, HR			
	US 20060205671	A1	20060914	US 2006-562757	20060328
PRAI	US 2003-484484P	P	20030702		
	WO 2004-US21387	W	20040702		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB Methods and compns. including combined use of a serine protease inhibitor and adenosine or adenosine agonist when administered as a single pharmaceutical composition, concomitantly or sequentially in any order to a living subject for preventing organ ischemia or reperfusion injury. The methods and compns. disclosed herein can be used in such procedures as cardiac surgery, non-surgical cardiac revascularization, organ transplantation, perfusion, ischemia, reperfusion, ischemia-reperfusion injury, oxidant injury, cytokine induced injury, shock induced injury, resuscitations injury or apoptosis.

IT 53296-10-9, CV1808

RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL


(Biological study); USES (Uses)

(use of a serine protease inhibitor and adenosine agonist for preventing organ ischemia and reperfusion injury in relation to alteration of G protein-coupled receptors and cAMP)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



L6 ANSWER 36 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 2004:829441 CAPLUS

DN 141:420038

TI 2-Substituted adenosine derivatives: affinity and efficacy at four subtypes of human adenosine receptors

AU Gao, Zhan-Guo; Mamedova, Liaman K.; Chen, Peiran; Jacobson, Kenneth A.

CS Molecular Recognition Section, Laboratory of Bioorganic Chemistry, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, MD, 20892, USA

SO Biochem. Pharmacol. (2004), 68(10), 1985-1993

CODEN: BCPA6; ISSN: 0006-2952

PB Elsevier B.V.

McIntosh


DT Journal

LA English

AB The affinity and efficacy at four subtypes (A1, A2A, A2B and A3) of human adenosine receptors (ARs) of a wide range of 2-substituted adenosine derivs. were evaluated using radioligand binding assays and a cAMP functional assay in intact CHO cells stably expressing these receptors. Similar to previous studies of the N6-position, several 2-substituents were found to be critical structural determinants for the A3AR activation. The following adenosine 2-ethers were moderately potent partial agonists ( $K_i$ , nM): benzyl (117), 3-chlorobenzyl (72), 2-(3-chlorophenyl)ethyl (41), and 2-(2-naphthyl)ethyl (130). The following adenosine 2-ethers were A3AR antagonists: 2,2-diphenylethyl, 2-(2-norbornan)ethyl, R- and S-2-phenylbutyl, and 2-(2-chlorophenyl)ethyl. 2-(S-2-Phenylbutyloxy)adenosine as an A3AR antagonist right-shifted the concentration-response curve for the inhibition by NECA of cAMP accumulation with a  $K_B$  value of 212 nM, which is similar to its binding affinity ( $K_i$  = 175 nM). These 2-substituted adenosine derivs. were generally less potent at the A1AR in comparison to the A3AR, but fully efficacious, with binding  $K_i$  values over 100 nM. The 2-phenylethyl moiety resulted in higher A3AR affinity ( $K_i$  in nM) when linked to the 2-position of adenosine through an ether group (54), than when linked through an amine (310) or thioether (1960). 2-[2-(1-Naphthyl)ethyloxy]adenosine ( $K_i$  = 3.8 nM) was found to be the most potent and selective (>50-fold) A2A agonist in this series. Mixed A2A/A3AR agonists have been identified. Interestingly, although most of these compds. were extremely weak at the A2BAR, 2-[2-(2-naphthyl)ethyloxy]adenosine ( $EC_{50}$  = 1.4  $\mu$ M) and 2-[2-(2-thienyl)-ethyloxy]adenosine ( $EC_{50}$  = 1.8  $\mu$ M) were found to be relatively potent A2B agonists, although less potent than NECA ( $EC_{50}$  = 140 nM).


IT 50257-82-4 50257-85-7 50257-95-9,  
2-(Hexyloxy)adenosine 131865-78-6 131865-81-1  
131865-82-2 131933-15-8 131933-17-0  
131933-20-5 131933-26-1 131973-26-7  
137817-83-5 137817-84-6 194154-31-9  
194154-32-0 794597-49-2,  
2-(3-Chlorobenzoyloxy)adenosine  
RL: PAC (Pharmacological activity); PRP (Properties); THU (Therapeutic  
use); BIOL (Biological study); USES (Uses)  
(affinity and efficacy at four subtypes of human adenosine receptors of  
2-substituted adenosine derivs.)  
RN 50257-82-4 CAPLUS  
CN Adenosine, 2-phenoxy- (CA INDEX NAME)

Absolute stereochemistry.



RN 50257-85-7 CAPLUS  
CN Adenosine, 2-(pentyloxy)- (CA INDEX NAME)


Absolute stereochemistry.



10/598,520


RN 50257-95-9 CAPLUS  
CN Adenosine, 2-(hexyloxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 131865-78-6 CAPLUS  
CN Adenosine, 2-(phenylmethoxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 131865-81-1 CAPLUS  
CN Adenosine, 2-(4-phenylbutoxy)- (CA INDEX NAME)

Absolute stereochemistry.



RN 131865-82-2 CAPLUS  
CN Adenosine, 2-[(5-phenylpentyl)oxy]- (CA INDEX NAME)

Absolute stereochemistry.




RN 131933-15-8 CAPLUS  
CN Adenosine, 2-(3-methylbutoxy)- (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 131933-17-0 CAPLUS  
CN Adenosine, 2-(cyclohexylmethoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 131933-20-5 CAPLUS  
CN Adenosine, 2-(4-cyclohexylbutoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 131933-26-1 CAPLUS  
CN Adenosine, 2-(2-ethylbutoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 131973-26-7 CAPLUS  
CN Adenosine, 2-[(4-methylpentyl)oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 137817-83-5 CAPLUS  
CN Adenosine, 2-[(2R)-2-phenylbutoxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 137817-84-6 CAPLUS  
CN Adenosine, 2-[(2S)-2-phenylbutoxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 194154-31-9 CAPLUS  
CN Adenosine, 2-[[ (3S)-3,7-dimethyl-6-octenyl]oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 194154-32-0 CAPLUS  
CN Adenosine, 2-[[ (3R)-3,7-dimethyl-6-octenyl]oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520



RN 794597-49-2 CAPLUS  
CN Adenosine, 2-[(3-chlorophenyl)methoxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 34 THERE ARE 34 CAPLUS RECORDS THAT CITE THIS RECORD (34 CITINGS)  
RE.CNT 37 THERE ARE 37 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 37 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 2004:756969 CAPLUS

DN 141:254620

TI Identification of therapeutic compounds

IN Richardson, Peter

PA Cambridge Biotechnology Ltd., UK

SO PCT Int. Appl., 44 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2004079329	A2	20040916	WO 2004-GB902	20040305
	WO 2004079329	A3	20041209		
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI			
	RW:	BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
	AU 2004217731	A1	20040916	AU 2004-217731	20040305
	AU 2004217731	B2	20090604		
	CA 2514338	A1	20040916	CA 2004-2514338	20040305
	EP 1604211	A2	20051214	EP 2004-717679	20040305
	EP 1604211	B1	20080430		
	R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK			
	JP 2006519602	T	20060831	JP 2006-505911	20040305
	US 20070059773	A1	20070315	US 2004-547462	20040305
	AT 393917	T	20080515	AT 2004-717679	20040305
	PT 1604211	E	20080704	PT 2004-717679	20040305
	ES 2305741	T3	20081101	ES 2004-717679	20040305
	AU 2005218997	A1	20050915	AU 2005-218997	20050304
	CA 2557285	A1	20050915	CA 2005-2557285	20050304
	WO 2005084653	A2	20050915	WO 2005-GB800	20050304
	WO 2005084653	A3	20060518		
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH,			

McIntosh

CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW  
 RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

EP 1749016 A2 20070207 EP 2005-717878 20050304  
 R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR

CN 1946732 A 20070411 CN 2005-80007119 20050304  
 BR 2005008488 A 20070731 BR 2005-8488 20050304  
 JP 2007526291 T 20070913 JP 2007-501345 20050304  
 SG 144146 A1 20080729 SG 2008-4350 20050304  
 NZ 549235 A 20100129 NZ 2005-549235 20050304  
 MX 2006010075 A 20070410 MX 2006-10075 20060904  
 NO 2006004365 A 20061122 NO 2006-4365 20060926  
 KR 2007004792 A 20070109 KR 2006-7020304 20060929  
 US 20080221060 A1 20080911 US 2007-598520 20071207

PRAI GB 2003-5153 A 20030307  
 GB 2004-5009 A 20040305  
 GB 2004-5012 A 20040305  
 WO 2004-GB902 W 20040305  
 GB 2004-12261 A 20040602  
 GB 2004-12262 A 20040602  
 GB 2004-13627 A 20040618  
 GB 2004-19718 A 20040906  
 GB 2004-20063 A 20040909  
 GB 2004-20615 A 20040916  
 WO 2005-GB800 W 20050304

# ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB Methods for identifying potential therapeutic agents involve determining the affinity and/or efficacy of a test compound for an adenosine receptor at a relatively high pH and at a relatively low pH. Compds. with greater affinity and/or efficacy at the low pH are identified as potential therapeutic agents, in particular for the treatment of pain or inflammation.


IT 50257-82-4 50257-84-6 50257-89-1  
 50257-95-9 53296-10-9, 2-Phenylaminoadenosine  
 53296-19-8 71231-79-3 756818-70-9  
 756818-71-0 756818-72-1 756818-73-2  
 756818-76-5 756818-78-7

RL: ADV (Adverse effect, including toxicity); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (identification of therapeutic compds.)

RN 50257-82-4 CAPLUS

CN Adenosine, 2-phenoxy- (CA INDEX NAME)

Absolute stereochemistry.




RN 50257-84-6 CAPLUS

CN Adenosine, 2-butoxy- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




10/598,520




RN 50257-89-1 CAPLUS  
CN Adenosine, 2-(2-hydroxyethoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 50257-95-9 CAPLUS  
CN Adenosine, 2-(hexyloxy)- (CA INDEX NAME)

Absolute stereochemistry.



RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.




RN 53296-19-8 CAPLUS  
CN Adenosine, 2-[(4-methoxyphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 71231-79-3 CAPLUS  
CN Adenosine, 2-[(4-fluorophenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 756818-70-9 CAPLUS  
CN Adenosine, 2-(4-cyanophenoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 756818-71-0 CAPLUS  
CN Adenosine, 2-([1,1'-biphenyl]-3-yloxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 756818-72-1 CAPLUS  
CN Adenosine, 2-[(2,3-dihydro-1H-inden-5-yl)oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

McIntosh


10/598,520



RN 756818-73-2 CAPLUS

CN Adenosine, 2-[3-(1-methylethyl)phenoxy]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 756818-76-5 CAPLUS

CN Adenosine, 2-(cyclopentylmethoxy)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 756818-78-7 CAPLUS

CN Adenosine, 2-heptyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)  
RE.CNT 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 38 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 2004:754440 CAPLUS

DN 141:271600

TI Use of adenosine receptor agonists in therapy

IN Richardson, Peter

McIntosh

10/598,520


PA Cambridge Biotechnology Ltd., UK  
SO PCT Int. Appl., 25 pp.  
CODEN: PIXXD2  
DT Patent  
LA English  
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2004078184	A1	20040916	WO 2004-GB952	20040305
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI			
	RW:	BW, CH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
	AU 2004216891	A1	20040916	AU 2004-216891	20040305
	AU 2004216891	B2	20090618		
	CA 2514848	A1	20040916	CA 2004-2514848	20040305
	EP 1603576	A1	20051214	EP 2004-717693	20040305
	R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK			
	CN 1809365	A	20060726	CN 2004-80005723	20040305
	JP 2006519824	T	20060831	JP 2006-505924	20040305
	NZ 541587	A	20080829	NZ 2004-541587	20040305
	NO 2005004475	A	20050927	NO 2005-4475	20050927
	IN 2005CN02547	A	20070831	IN 2005-CN2547	20051005
	US 20060234975	A1	20061019	US 2006-547454	20060628
	US 7790698	B2	20100907		
PRAI	GB 2003-5150	A	20030307		
	WO 2004-GB952	W	20040305		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OS MARPAT 141:271600

GI



AB The invention describes the use of compds. I (R = C1-4 alkoxy; X = H, OH) for the prevention, treatment, or amelioration of cancer, inflammation, autoimmune disease, ischemia-reperfusion injury, epilepsy, sepsis, septic shock, neurodegeneration (including Alzheimer's Disease), muscle fatigue or muscle cramp. The compds. are effective at very low doses, and so can be administered at doses at which serious side effects are not observed


IT 50257-84-6

RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(adenosine receptor agonists for therapy)

RN 50257-84-6 CAPLUS

CN Adenosine, 2-butoxy- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD (3 CITINGS)  
 RE.CNT 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 39 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
 AN 2004:570030 CAPLUS  
 DN 141:99661  
 TI Identification of compounds suitable as agonists and/or antagonists of adenosine A2A receptor coupled to specific G proteins, and use of identified compounds in treatment of various disorders in mammals  
 IN Fredholm, Bertil B.; Kull, Bjoern  
 PA Actar Ab, Swed.  
 SO PCT Int. Appl., 22 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004058974	A1	20040715	WO 2003-SE2086	20031229
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2003291608	A1	20040722	AU 2003-291608	20031229
PRAI US 2002-436480P	P	20021227		
WO 2003-SE2086	W	20031229		


AB The invention discloses a method of drug screening to select chemical compds. suitable as receptor agonists or antagonists that act on a receptor belonging to the family of G protein coupled receptors. The method involves constructing a biol. preparation comprising a receptor coupled to a specific G protein, bringing a compound in contact with said preparation, and studying the functional properties of said compound in biol. preparation. The invention also discloses the use of identified compound as a drug for treatment of CNS disorders, cardiovascular disorders, inflammatory disorders, metabolic disorders, cancer and other hyperproliferative disorders in a mammal. Specifically, the invention discloses a novel approach to identify and test compds. based on changes in adenosine A2A receptor binding dependent on the nature of the coupled G protein. The invention related that this approach seemed feasible due to the evidence that G proteins, such as Gs, Gi or Golf, influence the binding properties of A2A receptors. The invention also related that a truly efficient drug mol., in addition to affinity towards A2A, can be specific towards the different G proteins associated with A2A. In the examples, the invention used transformed CHO cells expressing A2A receptors linked to Gs or Golf G proteins, and demonstrated the existence of G-protein-dependent ligand specificity. Specifically, the examples demonstrated that substance KF 17837 has higher affinity to the A2A-Golf complex in striatum than to A2A-Gs complex in leukocytes. The examples also showed how some compds. influence A2A receptors coupled to Golf more readily than A2A receptors coupled to Gs.

IT 53296-10-9  
 RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (binding of chemical compds. to adenosine A2A receptors in membrane preparation derived from pig brain (A2A receptors coupled to Golf) or from pig

10/598,520

lymphocytes (A2A receptors coupled to Gs))  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)  
RE.CNT 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 40 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 2004:567572 CAPLUS

DN 142:89098

TI WGA-Coated Yttrium Oxide Beads Enable an Imaging-Based Adenosine 2a  
Receptor Binding Scintillation Proximity Assay Suitable for High  
Throughput Screening

AU Bryant, Robert; McGuinness, Debra; Turek-Etienne, Tammy; Guyer, Deborah;  
Yu, Liming; Howells, Leighton; Caravano, Joseph; Zhai, Ying; Lachowicz,  
Jean

CS Schering-Plough Research Institute, Kenilworth, NJ, USA

SO Assay and Drug Development Technologies (2004), 2(3), 290-299

CODEN: ADDTAR; ISSN: 1540-658X

PB Mary Ann Liebert, Inc.

DT Journal

LA English

AB Adenosine receptors belong to the superfamily of G protein-coupled  
receptors and are involved in a variety of physiolo. functions.  
Traditionally, binding assays to detect adenosine 2a (A2a) antagonists and  
agonists have used filtration methods that are cumbersome to run and not  
amenable to HTS. We developed scintillation proximity assays (SPA)  
utilizing HEK293 RBHA2AM cell membranes, either wheat germ agglutinin  
(WGA)-coated yttrium silicate (YSi) or red-shifted yttrium oxide (YO)  
beads and the A2a-selective radioligand [3H]SCH 58261. Both beads gave  
windows (total binding/nonspecific binding) of >5 and Kd values of 2-3 nM  
for the radioligand, in agreement with results obtained by filtration. In  
contrast, WGA-polyvinyltoluene as well as other bead types had windows of  
<3 and significant radioligand binding to the uncoated beads. A 384-well  
WGA-YO bead SPA was optimized utilizing a LEADseeker imaging system and an  
automated trituration process for dispensing the dense yttrium-based  
beads. Signals were stable after 4 h, and Z' values were 0.7-0.8. The  
LEADseeker imaging assay tolerated 2% DMSO and generated IC50 values of  
3-5 nM for the A2a antagonist CGS 15943, comparable to that obtained by  
the filtration method. A number of adenosine and xanthine analogs were  
identified as hits in the Library of Pharmacol. Active Compds. (LOPAC).  
This imaging-based A2a SPA enables HTS and is a major improvement over the  
filtration method.

IT 53296-10-9, 2-Phenylaminoadenosine

RL: ANT (Analyte); BSU (Biological study, unclassified); ANST (Analytical  
study); BIOL (Biological study)

(wheat germ agglutinin-coated yttrium oxide beads enable imaging-based  
adenosine 2a receptor binding scintillation proximity assay suitable  
for high throughput screening)


RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.

McIntosh

10/598,520




OSC.G 7 THERE ARE 7 CAPLUS RECORDS THAT CITE THIS RECORD (7 CITINGS)  
RE.CNT 17 THERE ARE 17 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 41 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 2004:432750 CAPLUS  
DN 141:11973  
TI Use of adenosine or its analogue in cosmetics for smoothing wrinkles  
IN Galey, Jean Baptiste  
PA L'Oreal, Fr.  
SO Fr. Demande, 17 pp.  
CODEN: FRXXBL  
DT Patent  
LA French  
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	FR 2847469	A1	20040528	FR 2002-14828	20021126
	FR 2847469	B1	20060407		
	EP 1424064	A1	20040602	EP 2003-292633	20031022
	EP 1424064	B1	20070606		
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
	AT 363888	T	20070615	AT 2003-292633	20031022
	ES 2287432	T3	20071216	ES 2003-292633	20031022
	US 20040146474	A1	20040729	US 2003-701495	20031106
PRAI	FR 2002-14828	A	20021126		
	US 2002-432634P	P	20021212		
AB	A cosmetic method to reduce the wrinkles of the face and/or relax the skin, comprises topical application of a composition containing, adenosine or its analogs on the skin. A cosmetic composition contained adenosine 0.10, stearic acid 3.00, a mixture of glyceryl mono-stearate and polyethylene glycol stearate 2.50, polyethylene glycol stearate 1.00, cyclopentadimethylsiloxane 10.00, excipients 3.00, vegetable oils 7.00, synthetic oil 6.00, preservative 1.20, polyoxyethylene methoxy dimethylsiloxane (16 EO) 1.00, silicone gum 0.20, acrylic copolymer in inverse emulsion (Simulgel 600) 1.700, stearyl alc. 1.00, and water q.s. 100%.				
IT	53296-10-9, 2-Phenylaminoadenosine RI: COS (Cosmetic use); BIOL (Biological study); USES (Uses) (use of adenosine or its analog in cosmetics for smoothing wrinkles)				
RN	53296-10-9 CAPLUS				
CN	Adenosine, 2-(phenylamino)- (CA INDEX NAME)				

Absolute stereochemistry.




OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)  
RE.CNT 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

McIntosh

10/598,520

L6 ANSWER 42 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 2004:406955 CAPLUS  
DN 141:64408  
TI A TOPS-MODE approach to predict affinity for A1 adenosine receptors.  
2-(Arylamino)adenosine analogues  
AU Perez Gonzalez, Maykel; Teran Moldes, Maria del Carmen  
CS Experimental Sugar Cane Station "Villa Clara-Cienfuegos", Services Unit,  
Drug Design Department, Ranchuelo, 53100, Cuba  
SO Bioorganic & Medicinal Chemistry (2004), 12(11), 2985-2993  
CODEN: BMECEP; ISSN: 0968-0896  
PB Elsevier Ltd.  
DT Journal  
LA English  
AB The TOPol. Sub-Structural Mol. Design (TOPS-MODE) approach has been  
applied to the study of the affinity of A1 adenosine receptor of different  
2-(arylamino)adenosine analogs. A model able to describe closed to 79% of  
the variance in the values for binding expts. of 32 analogs of these  
compsds. through multilinear regression anal. (MRA) was developed with the  
use of the mentioned approach. In contrast, no one of seven different  
approaches, including the use of Constitutional, Topol., Mol. walk counts,  
BCUT, Randic Mol. profiles, Geometrical, and RDF descriptors was able to  
explain more than 70% of the variance in the mentioned property with the  
same number of descriptors. In addition, the TOPS-MODE approach permitted to  
find the contribution of different fragments to the biol. property giving  
to the model a straightforward structural interpretability.  
IT 53296-10-9  
RL: PAC (Pharmacological activity); PRP (Properties); BIOL (Biological  
study)  
(TOPS-MODE approach to predict affinity for A1 adenosine receptors,  
studied using 2-(arylamino)adenosine analogs)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



RE.CNT 40 THERE ARE 40 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 43 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 2003:796432 CAPLUS  
DN 139:302061  
TI Synergy of dopamine D2 and adenosine A2 receptors activates protein kinase  
A (PKA) signaling via  $\beta/\gamma$  dimers, and use in the treatment of  
drug abuse and drug withdrawal  
IN Gordon, Adrienne S.; Diamond, Ivan F.; Yao, Lina  
PA The Regents of the University of California, USA  
SO PCT Int. Appl., 152 pp.  
CODEN: PIXXD2  
DT Patent  
LA English  
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2003082211	A2	20031009	WO 2003-US9629	20030327
	WO 2003082211	A3	20041216		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT,				

McIntosh



TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW  
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY,  
 KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES,  
 FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR,  
 BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG  
 AU 2003241281 A1 20031013 AU 2003-241281 20030327  
 US 20090137662 A1 20090528 US 2007-550331 20070222  
 PRAI US 2002-368417P P 20020327  
 WO 2003-US9629 W 20030327


# ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB The invention pertains to the discovery that a dopamine receptor agonist can activate PKA signaling and/or can act synergistically with an adenosine receptor to activate such signaling. In various embodiments, the invention exploits the synergy between the dopamine receptor pathway and an adenosine receptor pathway to provide methods of mitigating one or more symptoms produced by the chronic consumption of a substance of abuse or to mitigate one or more physiol. and/or behavioral symptoms associated with cessation of chronic consumption of a substance of abuse. In certain embodiments, the methods involve administering to a mammal an effective amount of an adenosine receptor antagonist and an effective amount of a dopamine receptor antagonist, where the effective amount of the adenosine receptor antagonist is lower than the effective amount of an adenosine receptor antagonist administered without the dopamine receptor antagonist.

IT 53296-10-9, 2-Phenylaminoadenosine  
 RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (synergy of dopamine D2 and adenosine A2 receptors activates protein kinase A signaling via  $\beta/\gamma$  dimers, and use in treatment of drug abuse and drug withdrawal)

RN 53296-10-9 CAPLUS  
 CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



L6 ANSWER 44 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
 AN 2003:335118 CAPLUS  
 DN 138:338409  
 TI Synthesis of 2-aralkoxyadenosines and 2-alkoxyadenosines via hydrolysis, alkylation, and amination reactions  
 IN Moorman, Allan R.  
 PA King Pharmaceuticals Research and Development, Inc., USA  
 SO PCT Int. Appl., 38 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 FAN.CNT 2


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003035662	A1	20030501	WO 2002-US34313	20021024
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
CA 2465264	A1	20030501	CA 2002-2465264	20021024

CA 2465264	C	20100907		
AU 2002336668	A1	20030506	AU 2002-336668	20021024
AU 2002336668	B2	20080110		
US 20030199686	A1	20031023	US 2002-281291	20021024
US 6951932	B2	20051004		
EP 1446413	A1	20040818	EP 2002-773916	20021024
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK				
BR 2002013557	A	20041026	BR 2002-13557	20021024
CN 1608076	A	20050420	CN 2002-826143	20021024
JP 2005511551	T	20050428	JP 2003-538175	20021024
NZ 532541	A	20050930	NZ 2002-532541	20021024
CN 1935152	A	20070328	CN 2006-10099934	20021024
IL 161608	A	20100531	IL 2002-161608	20021024
NO 2004002126	A	20040623	NO 2004-2126	20040524
NO 326625	B1	20090119		
US 20060135466	A1	20060622	US 2005-155212	20050616
US 7342003	B2	20080311		
PRAI US 2001-335169P	P	20011025		
US 2002-375723P	P	20020426		
CN 2002-826143	A3	20021024		
US 2002-281291	A2	20021024		
WO 2002-US34313	W	20021024		
US 2003-508804P	P	20031003		
US 2004-958470	A1	20041004		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OS CASREACT 138:338409; MARPAT 138:338409

GI



AB 2-Aralkyloxyadenosines and 2-alkoxyadenosines I, wherein R1 and R2 are independently hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aralkyl, substituted aralkyl, aryl, substituted aryl, heteroaryl, heterocyclic, or if taken together with the nitrogen atom, forms an azetidine ring or a 5-6 membered heterocyclic ring containing a total of one to four heteroatoms selected from nitrogen, oxygen, and sulfur; R3 is alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aralkyl, substituted aralkyl, aralkenyl, or substituted aralkenyl, were prepared via hydrolysis, alkylation, and amination reactions. Preferred methods of the invention include activating a guanosine compound followed by hydrolysis; alkylating the hydrolyzed compound with subsequent amination to provide a 2-aralkyloxyadenosine or a 2-alkoxyadenosine compound. The invention is particularly useful for synthesis of 2-[2-(4-chlorophenyl)ethoxy]adenosine.

IT 50257-84-6P 131865-78-6P 515138-96-2P  
515138-97-3P

RL: IMF (Industrial manufacture); SPN (Synthetic preparation); PREP (Preparation)


(synthesis of aralkoxyadenosines and alkoxyadenosines via hydrolysis, alkylation, and amination reactions)

RN 50257-84-6 CAPLUS

CN Adenosine, 2-butoxy- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

10/598,520




RN 131865-78-6 CAPLUS  
CN Adenosine, 2-(phenylmethoxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 515138-96-2 CAPLUS  
CN Adenosine, 2-[(4-nitrophenyl)methoxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 515138-97-3 CAPLUS  
CN Acetic acid, [(6-amino-9- $\beta$ -D-ribofuranosyl-9H-purin-2-yl)oxy]- (9CI)  
(CA INDEX NAME)

Absolute stereochemistry.



OSC.G 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)  
RE.CNT 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 45 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 2003:251382 CAPLUS  
DN 139:223688  
TI Allosteric interactions and QSAR: on the role of ligand hydrophobicity  
AU Hansch, Corwin; Garg, Rajni; Kurup, Alka; Mekapati, Suresh Babu

McIntosh

CS Department of Chemistry, Pomona College, Claremont, CA, 91711, USA

SO Bioorganic & Medicinal Chemistry (2003), 11(9), 2075-2084

CODEN: BMECEP; ISSN: 0968-0896

PB Elsevier Science Ltd.

DT Journal

LA English

AB A study of a very large database of QSAR (9100) has uncovered a few unusual examples where as one increases the hydrophobicity of the members of a set of congeners, activity decreases until at a certain point, activity begins to increase. Obviously a change in mechanism is involved. The only way we have found to rationalize this unusual event is by a change in the structure of the receptor. We have found this to occur with Hb, a substance first used to define allosteric reactions.

IT 50257-84-6 50257-85-7 50257-89-1

50257-95-9 131933-15-8 131933-17-0

131933-20-5 131933-26-1 131933-27-2

131973-26-7


RL: PAC (Pharmacological activity); PRP (Properties); BIOL (Biological study)

(A1-adenoreceptor activity prolongation; parabolic relationship between ligand hydrophobicity and activity in QSAR studies in relation to allosteric interactions)

RN 50257-84-6 CAPLUS

CN Adenosine, 2-butoxy- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 50257-85-7 CAPLUS

CN Adenosine, 2-(pentyloxy)- (CA INDEX NAME)


Absolute stereochemistry.



RN 50257-89-1 CAPLUS

CN Adenosine, 2-(2-hydroxyethoxy)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



10/598,520


RN 50257-95-9 CAPLUS  
CN Adenosine, 2-(hexyloxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 131933-15-8 CAPLUS  
CN Adenosine, 2-(3-methylbutoxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 131933-17-0 CAPLUS  
CN Adenosine, 2-(cyclohexylmethoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 131933-20-5 CAPLUS  
CN Adenosine, 2-(4-cyclohexylbutoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 131933-26-1 CAPLUS  
CN Adenosine, 2-(2-ethylbutoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 131933-27-2 CAPLUS  
CN Adenosine, 2-[(3-ethylpentyl)oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 131973-26-7 CAPLUS  
CN Adenosine, 2-[(4-methylpentyl)oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 13 THERE ARE 13 CAPLUS RECORDS THAT CITE THIS RECORD (13 CITINGS)  
RE.CNT 30 THERE ARE 30 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 46 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 2002:906668 CAPLUS  
DN 137:380042  
TI Methods and formulations for increasing the affinity of A1 adenosine  
receptor ligands for the A1 adenosine receptor using glycolipids  
IN Wilson, Constance Neely  
PA Endacea Inc., USA  
SO PCT Int. Appl., 36 pp.  
CODEN: PIXXD2  
DT Patent  
LA English  
FAN.CNT 1


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002095391	A1	20021128	WO 2002-US16218	20020523
	W: AU, CA, JP, US				
	RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR				
	CA 2441801	A1	20021128	CA 2002-2441801	20020523
	AU 2002311987	A1	20021203	AU 2002-311987	20020523
	EP 1390740	A1	20040225	EP 2002-739334	20020523
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				

McIntosh

10/598,520

JP 2005518331 T 20050623 JP 2002-591814 20020523  
US 20040121406 A1 20040624 US 2003-475925 20031024  
PRAI US 2001-293362P P 20010524  
WO 2002-US16218 W 20020523  
AB Glycolipids are useful for enhancing the affinity of A1 adenosine receptor ligands for the A1 adenosine receptor. Glycolipids are accordingly useful in diagnostic and therapeutic methods that require the delivery or administration of A1 adenosine ligands.  
IT 53296-10-9, CV 1808  
RL: BSU (Biological study, unclassified); DGN (Diagnostic use); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(A1 adenosine receptor ligand; methods and formulations for increasing the affinity of A1 adenosine receptor ligands for A1 adenosine receptor using glycolipids in relation to diagnostic and therapeutic uses)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.




OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)  
RE.CNT 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 47 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 2002:623386 CAPLUS  
DN 138:51514  
TI Adenosine A2A receptor agonists: CoMFA-based selection of the most predictive conformation  
AU Doytchinova, I.; Valkova, I.; Natcheva, R.  
CS Department of Chemistry, Faculty of Pharmacy, Medical University-Sofia, Sofia, 1000, Bulg.  
SO SAR and QSAR in Environmental Research (2002), 13(2), 227-235  
CODEN: SQERED; ISSN: 1062-936X  
PB Taylor & Francis Ltd.  
DT Journal  
LA English  
AB A step-wise comparative mol. field anal. (CoMFA)-based procedure was applied to a series of 51 2-oxyadenosines in order to select the most predictive conformation for binding to A2A adenosine receptor (AR). The highest correlation and predictive power were found for conformers with side chain at 2nd position oriented in the direction opposite to the exocyclic amino group on the adenine ring (torsion N1C2OR = 120°) and fully extended. The interaction of ligand and receptor is under steric and electrostatic control. The steric contribution is of a greater importance for the predictivity than the electrostatic one. Hydrophobicity of the compds. investigated does not affect significantly either the affinity to A2A AR, nor the predictivity of the models.  
IT 50257-82-4 50257-84-6 50257-85-7  
50257-89-1 50257-95-9 131865-78-6  
131865-81-1 131865-82-2 131933-15-8  
131933-17-0 131933-20-5 131933-26-1  
131933-27-2 131973-26-7 137817-83-5  
137817-84-6 145747-87-1 194154-31-9  
194154-32-0  
RL: BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)  
(step-wise comparative mol. field anal. of 2-oxyadenosine derivs. conformation and binding to adenosine A2A adenosine receptor)  
RN 50257-82-4 CAPLUS  
CN Adenosine, 2-phenoxy- (CA INDEX NAME)

McIntosh


10/598,520

Absolute stereochemistry.




RN 50257-84-6 CAPLUS  
CN Adenosine, 2-butoxy- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 50257-85-7 CAPLUS  
CN Adenosine, 2-(pentyloxy)- (CA INDEX NAME)

Absolute stereochemistry.



RN 50257-89-1 CAPLUS  
CN Adenosine, 2-(2-hydroxyethoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 50257-95-9 CAPLUS  
CN Adenosine, 2-(hexyloxy)- (CA INDEX NAME)

Absolute stereochemistry.

McIntosh




10/598,520




RN 131865-78-6 CAPLUS  
CN Adenosine, 2-(phenylmethoxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 131865-81-1 CAPLUS  
CN Adenosine, 2-(4-phenylbutoxy)- (CA INDEX NAME)

Absolute stereochemistry.



RN 131865-82-2 CAPLUS  
CN Adenosine, 2-[(5-phenylpentyl)oxy]- (CA INDEX NAME)

Absolute stereochemistry.




RN 131933-15-8 CAPLUS  
CN Adenosine, 2-(3-methylbutoxy)- (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 131933-17-0 CAPLUS  
CN Adenosine, 2-(cyclohexylmethoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 131933-20-5 CAPLUS  
CN Adenosine, 2-(4-cyclohexylbutoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 131933-26-1 CAPLUS  
CN Adenosine, 2-(2-ethylbutoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 131933-27-2 CAPLUS  
CN Adenosine, 2-[(3-ethylpentyl)oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 131973-26-7 CAPLUS  
CN Adenosine, 2-[(4-methylpentyl)oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 137817-83-5 CAPLUS  
CN Adenosine, 2-[(2R)-2-phenylbutoxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 137817-84-6 CAPLUS  
CN Adenosine, 2-[(2S)-2-phenylbutoxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 145747-87-1 CAPLUS  
CN Adenosine, 2-[(3S)-3-phenylbutoxy]- (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 194154-31-9 CAPLUS  
CN Adenosine, 2-[[[(3S)-3,7-dimethyl-6-octenyl]oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 194154-32-0 CAPLUS  
CN Adenosine, 2-[[[(3R)-3,7-dimethyl-6-octenyl]oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 6 THERE ARE 6 CAPLUS RECORDS THAT CITE THIS RECORD (6 CITINGS)  
RE.CNT 17 THERE ARE 17 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 48 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 2002:290506 CAPLUS  
DN 137:28514  
TI Tonic activity of the rat adipocyte A1-adenosine receptor  
AU Liang, Hui-Xiu; Belardinelli, Luiz; Ozeck, Mark J.; Shryock, John C.  
CS Division of Cardiovascular Medicine, Department of Medicine, University of  
Florida, Gainesville, FL, 32610, USA  
SO British Journal of Pharmacology (2002), 135(6), 1457-1466  
CODEN: BJPCBM; ISSN: 0007-1188  
PB Nature Publishing Group  
DT Journal  
LA English  
AB Adipocyte A1-adenosine receptors (A1 AdoR) tonically inhibit adenylyl cyclase and lipolysis. Three potential explanations for tonic activity of A1AdoR of rat epididymal adipocytes were investigated: high affinity of adenosine for the receptor, efficient coupling of receptor activation to response, and spontaneous activity of the receptor in the absence of agonist. The affinity of adenosine for the adipocyte A1AdoR was determined as 4.6  $\mu$ M by anal. of effects of an irreversible receptor antagonist on agonist concentration-response relationships. In contrast, the potency of adenosine to decrease cAMP in isolated adipocytes was 1.4 nM. Occupancy by agonist of the A1AdoR was efficiently coupled to functional response (decrease of adipocyte cAMP content). Activation by adenosine of less

McIntosh

than 1% of A1AdoRs caused a near-maximal decrease of cAMP in adipocytes. Thus the receptor reserve for adenosine to decrease cAMP content of adipocytes was greater than 99%. Affinities and receptor reserves for other A1AdoR agonists were determined. Agonists appeared to differ more in their affinity for the receptor than in their intrinsic efficacy to activate it. A1AdoRs were inactive in the absence of agonist. It is concluded that adipocyte A1AdoR are tonically activated by endogenous adenosine at nanomolar concns. The expression of a high d. of A1AdoR that are efficiently coupled to a functional response enables the adipocyte to respond with high sensitivity to the low-affinity agonist, adenosine. Adipocytes may be a model for cells whose functions are tonically modulated by adenosine present in the interstitium of well-oxygenated tissues.

IT 53296-10-9, 2-Phenylaminoadenosine

RL: BSU (Biological study, unclassified); BIOL (Biological study)


(A1 adenosine receptor agonist; tonic activity of rat adipocyte

A1-adenosine receptor in regulation of adenylate cyclase and lipolysis)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 16 THERE ARE 16 CAPLUS RECORDS THAT CITE THIS RECORD (16 CITINGS)

RE.CNT 42 THERE ARE 42 CITED REFERENCES AVAILABLE FOR THIS RECORD

ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 49 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 2001:780668 CAPLUS

DN 135:335153

TI Treatment of neurodegenerative disease

IN Bamdad, R. Shoshanna; Bamdad, Cynthia C.

PA Minerva Biotechnologies Corporation, USA

SO PCT Int. Appl., 82 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2001078709	A2	20011025	WO 2001-US12484	20010412
	WO 2001078709	A3	20030417		
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW			
	RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
	CA 2404858	A1	20011025	CA 2001-2404858	20010412
	US 20030060487	A1	20030327	US 2001-835099	20010412
	EP 1328261	A2	20030723	EP 2001-927116	20010412
	R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			
	JP 2003530432	T	20031014	JP 2001-576010	20010412
PRAI	US 2000-196497P	P	20000412		
	US 2000-214221P	P	20000623		
	US 2000-248890P	P	20001115		
	WO 2001-US12484	W	20010412		

## ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OS MARPAT 135:335153

AB The invention relates to treatments for peptide aggregation associated with disease states such as neurodegenerative disease, particularly physiologic disease associated with Alzheimer's Disease, and non-neurodegenerative disease aggregation. Other aspects of the invention also provides a variety of novel assays for screening candidate drugs. Yet another aspects of the present invention also provides a series of compounds useful for treatment of neurologic disease as determined from these assays. These compounds can be packaged in kits. Other aspects of the invention also relate to the use of these compounds for the treatment and/or prevention of patients susceptible to or exhibiting of a disease characteristic of fibril formation or aberrant protein aggregation. Examples are given for monitoring drug activity as a function of time for drug profiling and cell-based screening assay for candidate drugs for affecting aggregate formation at a variety of stages of biochemical progression.


IT 53296-10-9, 2-Phenylaminoadenosine

RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(treatment of neurodegenerative disease)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)  
RE.CNT 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 50 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 2001:657146 CAPLUS

DN 136:175

TI CoMFA study on adenosine A2A receptor agonists

AU Doytchinova, Irini; Valkova, Iva; Natcheva, Roumiana

CS Department of Chemistry, Faculty of Pharmacy, Medical University - Sofia,  
Sofia, 1000, Bulg.

SO Quantitative Structure-Activity Relationships (2001), 20(2), 124-129  
CODEN: QSARDI; ISSN: 0931-8771

PB Wiley-VCH Verlag GmbH

DT Journal

LA English

AB A step-wise CoMFA-based procedure was applied to a series of 51 C2-oxyadenosines to select the most predictive conformation for binding to A2A adenosine receptor. The highest correlation and predictive power was found for conformers with the side chain at the 2-position oriented in the direction opposite to the exocyclic amino group on the adenine ring (torsion N1C2OR = 120°) and fully extended. The interaction of ligand and receptor is under steric and electrostatic control. The steric contribution is of greater importance for the predictivity than the electrostatic one. Hydrophobicity of the compounds investigated does not affect significantly either the affinity to A2A adenosine receptor, nor the predictivity of the models.

IT 50257-82-4 50257-84-6 50257-85-7  
50257-89-1 50257-95-9 131865-78-6  
131865-81-1 131865-82-2 131933-15-8  
131933-17-0 131933-20-5 131933-26-1  
131933-27-2 131973-26-7 137817-83-5  
137817-84-6 145747-87-1 194154-31-9  
194154-32-0


RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(CoMFA study on adenosine A2A receptor agonists)

RN 50257-82-4 CAPLUS

10/598,520

CN Adenosine, 2-phenoxy- (CA INDEX NAME)


Absolute stereochemistry.



RN 50257-84-6 CAPLUS

CN Adenosine, 2-butoxy- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 50257-85-7 CAPLUS

CN Adenosine, 2-(pentyloxy)- (CA INDEX NAME)


Absolute stereochemistry.



RN 50257-89-1 CAPLUS

CN Adenosine, 2-(2-hydroxyethoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 50257-95-9 CAPLUS

CN Adenosine, 2-(hexyloxy)- (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 131865-78-6 CAPLUS  
CN Adenosine, 2-(phenylmethoxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 131865-81-1 CAPLUS  
CN Adenosine, 2-(4-phenylbutoxy)- (CA INDEX NAME)

Absolute stereochemistry.



RN 131865-82-2 CAPLUS  
CN Adenosine, 2-[(5-phenylpentyl)oxy]- (CA INDEX NAME)

Absolute stereochemistry.




RN 131933-15-8 CAPLUS  
CN Adenosine, 2-(3-methylbutoxy)- (CA INDEX NAME)

Absolute stereochemistry.

McIntosh




10/598,520




RN 131933-17-0 CAPLUS  
CN Adenosine, 2-(cyclohexylmethoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 131933-20-5 CAPLUS  
CN Adenosine, 2-(4-cyclohexylbutoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 131933-26-1 CAPLUS  
CN Adenosine, 2-(2-ethylbutoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 131933-27-2 CAPLUS  
CN Adenosine, 2-[(3-ethylpentyl)oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 131973-26-7 CAPLUS  
CN Adenosine, 2-[(4-methylpentyl)oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 137817-83-5 CAPLUS  
CN Adenosine, 2-[(2R)-2-phenylbutoxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 137817-84-6 CAPLUS  
CN Adenosine, 2-[(2S)-2-phenylbutoxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 145747-87-1 CAPLUS  
CN Adenosine, 2-[(3S)-3-phenylbutoxy]- (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 194154-31-9 CAPLUS  
CN Adenosine, 2-[[[(3S)-3,7-dimethyl-6-octenyl]oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 194154-32-0 CAPLUS  
CN Adenosine, 2-[[[(3R)-3,7-dimethyl-6-octenyl]oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 9 THERE ARE 9 CAPLUS RECORDS THAT CITE THIS RECORD (9 CITINGS)  
RE.CNT 17 THERE ARE 17 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 51 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 2001:597739 CAPLUS  
DN 135:162508  
TI Adenosine A2a receptor antagonist for treating and preventing hepatic  
fibrosis, cirrhosis and fatty liver  
IN Cronstein, Bruce N.; Chan, Edwin  
PA New York University, USA  
SO PCT Int. Appl., 48 pp.  
CODEN: PIXXD2  
DT Patent  
LA English  
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2001058241	A2	20010816	WO 2001-US4341	20010212
	WO 2001058241	A9	20021017		
	W: AU, CA, JP				
	RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR				
	CA 2398908	A1	20010816	CA 2001-2398908	20010212
	CA 2398908	C	20091215		
	AU 2001038124	A	20010820	AU 2001-38124	20010212
	US 20020002145	A1	20020103	US 2001-780365	20010212

McIntosh

US 6555545	B2	20030429		
JP 2004502640	T	20040129	JP 2001-557366	20010212
AU 2001238124	B2	20060525	AU 2001-238124	20010212
EP 1272897	B1	20080507	EP 2001-910529	20010212
R: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE, TR				
AT 394104	T	20080515	AT 2001-910529	20010212
PT 1272897	E	20080818	PT 2001-910529	20010212
ES 2307593	T3	20081201	ES 2001-910529	20010212
AU 2006203699	A1	20060921	AU 2006-203699	20060825
AU 2006203699	B2	20100204		
PRAI US 2000-181546P	P	20000210		
AU 2001-238124	A3	20010212		
WO 2001-US4341	W	20010212		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT


AB Adenosine A2a receptor antagonists such as CGS-21680 or adenosine derivs are used for treating and preventing hepatic fibrosis, cirrhosis and fatty liver. The adenosine A2a receptor antagonist CGS-21680 increased collagen production by rHSC.

IT 53296-10-9, 2-Phenylaminoadenosine  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (adenosine A2a receptor antagonists for treating and preventing hepatic fibrosis, cirrhosis and fatty liver)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD (3 CITINGS)

L6 ANSWER 52 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 2001:39215 CAPLUS

DN 134:216814

TI Design, Synthesis, and Evaluation of Novel A2A Adenosine Receptor Agonists

AU Rieger, Jayson M.; Brown, Milton L.; Sullivan, Gail W.; Linden, Joel; Macdonald, Timothy L.

CS Departments of Chemistry and Medicine, University of Virginia, Charlottesville, VA, 22904-4319, USA

SO Journal of Medicinal Chemistry (2001), 44(4), 531-539

CODEN: JMCMAR; ISSN: 0022-2623

PB American Chemical Society

DT Journal

LA English

OS CASREACT 134:216814

AB The authors have been interested in the design, synthesis, and evaluation of novel adenosine A2A agonists. Through the use of comparative mol. field anal. (CoMFA) the authors have generated a training model that includes 78 structurally diverse A2A agonists and correlated their affinity for isolated rat brain receptors with differences in their structural and electrostatic properties. The authors validated this model by predicting the activity of a test set that included 24 addnl. A2A agonists. Our CoMFA model, which incorporates the physiochem. property of dipole and selects against A1 receptor activity, generated a correlated final model ( $r^2 = 0.891$ ) that provides for enhanced A2A selectivity and predictability. Synthesis, pharmacol. evaluation, and modeling of four novel ligands further validate the utility and predictive power ( $r^2 = 0.626$ ) of the CoMFA model.

IT 50257-82-4 50257-84-6 50257-89-1  
 50257-95-9 131865-78-6 131865-81-1  
 131865-82-2 131933-17-0 131933-20-5  
 131933-26-1 131933-27-2 131973-26-7

10/598,520


137817-84-6 145747-87-1

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)  
(design and synthesis and evaluation of novel A2A adenosine receptor agonists)

RN 50257-82-4 CAPLUS

CN Adenosine, 2-phenoxy- (CA INDEX NAME)


Absolute stereochemistry.



RN 50257-84-6 CAPLUS

CN Adenosine, 2-butoxy- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 50257-89-1 CAPLUS

CN Adenosine, 2-(2-hydroxyethoxy)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 50257-95-9 CAPLUS

CN Adenosine, 2-(hexyloxy)- (CA INDEX NAME)

Absolute stereochemistry.




McIntosh

10/598,520


RN 131865-78-6 CAPLUS  
CN Adenosine, 2-(phenylmethoxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 131865-81-1 CAPLUS  
CN Adenosine, 2-(4-phenylbutoxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 131865-82-2 CAPLUS  
CN Adenosine, 2-[(5-phenylpentyl)oxy]- (CA INDEX NAME)

Absolute stereochemistry.



RN 131933-17-0 CAPLUS  
CN Adenosine, 2-(cyclohexylmethoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 131933-20-5 CAPLUS  
CN Adenosine, 2-(4-cyclohexylbutoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 131933-26-1 CAPLUS  
CN Adenosine, 2-(2-ethylbutoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 131933-27-2 CAPLUS  
CN Adenosine, 2-[(3-ethylpentyl)oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 131973-26-7 CAPLUS  
CN Adenosine, 2-[(4-methylpentyl)oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 137817-84-6 CAPLUS  
CN Adenosine, 2-[(2S)-2-phenylbutoxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520



RN 145747-87-1 CAPLUS  
CN Adenosine, 2-[(3S)-3-phenylbutoxy]- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 64 THERE ARE 64 CAPLUS RECORDS THAT CITE THIS RECORD (64 CITINGS)  
RE.CNT 33 THERE ARE 33 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 53 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 2000:573631 CAPLUS  
DN 133:182707  
TI Hair growth stimulants containing purinoceptor stimulants and their  
screening method  
IN Nakaya, Yutaka; Arase, Seiji; Imamura, Koji  
PA Taisho Pharmaceutical Co., Ltd., Japan  
SO PCT Int. Appl., 62 pp.  
CODEN: PIXXD2  
DT Patent  
LA Japanese  
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2000047172	A1	20000817	WO 2000-JP694	20000209
	W: AU, CA, CN, KR, US				
	RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
	JP 2000297015	A	20001024	JP 2000-34181	20000210
	JP 4465779	B2	20100519		
	JP 2009142281	A	20090702	JP 2009-423	20090105
PRAI	JP 1999-33502	A	19990210		
	JP 1999-33503	A	19990210		
	JP 1999-33504	A	19990210		
	JP 1999-33505	A	19990210		
	JP 2000-34181	A3	20000210		

OS MARPAT 133:182707

AB Disclosed are excellent hair growth stimulants having novel function mechanisms different from the conventional hair growth stimulants and a method for screening the same. The hair growth stimulants contain as the active ingredient compds. exerting an effect of stimulating purine receptors (adenosine receptor, ATP receptor, etc.), an effect of potentiating the above effect, and an effect of liberating compds. having an effect of stimulating purine receptors (adenosine, adenosine derivs., adenosine metabolites, etc.) from cells. The screening method comprises adding a test substance to cells which have been transformed with an ABC transporter gene and a purine derivative receptor gene and using the calcium influx at this point as an indication. A hair gel containing N6-(L-2-phenylisopropyl)adenosine 0.5, polyethyleneglycol monostearate 1, 1,3-butylene glycol 7, carboxyvinyl polymer 1.5, diisopropanol amine q.s.,


McIntosh



10/598,520

ethanol 40, and water q.s. to 100 % was prepared  
IT 53296-10-9, 2-Phenylaminoadenosine  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(hair growth stimulants containing purinoceptor stimulants and their screening method)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD (5 CITINGS)  
RE.CNT 40 THERE ARE 40 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 54 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 2000:405019 CAPLUS  
DN 133:115443

TI Thermodynamically distinct high and low affinity states of the A1 adenosine receptor induced by G protein coupling and guanine nucleotide ligation states of G proteins

AU Lorenzen, Anna; Guerra, Laura; Campi, Franca; Lang, Heidrun; Schwabe, Ulrich; Borea, Pier Andrea

CS Pharmakologisches Institut der Universitat Heidelberg, Heidelberg, D-69120, Germany

SO British Journal of Pharmacology (2000), 130(3), 595-604  
CODEN: BJPCBM; ISSN: 0007-1188

PB Nature Publishing Group

DT Journal

LA English

AB 1 The influence of the receptor-G protein coupling state and the guanine nucleotide ligation state of the G protein on the binding mechanism of A1 adenosine receptor ligands has been investigated in [3H]-1,3-dipropyl-8-cyclopentylxanthine ([3H]-DPCPX) binding studies in rat brain membranes. Thermodyn. parameters of binding of A1 adenosine receptor ligands of different intrinsic activities were determined in the absence or presence of GDP and compared to the binding mechanism after receptor-G protein uncoupling. 2 In agreement with previous studies, it was found that xanthine and non-xanthine antagonists showed an enthalpy- or enthalpy- and entropy-driven binding mechanism under all conditions. 3 In contrast to antagonists, the binding mechanism of agonists was strongly affected by the G protein coupling state or the absence or presence of guanine nucleotides. Binding of full and partial agonists to the high-affinity state of the A1 receptor was entropy-driven in the absence of GDP, and a good correlation between intrinsic activities and the contribution of entropy was observed. In the absence of GDP, binding of full and partial agonists and antagonists to the high affinity state of the receptor was thermodynamically discriminated. In contrast, no such discrimination was found in the presence of GDP. 4 The binding mechanism of agonists to the low-affinity state of the receptor was identical to that of antagonists only after uncoupling of the receptor from G proteins by pretreatment with N-ethylmaleimide or guanosine-5'-( $\gamma$ -thio)-triphosphate (GTP $\gamma$ S). 5 These results indicate the existence of two thermodynamically distinct high- and low-affinity states of the A1 adenosine receptor.

IT 53296-10-9, CV 1808


RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
(thermodynamically distinct high and low affinity states of A1 adenosine receptor induced by G protein coupling and guanine nucleotide

McIntosh

10/598,520

ligation states of G proteins)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 13 THERE ARE 13 CAPLUS RECORDS THAT CITE THIS RECORD (13 CITINGS)  
RE.CNT 26 THERE ARE 26 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 55 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 2000:258227 CAPLUS

DN 133:37724

TI Molecular modeling of 2-alkyloxy- and 2-aralkyloxy-adenosine A1- and A2-agonists

AU Matova, Mariana M.; Nacheva, Rumiana N.; Boicheva, Sirma V.

CS Department of Chemistry and Biochemistry, Faculty of Medicine, Medical University, Sofia, 1431, Bulg.

SO Drug Design and Discovery (2000), 16(4), 255-270, 5 plates

CODEN: DDDIEV; ISSN: 1055-9612

PB Harwood Academic Publishers

DT Journal

LA English

AB The C2-region of adenosine A1- and A2-receptors by a mol. modeling technique has been extended and applied to a series of 2-substituted adenosines reported by Olsson, et al. The similarity and dissimilarity of the structure maps obtained by mol. modeling have been used as a basis for the mapping of the analyzed receptor domain. The proposed model of the C2-region of the A1-receptor consists of a narrow and sterically limited area that interacts well electrostatically with small and electron rich moieties. Olsson's provisional model of the C2-region of the A2-receptor has been extended with two subsites, as well as with a forbidden area near the C2-position of the purine ring. The conformational anal. performed in the study does not support the hypothesis of Olsson et al. that adenosine C2 substituents may partly occupy the same receptor domain as the N6 substituents of the A1-receptor. The occupation of the cycloalkyl subsite increases the receptor selectivity while the occupation of the other subsite by aryl rings, fixed at a parallel position to the purine system, highly enhances the receptor affinity.

IT 50257-82-4 50257-84-6 50257-85-7  
50257-89-1 50257-95-9 131865-78-6  
131865-81-1 131865-82-2 131933-15-8  
131933-17-0 131933-20-5 131933-26-1  
131933-27-2 131973-26-7 137817-83-5  
137817-84-6 145747-87-1 194154-31-9  
194154-32-0


RL: BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)  
(mol. modeling of 2-alkyloxy- and 2-aralkyloxy-adenosine A1- and A2-agonists)

RN 50257-82-4 CAPLUS

CN Adenosine, 2-phenoxy- (CA INDEX NAME)


Absolute stereochemistry.

10/598,520




RN 50257-84-6 CAPLUS  
CN Adenosine, 2-butoxy- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 50257-85-7 CAPLUS  
CN Adenosine, 2-(pentyloxy)- (CA INDEX NAME)

Absolute stereochemistry.



RN 50257-89-1 CAPLUS  
CN Adenosine, 2-(2-hydroxyethoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 50257-95-9 CAPLUS  
CN Adenosine, 2-(hexyloxy)- (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 131865-78-6 CAPLUS  
CN Adenosine, 2-(phenylmethoxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 131865-81-1 CAPLUS  
CN Adenosine, 2-(4-phenylbutoxy)- (CA INDEX NAME)

Absolute stereochemistry.



RN 131865-82-2 CAPLUS  
CN Adenosine, 2-[(5-phenylpentyl)oxy]- (CA INDEX NAME)

Absolute stereochemistry.




RN 131933-15-8 CAPLUS  
CN Adenosine, 2-(3-methylbutoxy)- (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 131933-17-0 CAPLUS  
CN Adenosine, 2-(cyclohexylmethoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 131933-20-5 CAPLUS  
CN Adenosine, 2-(4-cyclohexylbutoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 131933-26-1 CAPLUS  
CN Adenosine, 2-(2-ethylbutoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 131933-27-2 CAPLUS  
CN Adenosine, 2-[(3-ethylpentyl)oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 131973-26-7 CAPLUS  
CN Adenosine, 2-[(4-methylpentyl)oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 137817-83-5 CAPLUS  
CN Adenosine, 2-[(2R)-2-phenylbutoxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 137817-84-6 CAPLUS  
CN Adenosine, 2-[(2S)-2-phenylbutoxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 145747-87-1 CAPLUS  
CN Adenosine, 2-[(3S)-3-phenylbutoxy]- (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 194154-31-9 CAPLUS  
CN Adenosine, 2-[[[(3S)-3,7-dimethyl-6-octenyl]oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 194154-32-0 CAPLUS  
CN Adenosine, 2-[[[(3R)-3,7-dimethyl-6-octenyl]oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)  
RE.CNT 28 THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 56 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1999:819059 CAPLUS  
DN 132:35992  
TI Preparation of novel arabinosyladenine derivatives having a resistance to  
the metabolism by ADA and antiviral action  
IN Yamada, Toshio; Yamanishi, Koichi  
PA Nippon Zoki Pharmaceutical Co., Ltd., Japan  
SO Eur. Pat. Appl., 19 pp.  
CODEN: EPXXDW  
DT Patent  
LA English  
FAN.CNT 1


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	EP 967220	A1	19991229	EP 1999-111945	19990623
	EP 967220	B1	20030409		
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
	JP 2000007695	A	20000111	JP 1998-177209	19980624
	JP 3619017	B2	20050209		
	KR 2000006339	A	20000125	KR 1999-23448	19990622
	AT 236923	T	20030415	AT 1999-111945	19990623
	TW 564249	B	20031201	TW 1999-110523	19990623

McIntosh

10/598,520

ES 2196676	T3	20031216	ES 1999-111945	19990623
CN 1242373	A	20000126	CN 1999-108908	19990624
CN 1146573	C	20040421		
US 6242429	B1	20010605	US 1999-339257	19990624

PRAI JP 1998-177209 A 19980624  
ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT  
OS MARPAT 132:35992  
GI




AB Novel 2-substituted arabinosyladenine derivs. I (where Z = an alkyl group having more than 4 carbon atoms, alkenyl or alkynyl; R = hydrogen or lower alkyl group) are prepared. For example, 5-amino-1-( $\beta$ -D-arabinofuranosyl)-4-cyanoimidazole is made to react with a nitrile of the formula Z-CN (where Z = an alkyl group having more than 4 carbon atoms, alkenyl or alkynyl) to yield a 2-substituted Ara-A derivative. The compds. of the present invention have a resistance to the metabolism by ADA and are useful as therapeutic or preventive agents for diseases infected by DNA virus such as herpes simplex virus (HSV), herpes zoster virus, cytomegalovirus (CMV), adenovirus, hepatitis virus or vaccinia virus.

IT 252552-21-9P 252552-23-1P  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)  
(preparation of novel arabinosyladenine derivs. having a resistance to the metabolism by ADA and antiviral action)

RN 252552-21-9 CAPLUS  
CN 9H-Purin-6-amine, 9- $\beta$ -D-arabinofuranosyl-2-octyl- (CA INDEX NAME)


Absolute stereochemistry.



RN 252552-23-1 CAPLUS  
CN 9H-Purin-6-amine, 9- $\beta$ -D-arabinofuranosyl-2-dodecyl- (CA INDEX NAME)

Absolute stereochemistry.





OSC.G 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD (4 CITINGS)  
 RE.CNT 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 57 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1999:771170 CAPLUS

DN 132:102410

TI Molecular Recognition of Modified Adenine Nucleotides by the  
 P2Y1-Receptor. 1. A Synthetic, Biochemical, and NMR Approach

AU Halbfinger, Efrat; Major, Dan T.; Ritzmann, Marco; Ubl, Joachim; Reiser,  
 Georg; Boyer, Jose L.; Harden, Kendall T.; Fischer, Bilha

CS Department of Chemistry Gonda-Goldschmied Center, Bar-Ilan University,  
 Ramat-Gan, 52900, Israel

SO Journal of Medicinal Chemistry (1999), 42(26), 5325-5337

CODEN: JMCMAR; ISSN: 0022-2623

PB American Chemical Society

DT Journal

LA English

AB The remarkably high potencies of 2-thioether-adenine nucleotides regarding  
 the activation of the P2Y1-receptor (P2Y1-R) in turkey erythrocyte  
 membranes represent some of the largest substitution-promoted increases in  
 potencies over that of a natural receptor ligand. This paper describes  
 the investigation regarding the origin of the high potency of these P2Y1-R  
 ligands over that of ATP. For this study, an integrated approach was  
 employed combining the synthesis of new ATP analogs, their biochem.  
 evaluation, and their SAR anal. involving NMR expts. and theor. calcns.  
 These expts. and calcns. were performed to elucidate the conformation and  
 to evaluate the electronic nature of the investigated P2Y1-R ligands. ATP  
 analogs synthesized included derivs. where C2 or C8 positions were  
 substituted with electron-donating groups such as ethers, thioethers, or  
 amines. The compds. were tested for their potency to induce  
 P2Y1-R-mediated activation of phospholipase C in turkey erythrocytes and  
 Ca2+ response in rat astrocytes. 8-Substituted ATP and AMP derivs. had  
 little or no effect on phospholipase C or on calcium levels, whereas the  
 corresponding 2-substituted ATP analogs potentially increased the levels of  
 inositol phosphates and [Ca2+]i. AMP analogs were ineffective except for  
 2-butylthio-AMP which induced a small Ca2+ response. P2Y1-R activity of  
 these compds. was demonstrated by testing these ligands also on NG108-15  
 neuroblastoma + glioma hybrid cells. NMR data together with theor.  
 calcns. imply that steric, rather than electronic, effects play a major  
 role in ligand binding to the P2Y1-R. Hydrophobic interactions and  
 H-bonds of the C2 substituent appear to be important determinants of a  
 P2Y1-R ligand affinity.

IT 50257-84-6P, 2-(Butoxy)adenosine

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
 (Reactant or reagent)


(preparation and structure activity relations of modified adenine  
 nucleotides as P2Y1 receptor agonists)

RN 50257-84-6 CAPLUS

CN Adenosine, 2-butoxy- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


10/598,520



OSC.G 35 THERE ARE 35 CAPLUS RECORDS THAT CITE THIS RECORD (36 CITINGS)  
RE.CNT 51 THERE ARE 51 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 58 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1999:173299 CAPLUS  
DN 130:276244  
TI Using theoretical descriptors in a correlation analysis of adenosine activity  
AU Famini, George R.; Loubnev, Valery P.; Frykman, Eric K.; Wilson, Leland Y.  
CS Development Engineering Center, Edgewood Research, Aberdeen, MD, 21010, USA  
SO Quantitative Structure-Activity Relationships (1998), 17(6), 558-564  
CODEN: QSARDI; ISSN: 0931-8771  
PB Wiley-VCH Verlag GmbH  
DT Journal  
LA English  
AB A theor. linear solvation energy relationship (TLSE) is used as a model for relating 2 guinea pig heart muscle activities to a set of computationally derived mol. descriptors for a set of 24 2-alkoxy and 25 2-aryloxy adenosines. The resulting equations are consistent with the structure activity relationship (SAR) study showing an increase in activity at 1 site with increase in substituent size and a hydrophobicity index.  
IT 50257-82-4 50257-84-6 50257-85-7  
50257-89-1 50257-95-9 131865-78-6  
131865-81-1 131865-82-2 131933-15-8  
131933-17-0 131933-20-5 131933-26-1  
131933-27-2 131973-26-7  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(using theor. descriptors in a correlation anal. of adenosine activity)  
RN 50257-82-4 CAPLUS  
CN Adenosine, 2-phenoxy- (CA INDEX NAME)

Absolute stereochemistry.




RN 50257-84-6 CAPLUS  
CN Adenosine, 2-butoxy- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 50257-85-7 CAPLUS  
CN Adenosine, 2-(pentyloxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 50257-89-1 CAPLUS  
CN Adenosine, 2-(2-hydroxyethoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 50257-95-9 CAPLUS  
CN Adenosine, 2-(hexyloxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 131865-78-6 CAPLUS  
CN Adenosine, 2-(phenylmethoxy)- (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 131865-81-1 CAPLUS  
CN Adenosine, 2-(4-phenylbutoxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 131865-82-2 CAPLUS  
CN Adenosine, 2-[(5-phenylpentyl)oxy]- (CA INDEX NAME)

Absolute stereochemistry.



RN 131933-15-8 CAPLUS  
CN Adenosine, 2-(3-methylbutoxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 131933-17-0 CAPLUS  
CN Adenosine, 2-(cyclohexylmethoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 131933-20-5 CAPLUS  
CN Adenosine, 2-(4-cyclohexylbutoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 131933-26-1 CAPLUS  
CN Adenosine, 2-(2-ethylbutoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 131933-27-2 CAPLUS  
CN Adenosine, 2-[(3-ethylpentyl)oxy]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 131973-26-7 CAPLUS  
CN Adenosine, 2-[(4-methylpentyl)oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

McIntosh




OSC.G 10 THERE ARE 10 CAPLUS RECORDS THAT CITE THIS RECORD (10 CITINGS)  
 RE.CNT 33 THERE ARE 33 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 59 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
 AN 1999:27719 CAPLUS  
 DN 130:90521  
 TI Methods for the inhibition of neuronal activity and treatment of pain  
 syndromes or epilepsy by local delivery of adenosine  
 IN Mohler, Hanns; Boison, Detlev  
 PA Switz.  
 SO PCT Int. Appl., 43 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9858653	A1	19981230	WO 1998-IB973	19980623
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG US 6110902 A 20000829 US 1997-881038 19970623 AU 9876711 A 19990104 AU 1998-76711 19980623 EP 996453 A1 20000503 EP 1998-924521 19980623 EP 996453 B1 20040428 R: CH, DE, GB, LI PRAI US 1997-881038 A 19970623 WO 1998-IB973 W 19980623				
AB The invention relates to the treatment of conditions associated with neuronal activity. Specifically, the invention is drawn to methods and compns. for administering adenosine to inhibit pain syndromes or epilepsy in a patient. IT 53296-10-9, 2-Phenylaminoadenosine RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (adenosine local delivery method for inhibition of neuronal activity and treatment of pain syndromes or epilepsy) RN 53296-10-9 CAPLUS CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)				


Absolute stereochemistry.



RE.CNT 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 60 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1998:810073 CAPLUS  
DN 130:177447  
TI Differences in the order of potency for agonists but not antagonists at human and rat adenosine A2A receptors  
AU Kull, Bjorn; Arslan, Guilia; Nilsson, Christer; Owman, Christer; Lorenzen, Anna; Schwabe, Ulrich; Fredholm, Bertil B.  
CS Department of Physiology and Pharmacology, Section of Molecular Neuropharmacology, Karolinska Institutet, Stockholm, S-171 77, Swed.  
SO Biochemical Pharmacology (1999), 57(1), 65-75  
CODEN: BCPA6; ISSN: 0006-2952  
PB Elsevier Science Inc.  
DT Journal  
LA English  
AB To examine possible species differences in pharmacol., rat adenosine A2A receptors were studied in PC12 (pheochromocytoma) cells, and human receptors in Chinese hamster ovary (CHO) cells transfected with the cloned human A2A receptor cDNA. Using [3H]-5-amino-7-(2-phenylethyl)-2-(2-furyl)pyrazolo[4,3-e]-1,2,4-triazolo[1,5-c]pyrimidine ([3H]-SCH 58261) as radioligand, the estimated Bmax (maximal binding) was 538 and 2085 fmol/mg in CHO and PC12 cells, resp. The Kd (dissociation constant) values for [3H]-SCH 58261 were 1.05 and 5.6 nM in the 2 cell types, resp. The order of potency of antagonists and most agonists was the same in both cell types, but 2-phenylaminoadenosine and 2-chloroadenosine were relatively less potent in PC12 cells than in CHO cells. In the functional assay, using cAMP accumulation, all agonists tested were more potent in CHO than in PC12 cells, but this could not be readily explained by differences in adenylyl cyclase or in the expression of G proteins. As in the case of binding, the relative agonist potencies were similar for most compds., but 2-phenylaminoadenosine and 2-chloroadenosine were more potent at human A2A receptors in CHO cells than predicted from the data obtained on rat A2A receptors in PC12 cells. The antagonists were approx. equipotent in the 2 cells. These results show that, despite only small differences in receptor amino acid sequences and no difference in antagonist pharmacol., the relative order of potency of receptor agonists can differ between species homologues of the adenosine A2A receptor.  
IT 53296-10-9, CV 1808  
RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
(species differences in potency at human and rat adenosine A2A receptors of)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 30 THERE ARE 30 CAPLUS RECORDS THAT CITE THIS RECORD (30 CITINGS)  
RE.CNT 38 THERE ARE 38 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT


L6 ANSWER 61 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1998:789051 CAPLUS  
DN 130:29255  
TI Medicinal composition for prevention or treatment of hepatopathy  
IN Ozaki, Takayuki; Hirata, Yoshihisa; Tada, Shin-ichi  
PA Nippon Shinyaku Co., Ltd., Japan  
SO PCT Int. Appl., 26 pp.

10/598,520

CODEN: PIXXD2  
DT Patent  
LA Japanese  
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9852611	A1	19981126	WO 1998-JP2223	19980520
	W: AT, AU, BR, CA, CH, CN, DE, DK, ES, GB, HU, ID, IL, JP, KR, MX, NO, NZ, PT, RU, SE, UA, US, VN, AM, AZ, BY, KG, KZ, MD, TJ, TM				
	RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
	AU 9874495	A	19981211	AU 1998-74495	19980520
	EP 983768	A1	20000308	EP 1998-921742	19980520
	R: BE, CH, DE, ES, FR, GB, IT, LI, NL				
PRAI	JP 1997-133480	A	19970523		
	JP 1997-192555	A	19970717		
	WO 1998-JP2223	W	19980520		
OS	MARPAT 130:29255				
AB	A medicinal composition containing an adenosine A2 receptor agonist as an active ingredient, is effective in the prevention or treatment of hepatopathy. 1-[6-Amino-2-[[2-[4-(2-carboxyethyl)phenyl]ethyl]amino]-9H-purine-9-yl]-1-deoxy-N-ethyl-β-D-ribofuranuronamide was i.p. or orally administered to mice to show inhibitory activities against con A-induced liver damages. Tablet formulations containing the active ingredients are also provided.				
IT	53296-10-9, 2-(Phenylamino)adenosine				
	RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)				
	(adenosine A2 receptor agonists for treatment of hepatopathy)				
RN	53296-10-9 CAPLUS				
CN	Adenosine, 2-(phenylamino)- (CA INDEX NAME)				

Absolute stereochemistry.



OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)  
RE.CNT 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 62 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1998:727900 CAPLUS  
DN 130:90482  
TI Activation of various subtypes of G-protein  $\alpha$  subunits by partial agonists of the adenosine A1 receptor  
AU Lorenzen, Anna; Lang, Heidrun; Schwabe, Ulrich  
CS INSTITUTE OF PHARMACOLOGY, UNIVERSITY OF HEIDELBERG, HEIDELBERG, 69120, Germany  
SO Biochemical Pharmacology (1998), 56(10), 1287-1293  
CODEN: BCPA6; ISSN: 0006-2952  
PB Elsevier Science Inc.  
DT Journal  
LA English  
AB The activation of different G protein subtypes by the rat adenosine A1 receptor initiated by stimulation with the full agonist 2-chloro-N6-cyclopentyladenosine (CCPA) and by six structurally distinct partial agonists of this receptor was investigated. Endogenous G protein  $\alpha$  subunits in rat cortical membranes were inactivated by N-ethylmaleimide (NEM). Activation of rat recombinant myristoylated  $\alpha_0$ ,  $\alpha_{11}$ ,  $\alpha_{i2}$  and  $\alpha_{i3}$  by partial agonists in comparison to the full agonist was assessed by guanosine-5'-( $\gamma$ -[35S]thio)triphosphate ([35S]GTP $\gamma$ S) binding after reconstitution of G protein  $\alpha$  subunits with the adenosine A1

McIntosh



10/598,520

receptor in N-ethylmaleimide-treated membranes.  
2-Chloro-N6-cyclopentyladenosine and 3'-deoxy-N6-cyclopentyladenosine (3'-d-CPA), the partial agonist with the highest intrinsic activity, were significantly more potent in activation of  $\alpha_i$  subtypes than  $\alpha_o$ . In contrast, 5'-methylthioadenosine (MeSA), 2'-deoxy-2-chloroadenosine (cladribine), 2'-deoxy-N6-cyclopentyladenosine (2'-d-CPA), 2-phenylaminoadenosine (CV 1808) and C8-aminopropyl-N6-cyclopentyladenosine (C8-aminopropyl-CPA) did not exhibit higher potency for  $G_o$  or any  $G_i$  subtype. All partial agonists, although carrying structurally different modifications, showed higher relative intrinsic activities in activation of  $G_i$  than of  $G_o$ , indicating that  $G_i$ -coupled pathways may be activated selectively via the A1 receptor by partial agonists, but not  $G_o$ -mediated responses.


IT 53296-10-9, CV 1808

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)  
(activation of various subtypes of G-protein  $\alpha$  subunits by partial agonists of the adenosine A1 receptor)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 25 THERE ARE 25 CAPLUS RECORDS THAT CITE THIS RECORD (25 CITINGS)  
RE.CNT 33 THERE ARE 33 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 63 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1998:493732 CAPLUS  
DN 129:131238  
OREF 129:26693a,26696a  
TI Screening method for agents for treatment of eye disorders  
IN Trier, Klaus  
PA Aps, Klaus Trier, Den.  
SO PCT Int. Appl., 100 pp.  
CODEN: PIXXD2  
DT Patent  
LA English  
FAN.CNT 1


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9830900	A2	19980716	WO 1998-DK1	19980105
	WO 9830900	A3	19981210		
	W:	AL, AM, AT, AU, AZ, BA, BE, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW			
	RW:	GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG			
	CA 2276287	A1	19980716	CA 1998-2276287	19980105
	CA 2276287	C	20071030		
	AU 9853121	A	19980803	AU 1998-53121	19980105
	IN 1998CA00024	A	20051111	IN 1998-CA24	19980106
	US 6710051	B1	20040323	US 1999-341169	19990706
	US 20040013609	A1	20040122	US 2003-464750	20030619
PRAI	DK 1997-9	A	19970106		
	DK 1997-823	A	19970707		
	DK 1997-1383	A	19971201		
	WO 1998-DK1	W	19980105		

McIntosh

10/598,520

US 1999-341169 A3 19990706  
ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT  
OS MARPAT 129:131238  
AB A method is provided for identification of substances which are applicable for treatment or prevention of an insufficient longitudinal growth of the eye (hypermetropia) or for treatment or prevention of an excessive longitudinal growth of the eye (myopia); substances identified by the method for treating or preventing conditions related to the longitudinal growth of the eye; substances and mixts. of substances for the preparation of a pharmaceutical composition for the treatment or prevention of abnormal growth of the axial length of the eye. The identification involves measuring the effect of the substances on the retinal pigment epithelium of the eye, e.g. by detecting the metabolic effect of the substance on the retinal epithelium, the effect on the standing potential or the effect on the proteoglycans of the scleral tissue of the eye, by way of EOG examination, by way on the size of the so-called c-wave in ERG-recordings, or by the state of the Ca<sup>2+</sup>-channels or on the [3H]-ryanodine receptors of the retinal pigment epithelium.  
IT 53296-10-9, CV 1808  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(screening method for agents for treatment of eye disorders)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.




OSC.G 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)  
RE.CNT 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 64 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1998:441960 CAPLUS  
DN 129:109311  
OREF 129:22461a,22464a  
TI Preparation of nucleoside uronamides as A3 adenosine receptor agonists  
IN Jacobson, Kenneth A.; Gallo-Rodriguez, Carola; Van Galen, Philip J. M.; Von Lubitz, Dag K. J. E.; Jeong, Heaok Kim  
PA United States Dept. of Health and Human Services, USA  
SO U.S., 54 pp., Cont.-in-part of U. S. Ser. No. 163,324, abandoned.  
CODEN: USXXAM  
DT Patent  
LA English  
FAN.CNT 3

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	US 5773423	A	19980630	US 1994-274628	19940713
	US 5688774	A	19971118	US 1995-396111	19950228
PRAI	US 1993-91109	B2	19930713		
	US 1993-163324	B2	19931206		
	US 1994-274628	A2	19940713		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT  
OS MARPAT 129:109311  
GI

McIntosh



AB The present invention provides N6-benzyladenosine-5'-N-uronamide and related substituted compds. I (R1 = amide; R2 = halo, amino, alkenyl, alkynyl, thio, alkylthio; R3 = S-1-phenylethyl, Bn, phenylethyl), particularly those containing substituents on the benzyl and/or uronamide groups, and modified xanthine ribosides, as well as pharmaceutical compns. containing such compds. The present invention also provides a method of selectively activating an A3 adenosine receptor in a mammal, which method comprises acutely or chronically administering to a mammal in need of selective activation of its A3 adenosine receptor a therapeutically effective amount of a compound which binds with the A3 receptor so as to stimulate an A3 receptor-dependent response. Thus, N6-(3-iodobenzyl)adenosine was prepared tested for its affinity in binding at rat brain A1, A2, A3 adenosine receptors (Ki = 9.5-220.0 nM).


IT 53296-10-9P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(preparation of nucleoside uronamides as A3 adenosine receptor agonists)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 11 THERE ARE 11 CAPLUS RECORDS THAT CITE THIS RECORD (11 CITINGS)  
RE.CNT 35 THERE ARE 35 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 65 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1998:413029 CAPLUS

DN 129:145069

OREF 129:29475a,29478a

TI Pharmacological classification of adenosine receptors in the sinoatrial and atrioventricular nodes of the guinea pig

AU Meester, B. J.; Shankley, N. P.; Welsh, N. J.; Wood, J.; Meijler, F. L.; Black, J. W.

CS Rayne Institute, Analytical Pharmacology, King's College School of Medicine and Dentistry, London, SE5 9NU, UK

SO British Journal of Pharmacology (1998), 124(4), 685-692

CODEN: BJPCBM; ISSN: 0007-1188

PB Stockton Press

DT Journal


LA English

AB The effects of seven agonist and three antagonist adenosine receptor

ligands were compared on the guinea pig sinoatrial (SA) node (isolated right atrium) and atrioventricular (AV) node (perfused whole heart). Single agonist concentration-effect curves were obtained to 5'-N-ethylcarboxamidoadenosine, R(-)-N6-(2-phenylisopropyl)adenosine (R-PIA), N6-cyclohexyladenosine, 2-chloroadenosine, S(+)-N6-(2-phenylisopropyl)adenosine (L-PIA), 2-phenylaminoadenosine (CV 1808) and N6-aminoadenosine. Adenosine and/or NECA curves were obtained in the absence and presence of the antagonists 8-cyclopentyl-1,3-dipropylxanthine, CGS 15943 and N-0861. A formal comparison of the agonist and antagonist potency data was made by fitting the data to a straight line using a least squares procedure based on principal components anal. to account for the variance on both axes. The antagonist affinity ests. made on the two assays did not deviate significantly from the line of identity. The agonist p[A]50 data obtained on the two assays did not deviate from the line of identity, indicating that there were no significant differences in potencies between the two assays. The p[A]50 ratio of R-PIA and S-PIA was 1.24 in the SA node and 1.36 in the AV node, indicating no difference in the stereoselectivity of the PIA isomers between the two tissues. The agonist potency and antagonist affinity data obtained are consistent with previous findings showing that the AV and SA node data are pharmacol. indistinguishable and belong to the adenosine A1-receptor class. No evidence for the reported A3-receptor was found.

IT 53296-10-9, 2-Phenylaminoadenosine  
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
 (adenosine receptor pharmacol. classification in sinoatrial and atrioventricular nodes of guinea pigs)  
 RN 53296-10-9 CAPLUS  
 CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 13 THERE ARE 13 CAPLUS RECORDS THAT CITE THIS RECORD (13 CITINGS)  
 RE.CNT 34 THERE ARE 34 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 66 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
 AN 1998:329095 CAPLUS  
 DN 129:75990  
 OREF 129:15525a,15528a  
 TI A functional screening of adenosine analogs at the adenosine A2B receptor:  
 a search for potent agonists  
 AU De Zwart, Maarten; Link, Regina; Von Frijtag Drabbe Kunzel, Jacobien K.;  
 Cristalli, Gloria; Jacobson, Kenneth A.; Townsend-Nicholson, Andrea;  
 Ijzerman, Ad P.  
 CS Division of Medicinal Chemistry, Leiden/Amsterdam Center for Drug  
 Research, Leiden University, Leiden, 2300 RA, Neth.  
 SO Nucleosides & Nucleotides (1998), 17(6), 969-985  
 CODEN: NUNUD5; ISSN: 0732-8311  
 PB Marcel Dekker, Inc.  
 DT Journal  
 LA English  
 AB Various adenosine analogs were tested at the adenosine A2B receptor.  
 Agonist potencies were determined by measuring the cAMP production in Chinese  
 Hamster Ovary cells expressing human A2B receptors. 5'-N-Substituted  
 carboxamidoadenosines were most potent. 5'-N-Ethylcarboxamidoadenosine  
 (NECA) was most active with an EC50 value of 3.1  $\mu$ M. Other ribose  
 modified derivs. displayed low to negligible activity. Potency was  
 reduced by substitution on the exocyclic amino function (N6) of the purine  
 ring system. The most active N6-substituted derivative N6-methyl-NECA was 5

10/598,520

fold less potent than NECA. C8- and most C2-substituted analogs were virtually inactive. 1-Deaza-analogs had a reduced potency, 3- and 7-deazaanalogues were not active.

IT 53296-10-9


RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(functional screening of adenosine analogs at adenosine A2B receptor: search for potent agonists)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 45 THERE ARE 45 CAPLUS RECORDS THAT CITE THIS RECORD (46 CITINGS)

RE.CNT 28 THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS RECORD

ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 67 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1998:193066 CAPLUS

DN 128:257648

OREF 128:51007a,51010a

TI Synthetic nucleosides and nucleotides. 40. Selective inhibition of eukaryotic DNA polymerase  $\alpha$  by

9-( $\beta$ -D-arabinofuranosyl)-2-(p-n-butylanilino)adenine 5'-triphosphate (BuAaraATP) and its 2'-up azido analog: synthesis and enzymic evaluations

AU Tomikawa, Aki; Sato-Kiyotaki, Kunie; Ohtsuki, Chizuru; Hirai, Toshiaki; Yamaguchi, Toyofumi; Kawaguchi, Takeo; Yoshida, Shonen; Saneyoshi, Mineo

CS Dep. Biol. Sci., Teikyo Univ. Sci. Technol., Yamanashi, 409-01, Japan

SO Nucleosides & Nucleotides (1998), 17(1-3), 487-501

CODEN: NUNUD5; ISSN: 0732-8311

PB Marcel Dekker, Inc.

DT Journal

LA English

OS CASREACT 128:257648

AB Starting from 2',3',5'-tri-O-acetyl-2-iodoadenosine,

9-( $\beta$ -D-arabinofuranosyl)-2-(p-n-butylanilino)adenine and its 2'(S)-azido counterparts were synthesized in seven steps. These exhibited only moderate growth-inhibitory effects against mouse leukemic P388 cells (IC<sub>50</sub> = 13-24  $\mu$ M), although 5'-triphosphate derivs. showed strong and selective inhibitory action on calf thymus DNA polymerase  $\alpha$ , but not on  $\beta$ - and  $\epsilon$ -polymerases from eukaryotes.

IT 169687-98-3P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); RCT (Reactant); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent)


(preparation and enzymic evaluations of (arabinofuranosyl)(p-butylanilino)adenine triphosphate and its azido analog)

RN 169687-98-3 CAPLUS

CN 9H-Purine-2,6-diamine, 9- $\beta$ -D-arabinofuranosyl-N2-(4-butylphenyl)- (CA INDEX NAME)


Absolute stereochemistry.

10/598,520



IT 169687-92-7P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
(Reactant or reagent)  
(preparation and enzymic evaluations of  
(arabinofuranosyl)(p-butylanilino)adenine triphosphate and its azido  
analog)  
RN 169687-92-7 CAPLUS  
CN Adenosine, 2-[(4-butylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 23 THERE ARE 23 CAPLUS RECORDS THAT CITE THIS RECORD (23 CITINGS)  
RE.CNT 27 THERE ARE 27 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 68 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1997:576691 CAPLUS  
DN 127:243272  
OREF 127:47336a  
TI Method and composition using purines and other compounds for inhibiting  
cellular irreversible changes due to stress  
IN Miller, Guy; Lou, Lillian; Nakamura, John  
PA Galileo Laboratories, Inc., USA  
SO PCT Int. Appl., 31 pp.  
CODEN: PIXXD2  
DT Patent  
LA English  
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9730713	A1	19970828	WO 1997-US2945	19970220
	W:	AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN			
	RW:	KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG			
	US 5801159	A	19980901	US 1996-607022	19960223
	CA 2247461	A1	19970828	CA 1997-2247461	19970220
	AU 9719749	A	19970910	AU 1997-19749	19970220
	EP 935466	A1	19990818	EP 1997-907855	19970220
	R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI			
	JP 2000506834	T	20000606	JP 1997-530408	19970220
	NO 9803823	A	19981001	NO 1998-3823	19980820
PRAI	US 1996-607022	A	19960223		
	WO 1997-US2945	W	19970220		


ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

McIntosh

10/598,520

OS MARPAT 127:243272  
AB Formulations of naturally occurring physiol. acceptable compds. and their derivs. are provided for treatment of cellular stress, particularly hypoxia. By administering the formulations, comprising for the most part purines, sugars, amino acids and physiol. acceptable derivs. thereof, by themselves or in combination with each other and with other compds., particularly electron acceptor compds., the time to irreversible cellular changes, particularly mortality, can be greatly extended.  
IT 53296-10-9, 2-Phenylaminoadenosine  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(purines and other compds. for inhibition of cellular irreversible changes due to stress)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.




OSC.G 11 THERE ARE 11 CAPLUS RECORDS THAT CITE THIS RECORD (11 CITINGS)  
RE.CNT 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 69 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1997:534191 CAPLUS  
DN 127:200473  
OREF 127:38779a,38782a  
TI Adenosine A3 receptors promote degranulation of rat mast cells both in vitro and in vivo  
AU Reeves, J. J.; Jones, C. A.; Sheehan, M. J.; Vardey, C. J.; Whelan, C. J.  
CS Medicines Research Center, Glaxo Wellcome Research Development Ltd., Stevenage, SG1 2NY, UK  
SO Inflammation Research (1997), 46(5), 180-184  
CODEN: INREFB; ISSN: 1023-3830  
PB Birkhaeuser  
DT Journal  
LA English  
AB The effects were investigated of adenosine receptor agonists and antagonists on 5-HT release from rat isolated pleural mast cells and on plasma protein extravasation in the skin of conscious rats. In isolated mast cells, each adenosine agonist enhanced DNP-induced 5-HT release, N6-(3-iodobenzyl)-5-(N-methyl-carboxamidoadenosine) (IB-MECA), being the most potent agonist. The adenosine A1/A2 antagonist, 8-phenyltheophylline (8-PT), had no effect on the response to IB-MECA.  
3-(4-Amino-iodobenzyl)-8-[4-[[[carboxy]methyl]oxy]phenyl]-1-propylxanthine (I-ABOPX) inhibited (pA2 6.2) the IB-MECA responses. In the skin of conscious rats, intradermal IB-MECA produced a marked blood plasma protein extravasation (PPE) which was mimicked by N6-2-(4-aminophenyl)-ethyladenosine (APNEA). The PPE produced by IB-MECA was not affected by either 8-PT or CGS15943A, but was virtually abolished by cyproheptadine and in rats pre-treated with Compound 48/80. Thus, stimulation of adenosine A3 receptors both enhances degranulation in vitro and directly produces degranulation of rat mast cells in vivo.  
IT 53296-10-9, CV-1808  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)  
(effect on antigen-induced release of 5-HT from mast cells)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.

McIntosh



OSC.G 49 THERE ARE 49 CAPLUS RECORDS THAT CITE THIS RECORD (49 CITINGS)

L6 ANSWER 70 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1997:478039 CAPLUS

DN 127:171088

OREF 127:32965a,32968a

TI QSAR analysis of 2-alkyloxy and 2-aralkyloxy adenosine A1- and A2-agonists

AU Matova, M.; Nacheva, R.; Boicheva, S.

CS Department of Chemistry and Biochemistry, Faculty of Medicine, Medical University, Sofia, 1431, Bulg.

SO European Journal of Medicinal Chemistry (1997), 32(6), 505-513

CODEN: EJMCA5; ISSN: 0223-5234

PB Elsevier

DT Journal

LA English

AB A quant. structure-activity relationship (QSAR) anal. of a series 2-alkyloxy-, 2-aryloxy- and 2-aralkyloxy-adenosines has been performed. Various theor. 3-D electronic and topol. descriptors encoding their mol. structure were estimated and the structure-activity correlations were evaluated. A cluster anal. of the affinity consts. of the compds. was carried out, and according to the obtained results the QSAR anal. was developed at two levels. The results of this investigation allowed a distinction to be made between A1- and A2-receptor selectivity of the compds. due to structural reasons. It was shown that small and less lipophilic substituents may enhance the A1-receptor selectivity of the compds. Hydrophobic and bulky cycloalkyl substituents greatly enhance A2-receptor selectivity. The more lipophilic and rigid aromatic substituents increase the affinity, but decrease selectivity at both receptors. Adenosine agonist activity is also determined by the electron-donating properties of the purine ring and of certain atoms in this aromatic system: the N6 atom in A1-selective ligands and the N1, N7, C2, C5, C6, C8 atoms in A2-selective ligands appear to constitute part of the pharmacophore of the mols.

IT 50257-82-4 50257-84-6 50257-85-7

50257-89-1 50257-95-9 131865-78-6

131865-81-1 131865-82-2 131933-15-8

131933-17-0 131933-20-5 131933-26-1

131933-27-2 131973-26-7 137817-83-5

137817-84-6 145747-87-1 194154-31-9

194154-32-0


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)

(QSAR anal. of 2-alkyloxy and 2-aralkyloxy adenosine A1- and A2-agonists)

RN 50257-82-4 CAPLUS

CN Adenosine, 2-phenoxy- (CA INDEX NAME)

Absolute stereochemistry.






10/598,520


RN 50257-84-6 CAPLUS  
CN Adenosine, 2-butoxy- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 50257-85-7 CAPLUS  
CN Adenosine, 2-(pentyloxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 50257-89-1 CAPLUS  
CN Adenosine, 2-(2-hydroxyethoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 50257-95-9 CAPLUS  
CN Adenosine, 2-(hexyloxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 131865-78-6 CAPLUS  
CN Adenosine, 2-(phenylmethoxy)- (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 131865-81-1 CAPLUS  
CN Adenosine, 2-(4-phenylbutoxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 131865-82-2 CAPLUS  
CN Adenosine, 2-[(5-phenylpentyl)oxy]- (CA INDEX NAME)

Absolute stereochemistry.



RN 131933-15-8 CAPLUS  
CN Adenosine, 2-(3-methylbutoxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 131933-17-0 CAPLUS  
CN Adenosine, 2-(cyclohexylmethoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 131933-20-5 CAPLUS  
CN Adenosine, 2-(4-cyclohexylbutoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 131933-26-1 CAPLUS  
CN Adenosine, 2-(2-ethylbutoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 131933-27-2 CAPLUS  
CN Adenosine, 2-[(3-ethylpentyl)oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 131973-26-7 CAPLUS  
CN Adenosine, 2-[(4-methylpentyl)oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 137817-83-5 CAPLUS  
CN Adenosine, 2-[(2R)-2-phenylbutoxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 137817-84-6 CAPLUS  
CN Adenosine, 2-[(2S)-2-phenylbutoxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 145747-87-1 CAPLUS  
CN Adenosine, 2-[(3S)-3-phenylbutoxy]- (CA INDEX NAME)

Absolute stereochemistry.




RN 194154-31-9 CAPLUS  
CN Adenosine, 2-[[3S)-3,7-dimethyl-6-octenyl]oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520



RN 194154-32-0 CAPLUS  
CN Adenosine, 2-[[[(3R)-3,7-dimethyl-6-octenyl]oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 7 THERE ARE 7 CAPLUS RECORDS THAT CITE THIS RECORD (7 CITINGS)  
RE.CNT 34 THERE ARE 34 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT


L6 ANSWER 71 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1997:366482 CAPLUS  
DN 127:76385  
OREF 127:14461a,14464a  
TI Characterization of human A2A adenosine receptors with the antagonist radioligand [3H]-SCH 58261  
AU Dionisotti, Silvio; Ongini, Ennio; Zocchi, Cristina; Kull, Bjorn; Arslan, Giulia; Fredholm, Bertil B.  
CS Schering-Plough Research Institute, Milan, I-20132, Italy  
SO British Journal of Pharmacology (1997), 121(3), 353-360  
CODEN: BJPCBM; ISSN: 0007-1188  
PB Stockton  
DT Journal  
LA English  
AB The authors have characterized the binding of the new potent and selective antagonist radioligand [3H]-5-amino-7-(2-phenylethyl)-2-(2-furyl)-pyrazolo[4,3-e]-1,2,4-triazolo[1,5-c]pyrimidine, [3H]-SCH 58261, to human cloned A2A adenosine receptors. In Chinese hamster ovary (CHO) cells transfected with the human cloned A2A receptor, [3H]-SCH 58261 specific binding (about 70%) was rapid, saturable, reversible and proportional to protein concentration. The kinetic KD value was 0.75 nM. Saturation expts. showed that [3H]-SCH 58261 labeled a single class of recognition sites with high affinity (KD = 2.3 nM) and limited capacity (apparent Bmax = 526 fmol mg-1 protein). Competition expts. revealed that binding of 0.5 nM [3H]-SCH 58261 was displaced by adenosine receptor agonists with the following order of potency: 2-hexynyl-5'-N-ethylcarboxamidoadenosine (2HE-NECA) > 5'-N-ethylcarboxamidoadenosine (NECA) = 2-phenylaminoadenosine (CV 1808) > 2-[4-(2-carboxyethyl)-phenethylamino]-5'-N-ethylcarboxamidoadenosine (CGS 21680) > R-N6-phenylisopropyladenosine (R-PIA) ≥ N6-cyclohexyladenosine (CHA) > S-N6-phenylisopropyladenosine (S-PIA). Adenosine receptor antagonists inhibited [3H]-SCH 58261 binding with the following order: 5-amino-9-chloro-2-(2-furyl)-[1,2,4]-triazolo[1,5-c]quinazoline (CGS 15943) > SCH 58261 > xanthine amine congener (XAC) > (E,18%-Z,82%)7-methyl-8-(3,4-dimethoxystyryl)-1,3-dipropylxanthine (KF 17837S) > 8-cyclopentyl-1,3-dipropylxanthine (DPCPX) > theophylline. Affinity values and rank order of potency of both receptor agonists and antagonists were similar to those previously obtained in human platelet and rat striatal membranes, except for CV 1808 which was more potent than CGS 21680. SCH 58261 was a competitive antagonist at inhibiting NECA-induced cAMP accumulation in CHO cells transfected with human A2A

McIntosh

receptors. Good agreement was found between binding and functional data. Thus, the new antagonist radioligand is preferable to the receptor agonist radioligand [3H]-CGS 21680 hitherto used to examine the pharmacol. of human cloned A2A adenosine receptors.

IT 53296-10-9, 2-Phenylaminoadenosine  
 RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
 (characterization of human A2A adenosine receptors with antagonist radioligand [3H]-SCH 58261)  
 RN 53296-10-9 CAPLUS  
 CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 36 THERE ARE 36 CAPLUS RECORDS THAT CITE THIS RECORD (36 CITINGS)  
 RE.CNT 36 THERE ARE 36 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 72 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
 AN 1997:293390 CAPLUS  
 DN 127:1181

OREF 127:287a,290a

TI Binding of [125I]AB-MECA to the human cloned adenosine A3 receptor using the Semliki Forest virus expression system

AU Patel, M.; Harris, C.; Lundstrom, K.

CS Department of Receptor Pharmacology, Glaxo-Wellcome Medicines Research Centre, Herts, UK

SO Drug Development Research (1997), 40(1), 35-40

CODEN: DDREDK; ISSN: 0272-4391

PB Wiley-Liss

DT Journal

LA English

AB The cDNA for the human adenosine A3 receptor was introduced into the pSFV1 vector, and the in vitro transcribed RNA was electroporated into baby hamster kidney (BHK) cells with pSFV-Helper RNA. This protocol resulted in packaging of a high titer Semliki Forest Virus (SFV)-A3 virus stock. Infection of confluent Chinese hamster ovary (CHO) cells with the SFV-A3 virus stock resulted in an expression of human adenosine A3 receptors that was twofold more than that obtained with usual transfection methods (as determined by radioligand binding studies). The binding of [125I]N6-(4-amino-3-iodobenzyl)adenosine-5'-N-methyl-uronamide ([125I]AB-MECA) was specific and saturable ( $pK_d = 8.8$ ;  $B_{max} = 0.5$  pmol mg<sup>-1</sup> protein). Adenosine receptor ligands were evaluated for their binding affinities at the human cloned adenosine A3 receptor. The rank order of affinities of the ligands were: CGS 15943 > IB-MECA > APNEA > ligands with selectivity for adenosine A1, A2A, and A2B receptors. However, the A1 selective ligand, GR79236, had little or no affinity for the human adenosine A3 receptor. In conclusion, the SFV expression system can be used to express the human cloned adenosine A3 receptor at high levels in CHO cells. This study has examined the binding affinities at the human cloned adenosine A3 receptor, of an extensive range of ligands for the adenosine family of receptors. Furthermore, CGS 15943 has been identified as a ligand with high affinity at the human cloned adenosine A3 receptor.

IT 53296-10-9, CV 1808

RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)


(binding of [125I]AB-MECA to human cloned adenosine A3 receptor using Semliki Forest virus expression system)

RN 53296-10-9 CAPLUS

10/598,520

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 9 THERE ARE 9 CAPLUS RECORDS THAT CITE THIS RECORD (10 CITINGS)  
RE.CNT 27 THERE ARE 27 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 73 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1997:107356 CAPLUS

DN 126:113152

OREF 126:21729a,21732a

TI A method for measuring the adenosine A2a receptor binding activity of compounds of pharmacological interest by the use of the tritiated ligand [3H]-SCH 58261

IN Zocchi, Cristina; Baraldi, Pier Giovanni; Cacciari, Barbara; Dionisotti, Silvio; Ongini, Ennio

PA Schering-Plough S.P.A., Italy; Zocchi, Cristina; Baraldi, Pier Giovanni; Cacciari, Barbara; Dionisotti, Silvio; Ongini, Ennio

SO PCT Int. Appl., 13 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9638728	A1	19961205	WO 1996-EP2348	19960601
	W:	AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG			
	RW:	KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA			
	AU 9661238	A	19961218	AU 1996-61238	19960601
PRAI	IT 1995-MI1155	A	19950602		
	WO 1996-EP2348	W	19960601		

AB The invention relates to a method for evaluating the adenosine A2a receptor binding affinity of compds. of pharmacol. interest. Moreover, the invention relates to reagents and a kit particularly suitable for the above mentioned purpose. Tritiation of SCH 58261 is described, as are results of binding competition expts.

IT 53296-10-9, 2-Phenylaminoadenosine


RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(tritiated SCH 58261 preparation for adenosine A2a receptor binding activity determination for compds. of pharmacol. interest)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.




McIntosh

10/598,520

OSC.G 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD (3 CITINGS)

L6 ANSWER 74 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1996:760265 CAPLUS  
DN 126:54832  
OREF 126:10679a,10682a  
TI Comparison of nucleoside transport binding sites in rabbit iris-ciliary body and cultured rabbit nonpigmented ciliary epithelial cells  
AU Williams, Evan F.; Chu, Teh-Ching; Socci, Robin R.; Brown, Lester G.; Walker, Cassandra E.; Manor, Errol L.  
CS Dept. of Pharmacology and Toxicology, Morehouse School of Medicine, Atlanta, GA, USA  
SO Journal of Ocular Pharmacology and Therapeutics (1996), 12(4), 461-469  
CODEN: JOPTFU; ISSN: 1080-7683  
PB Liebert  
DT Journal  
LA English  
AB The iris-ciliary body (ICB) is a site of action for topically applied antiglaucoma drugs. Moreover, adenosine has been implicated as a modulator of aqueous humor dynamics. The present study compared the binding of a nucleoside transporter probe, [3H]nitrobenzylthioinosine ([3H]NBMPR), by homogenates prepared from rabbit ICB and a cultured rabbit nonpigmented ciliary epithelial cell line (NPE) to determine whether NPE can be used as an exptl. model to study the nucleoside transporter. Linear transformation of the saturation binding data revealed that [3H]NBMPR bound to a homogeneous population of binding sites with similar binding affinities in NPE and ICB (Kd 0.3 and 0.6 nM, resp.). However, the maximal binding capacity in NPC (Bmax 288 fmol/mg protein) was significantly higher than that in ICB (Bmax 154 fmol/mg protein). Selected inhibitors of the nucleoside transport system and structural analogs of adenosine inhibited the binding in both homogenate preps. with a similar rank order of potency: S-(p-nitrobenzyl)-6-thioinosine > dipyridamole > 2-phenylaminoadenosine > N6-cyclohexyladenosine > R- > S-(+)-N6-(2-phenylisopropyl)adenosine > 2-chloroadenosine > 5'-(N-ethylcarboxamido)adenosine. The results suggest that NPE is a model which could be used for characterizing the nucleoside transporter in ICB and for the screening of nucleoside transport inhibitors as potential antiglaucoma drugs.  
IT 53296-10-9, 2-Phenylaminoadenosine  
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
(nucleoside transport binding sites in iris-ciliary body and nonpigmented ciliary epithelial cells characterized by use of)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD (3 CITINGS)

L6 ANSWER 75 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1996:306232 CAPLUS  
DN 125:1181  
OREF 125:267a,270a  
TI Interaction of full and partial agonists of the A1 adenosine receptor with receptor/G protein complexes in rat brain membranes  
AU Lorenzen, Anna; Guerra, Laura; Vogt, Heidrun; Schwabe, Ulrich  
CS Ist. Farmacologia, Univ. Ferrara, Ferrara, I-44100, Italy  
SO Molecular Pharmacology (1996), 49(5), 915-926  
CODEN: MOPMA3; ISSN: 0026-895X  
PB Williams & Wilkins  
DT Journal

McIntosh




LA English  
 AB Full and partial agonists of the A1 adenosine receptor were characterized with respect to their influence on G protein activation and their thermodyn. parameters of receptor binding in rat brain membranes. G protein activation was determined through measurement of [35S]guanosine-5'-( $\gamma$ -thio)-triphosphate ([35S]GTP[S]) binding, and receptor binding was studied under identical conditions through the displacement of [3H]-1,3-dipropyl-8-cyclopentylxanthine ([3H]DPCPX) in equilibrium binding studies. The intrinsic activity in stimulating [35S]GTP[S] binding did not correlate with the affinity of the ligands. 5'-Deoxy-5'-methylthioadenosine, 2-phenylaminoadenosine, and 2-chloro-2'-deoxyadenosine were identified as partial A1 agonists in the G protein activation assay. Depending on the temperature, these ligands showed agonistic and antagonistic properties to varying extents. EC50 values for G protein stimulation and KH and KL values of the partial agonists decreased when the incubations were performed at lower temps., indicating a mainly enthalpy-driven process of interaction with the receptor. Thermodyn. parameters of receptor binding of the partial agonists resembled the characteristics of the antagonist DPCPX more closely than those of the agonist 2-chloro-N6-cyclopentyladenosine. In addition, partial agonists detected fewer A1 adenosine receptors in the high affinity state binding of [35S]GTP[S] is probably the consequence of an impaired ability of the partial agonists to release GDP from the G protein, as was shown by an impaired release of prebound [35S]GDP[S] from the membranes.

IT 53296-10-9, CV 1808  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)  
 (interaction of full and partial agonists of A1 adenosine receptor with receptor/G protein complexes in rat brain membranes)

RN 53296-10-9 CAPLUS  
 CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 63 THERE ARE 63 CAPLUS RECORDS THAT CITE THIS RECORD (63 CITINGS)

L6 ANSWER 76 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
 AN 1996:269910 CAPLUS  
 DN 124:333784  
 OREF 124:61713a,61716a  
 TI Pharmacological and biochemical characterization of purified A2a adenosine receptors in human platelet membranes by [3H]-CGS 21680 binding  
 AU Varani, Katia; Gessi, Stefania; Dalpiaz, Alessandro; Borea, Pier Andrea  
 CS Institute of Pharmacology, University of Ferrara, Ferrara, 44100, Italy  
 SO British Journal of Pharmacology (1996), 117(8), 1693-701  
 CODEN: BJPCBM; ISSN: 0007-1188  
 PB Stockton  
 DT Journal  
 LA English  
 AB The binding properties of human platelet A2a adenosine receptors, assayed with the A2a-selective agonist, [3H]-2-[p-(2-carboxyethyl)-phenethylamino]-5'-N-ethylcarboxamidoadenosine ([3H]-CGS 21680), are masked by a non-receptorial component, the adenotin site. To sep. A2a receptors from adenotin sites, human platelet membranes were solubilized with 1% 3-[(3-cholamidopropyl)dimethylammonio]-1-propane sulfonate (CHAPS). The soluble platelet extract was precipitated with polyethylene glycol (PEG) and the fraction enriched in adenosine receptors was isolated from the precipitate by differential centrifugation. The present paper describes the binding characteristics of the selective A2a agonist, [3H]-CGS 21680, to this purified platelet membrane preparation. In addition, receptor affinity and potency of several adenosine agonists and antagonists were determined in binding and

10/598,520

adenylyl cyclase studies. Saturation expts. revealed a single class of binding site with  $K_d$  and  $B_{max}$  values of 285 nM and 2.07 pmol/mg of protein resp. Adenosine receptor ligands competed for the binding of 50 nM [3H]-CGS 21680 to purified protein, showing a rank order of potency consistent with that typically found for interactions with the A2a adenosine receptors. In the adenylyl cyclase assay the compds. examined exhibited a rank order of potency very close to that observed in binding expts. Thermodyn. data indicated that [3H]-CGS 21680 binding to the purified receptor is totally entropy-driven in agreement with results obtained in rat striatal A2a adenosine receptors. It is concluded that in the purified platelet membranes there is a CGS 21680 binding site showing the characteristic properties of the A2a receptor. This makes it possible to use this compound for reliable radioligand binding studies on the A2a adenosine receptor of human platelets.

IT 53296-10-9, CV 1808


RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(adenosine A2a receptors of human platelet membranes solubilization and characterization)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 44 THERE ARE 44 CAPLUS RECORDS THAT CITE THIS RECORD (44 CITINGS)

L6 ANSWER 77 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1996:239903 CAPLUS

DN 124:279179

OREF 124:51395a,51398a

TI Ribosylpurine derivatives for treatment of cerebrovascular disorders by vascular permeability enhancer inhibition

IN Nagaoka, Akinobu; Imamoto, Tetsuji; Asano, Tsuneo; Sugiura, Yoshihiro; Goto, Gichi

PA Takeda Chemical Industries, Ltd., Japan

SO Can. Pat. Appl., 52 pp.

CODEN: CPXXEB

DT Patent

LA English


FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	CA 2150780	A1	19951203	CA 1995-2150780	19950601
	EP 704215	A2	19960403	EP 1995-108322	19950530
	EP 704215	A3	19980401		
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, NL, PT, SE				
	JP 08048631	A	19960220	JP 1995-134618	19950601
	US 5604210	A	19970218	US 1995-456723	19950601
PRAI	JP 1994-120947	A	19940602		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OS MARPAT 124:279179

GI



AB A composition is disclosed for preventing or treating brain edema, intracranial hemorrhage, and cerebral infarction by inhibiting a vascular permeability enhancer. The composition comprises I [A = halo, XR3 (X = O, S, NH, NHNH; R3 = H, acyl, (substituted) hydrocarbyl, (substituted) aromatic heterocyclyl), Y:R4 (Y = N:, NHN:; R4 = (substituted) divalent hydrocarbyl); R1 = H, halo, (substituted) hydrocarbyl, (substituted) heterocyclyl, ZR5 (Z = O, S, NH; R5 = H, (substituted) hydrocarbyl, (substituted) aromatic heterocyclyl); R2 = H, halo, (substituted) hydrocarbyl, (substituted) heterocyclyl; B = WR6 (W = CH2, C:O, C:S; R6 = OH, (substituted) alkoxy, acyloxy, alkylsulfonyl, alkylsulfonyl, O-phosphono, amino, or B together with E form cyclic phosphoric ester); D, E = H, (substituted) amino, azido, halo, (protected) OH] or a pharmaceutically acceptable salt thereof. Inhibitory activity of 42 compds. to a vascular permeability enhancer was determined 2',3'-O-(1-ethoxyethylidene)adenosine-5'-(N-ethylcarboxamide) was shown to have efficacy in preventing stroke in an animal model. Tablet and injection formulations of 6-[2-(9H-purin-6-yl)hydrazino]nebularine are included.

IT 53296-10-9 70590-23-7 70590-29-3  
71231-81-7 74615-32-0 74615-39-7  
75106-29-5 75106-32-0 75106-33-1  
102711-94-4 102711-99-9 102712-00-5  
169687-92-7 175552-71-3 175552-74-6


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(ribosylpurine derivs. for treatment of cerebrovascular disorders by vascular permeability enhancer inhibition)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.




RN 70590-23-7 CAPLUS

CN Adenosine, 2-[(3-methylphenyl)amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

10/598,520




RN 70590-29-3 CAPLUS  
CN Adenosine, 2-[(3-chloro-4-methoxyphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 71231-81-7 CAPLUS  
CN Adenosine, 2-[(4-ethylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 74615-32-0 CAPLUS  
CN Adenosine, 2-[[4-(aminocarbonyl)phenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 74615-39-7 CAPLUS  
CN Adenosine, 2-[[4-[(ethylamino)carbonyl]phenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 75106-29-5 CAPLUS  
CN Adenosine, 2-[(4-acetylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 75106-32-0 CAPLUS  
CN Adenosine, 2-[(4-acetyl-3-methoxyphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 75106-33-1 CAPLUS  
CN Adenosine, 2-[(4-acetyl-3-chlorophenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 102711-94-4 CAPLUS  
CN Adenosine, 2-[[4-(4-morpholinylcarbonyl)phenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 102711-99-9 CAPLUS  
CN Adenosine, 2-[(3-aminophenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 102712-00-5 CAPLUS  
CN Adenosine, 2-[[3-(1-oxopropyl)phenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 169687-92-7 CAPLUS  
CN Adenosine, 2-[(4-butylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 175552-71-3 CAPLUS  
CN Adenosine, 2-[(4-propylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

McIntosh


10/598,520



RN 175552-74-6 CAPLUS

CN Adenosine, 2-[[4-(1-methylpropyl)phenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)

L6 ANSWER 78 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1996:187715 CAPLUS

DN 124:279743

OREF 124:51535a,51538a

TI Functional characterization of adenosine A2 receptors in Jurkat cells and PC12 cells using adenosine receptor agonists

AU van der Ploeg, Ingeborg; Ahlberg, Susanne; Parkinson, Fiona E.; Olsson, Ray A.; Fredholm, Bertil B.

CS Department Physiology Pharmacology, Karolinska Institute, Stockholm, S-171 77, Swed.

SO Naunyn-Schmiedeberg's Archives of Pharmacology (1996), 353(3), 250-60  
CODEN: NSAPCC; ISSN: 0028-1298

PB Springer

DT Journal

LA English

AB The effect of several adenosine analogs on cAMP accumulation was examined in the rat pheochromocytoma cell PC12 and in the human T-cell leukemia cell Jurkat, selected as prototypes of cells predominantly expressing adenosine A2A or A2B receptors. Using the reverse transcription-polymerase chain reaction it was, however, demonstrated that the Jurkat cell and the PC12 cell express both A2A and A2B receptor mRNA, albeit in different relative proportions. In PC12 cells, the concentration required for half-maximal response (EC50) for the full agonist NECA was 30 times lower than in Jurkat cells. There was no significant difference in the pA2 for the antagonist CGS 15943 between the two cell types. In the presence of forskolin (1  $\mu$ M in PC12 cells; 10  $\mu$ M in Jurkat cells) the EC50 value for NECA was reduced two-to sixfold. Forskolin also increased the maximal cAMP accumulation twofold in PC12 cells and sevenfold in Jurkat cells. A series of 2-substituted adenosine analogs CV 1808, CV 1674, CGS 21680, and four 2-substituted isoguanosines, SHA 40, SHA 91, SHA 118, and SHA 125, all raised cAMP accumulation in PC12 cells, but had minimal or no effect in Jurkat cells. In the PC12 cells the addition of forskolin (1  $\mu$ M) reduced the EC50 by a factor of 2 (CV 1808) to 12 (SHA 125). In Jurkat cells all the analogs gave a significant, but submaximal, cAMP response in the presence of forskolin (10  $\mu$ M), but they were essentially inactive in its absence. The results show that a series of 2-substituted adenosine analogs can be used to discriminate between A2A and A2B receptors. The two receptor subtypes appear to coexist, even in clonal cells selected for typical pharmacol. A2 receptor pharmacol. can therefore be complex.

IT 50257-95-9, 2-Hexyloxyadenosine 53296-10-9, CV 1808

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BUU (Biological use, unclassified); BIOL (Biological

10/598,520


study); USES (Uses)

(adenosine A2 receptor subtype functional characterization in Jurkat cells and PC12 cells using adenosine receptor agonists)

RN 50257-95-9 CAPLUS

CN Adenosine, 2-(hexyloxy)- (CA INDEX NAME)


Absolute stereochemistry.



RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 43 THERE ARE 43 CAPLUS RECORDS THAT CITE THIS RECORD (43 CITINGS)

L6 ANSWER 79 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1996:162066 CAPLUS

DN 124:221345

OREF 124:40737a,40740a

TI Pharmacological probes for A1 and A2 adenosine receptors in vivo in feline pulmonary vascular bed

AU Neely, Constance Fisher; Matot, Idit

CS Sch. Med., Univ. Pennsylvania, Philadelphia, PA, 19104, USA

SO American Journal of Physiology (1996), 270(2, Pt. 2), H610-H619

CODEN: AJPHAP; ISSN: 0002-9513

PB American Physiological Society

DT Journal

LA English

AB Under conditions of controlled pulmonary blood flow and constant left atrial pressure, adenosine produces dose-dependent, tone-dependent responses in the pulmonary vascular (PV) bed of intact-chest, spontaneously breathing cats. The potency profile for adenosine receptor agonists to produce vasoconstriction at low baseline PV tone is 5'-(N-ethylcarboxamido)adenosine  $\geq$  CGS-21680  $\geq$  2-chloroadenosine (2-CADO)  $\geq$  [R]-N6-(2-phenylisopropyl)adenosine (R-PIA)  $\geq$  N6-cyclopentyladenosine > adenosine  $\gg$  CV-1808. After an increase in PV tone with the use of an intralobar infusion of the thromboxane mimic U-46619, the potency profile for adenosine receptor agonists to produce vasodilation at elevated PV tone is 2-CADO  $\geq$  CV-1808  $\geq$  CGS-21680 > R-PIA  $\geq$  adenosine. The selective A1 adenosine receptor antagonists xanthine amine congener (XAC) and 8-cyclopentyl-1,3-dipropylxanthine (DP-CPX) significantly antagonize the vasoconstrictor responses of adenosine and R-PIA at low baseline PV tone while having less effect on the vasodilator responses of adenosine, 2-CADO, and R-PIA at elevated PV tone. DPCX antagonizes the vasoconstrictor responses of CGS-21680 at low baseline PV tone. The nonselective A1 and A2 adenosine receptor antagonist BWA-1433U significantly antagonizes the vasoconstrictor responses of R-PIA and vasodilator responses of adenosine, 2-CADO, and R-PIA. These data support

McIntosh



that adenosine produces vasoconstriction at low baseline PV tone and vasodilation at elevated PV tone in the feline PV bed by acting on A1 and A2 adenosine receptors, resp. Compared with the adenosine receptor agonists tested in this in vivo model, R-PIA and CV-1808 are the most selective adenosine receptor agonists for A1 and A2 adenosine receptors, resp., in the feline PV bed. R-PIA, CV-1808, DPCPX, and XAC may be used in this in vivo model to define the roles of A1 and A2 adenosine receptors in acute lung injury and pathophysiol. changes in the pulmonary vasculature associated with pulmonary hypertension and edema formation in the same animal model.

IT 53296-10-9, CV-1808


RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

(pharmacol. probes for A1 and A2 adenosine receptors in vivo in feline pulmonary vascular bed)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 9 THERE ARE 9 CAPLUS RECORDS THAT CITE THIS RECORD (9 CITINGS)

L6 ANSWER 80 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1995:706870 CAPLUS

DN 123:102310

OREF 123:17911a,17914a

TI Therapeutic aspects of adenosine in relation to its anti-TNF properties.

AU Giroud, Jean-Paul; Lian Chen, Yan; Le Vraux, Valerie; Chauvelot-Moachon, Laurence

CS Departement de Pharmacologie, Hopital Cochin, Paris, 75679/14, Fr.

SO Bulletin de l'Academie Nationale de Medecine (Paris) (1995), 179(1), 79-101

CODEN: BANMAC; ISSN: 0001-4079

PB Academie Nationale de Medecine

DT Journal

LA French

AB Expts. tested the hypothesis that the antiinflammatory properties of adenosine occur via a down-regulation of tumor necrosis factor (TNF). Adenosine receptor agonists (ARA) and agents potentiating endogenous adenosine (APA) were evaluated for their effects on TNF production by endotoxin-stimulated human monocytes. Addnl., one of the most potent agonists, (R)-phenylisopropyladenosine (R-PIA), was tested in 2 exptl. models of acute-phase response: endotoxin shock and carrageenan-induced plantar edema. Several ARA and APA inhibited monocyte TNF production in a concentration-dependent manner. R-PIA and other ARA were active at micromolar concns. This property is pharmacol. relevant, since rats receiving a LD of endotoxin were protected by R-PIA, and the endotoxin-induced increase in serum TNF levels was abolished by pretreatment with R-PIA. Inhibitory effects on serum TNF production were obtained with similar concns. of dexamethasone and 100-fold higher concns. of pentoxifylline. R-PIA was also active on carrageenan-induced edema. The antiedema properties of R-PIA were associated with a marked reduction of locally produced TNF and were also observed after the administration of dexamethasone, pentoxifylline and a neutralizing anti-TNF antibody. The results indicate that adenosine is a potent inhibitor of TNF production induced by different stimuli. This property could lead to therapeutic applications in inflammatory diseases and other conditions in which TNF is known to play a pathogenic or aggravating role.

IT 53296-10-9, CV 1808

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)


10/598,520

(pharmacol. effects of adenosine and adenosine agonists in relation to inhibition of tumor necrosis factor production)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD (3 CITINGS)

L6 ANSWER 81 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1995:631030 CAPLUS

DN 123:286460

OREF 123:51354h,51355a

TI Synthesis and biological activities of sugar-modified 2-(p-n-butylanilino)-2'-deoxyadenosine analogs

AU Yamaguchi, Toyofumi; Kunie, Sato; Saneyoshi, Mineo

CS Department Biological Sciences, Nishi-Tokyo University, Yamanashi, 409-01, Japan

SO Nucleosides & Nucleotides (1995), 14(3-5), 529-32


CODEN: NUNUD5; ISSN: 0732-8311

PB Dekker

DT Journal

LA English

GI



AB Several sugar-modified 2-(p-n-butylanilino)-2'-deoxyadenosine analogs including arabino and 2'(R)-azido-2'-deoxy analogs I (R = H, OH, N3, R1 = H) and their 5'-triphosphates were synthesized. These nucleosides thus obtained exhibited moderate cytotoxicity against P-388 leukemic cells in culture (IC50 = 13-24  $\mu$ M). In contrast to above results, the 5'-triphosphates have been shown to exert strong and selective inhibitory effects on mammalian DNA polymerase  $\alpha$  (Ki = 0.02-0.04  $\mu$ M).

IT 169687-98-3P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); RCT (Reactant); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent) (synthesis and antitumor and antiviral activities of anilinodeoxyadenosines)


RN 169687-98-3 CAPLUS

CN 9H-Purine-2,6-diamine, 9- $\beta$ -D-arabinofuranosyl-N2-(4-butylphenyl)- (CA INDEX NAME)

McIntosh

10/598,520

Absolute stereochemistry.



IT 169687-92-7P


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(synthesis and antitumor and antiviral activities of anilinodeoxyadenosines)

RN 169687-92-7 CAPLUS

CN Adenosine, 2-[(4-butylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)

L6 ANSWER 82 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1995:590191 CAPLUS

DN 123:52110

OREF 123:9283a,9286a

TI Structure-activity relationship for the binding of nucleoside ligands to adenosine kinase from *Toxoplasma gondii*

AU Iltzsch, Max H.; Uber, Sheri S.; Tankersley, Kevin O.; el Kouni, Mahmoud H.

CS Dept. Biol. Sci., Univ. Cincinnati, Cincinnati, OH, 45221, USA

SO Biochemical Pharmacology (1995), 49(10), 1501-12

CODEN: BCPCA6; ISSN: 0006-2952

PB Elsevier

DT Journal

LA English


AB One hundred and twenty-eight purine nucleoside analogs were evaluated as ligands of *Toxoplasma gondii* adenosine kinase (EC 2.7.1.20) by examining their ability to inhibit this enzyme in vitro. Inhibition was quantified by determining apparent  $K_i$  (app $K_i$ ) values for those compds. that inhibited this enzyme by greater than 10% at a concentration of 1 mM. Two compds., N6-(p-methoxybenzoyl)adenosine and 7-iodo-7-deazaadenosine (iodotubercidin), were found to bind to the enzyme (app $K_i$  = 3.9 and 1.6  $\mu$ M, resp.) better than adenosine. On the basis of these data, a structure-activity relationship for the binding of ligands to *T. gondii* adenosine kinase was formulated using adenosine as a reference compound. It was concluded that the following structures features of purine nucleoside analogs are required or strongly preferred for binding: (1) "pyridine-type" endocyclic nitrogens at the 1- and 3-positions; (2) an exocyclic hydrogen at the 2-position; (3) 6-position exocyclic substituents in the lactim tautomeric form; (4) a "pyridine-type" endocyclic nitrogen at the 7-position or hydrophobic exocyclic substituents attached to an endocyclic carbon at the 7-position; (5) an endocyclic methine or "pyridine-type" nitrogen at the 8-position; (6) an endocyclic nitrogen at the 9-position; (7) a pentose or "pentose-like" (e.g. hydroxylated cyclopentene) moiety attached to the 9-position nitrogen; (8) hydroxyl groups at the 2'- and 3'-positions in a ribose

McIntosh

configuration; (9) a hydroxymethyl or Me (i.e. 5'-deoxy) group at the 5'-position; (10) a  $\beta$ -D-nucleoside configuration; and (11) an anti conformation around the N-glycosidic bond. In addition, there appears to be a "pocket" in the catalytic site of *T. gondii* adenosine kinase, adjacent to the 6-position of adenosine, that can accommodate large (preferably unsatd. or aromatic) substituents (e.g. phenyl). These findings provide the basis for the rational design of addnl. ligands of *T. gondii* adenosine kinase.

IT 53296-10-9, 2-Phenylaminoadenosine  
 RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)  
 (structure-activity relationship for the binding of nucleoside ligands to adenosine kinase from *Toxoplasma gondii*)  
 RN 53296-10-9 CAPLUS  
 CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 39 THERE ARE 39 CAPLUS RECORDS THAT CITE THIS RECORD (39 CITINGS)

L6 ANSWER 83 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1995:405318 CAPLUS

DN 122:255527

OREF 122:46321a,46324a

TI A theoretical structure-activity relationship study of  
 2-alkoxyadenosines: selective agonists at the coronary artery  
 A2-adenosine receptor

AU Ojha, T. N.; Singh, P.; Tiwari, Susheela; Sharma, R. C.

CS Department of Chemistry, SK Government College, Sikar, 332 001, India

SO Indian Journal of Biochemistry & Biophysics (1995), 32(1), 60-2

CODEN: IJBBBQ; ISSN: 0301-1208

PB Publications & Information Directorate, CSIR

DT Journal

LA English

AB A theor. explanation of the agonist actions of several adenosine derivs., elicited from its binding to the two subtypes of discrete membrane-bound adenosine receptors, A1AR and A2AR, has been provided on the basis of derived statistical correlations. The van der Waals volume (Vw) of R-group, which is a measure of bulk, also stands a measure of the hydrophobic nature of the R-substituent, as evidenced from its near linear relation with hydrophobicity index, k', for these ligands. Through the use of an indicator parameter, it could be inferred that if the substituent has more CH2 instead of secondary CH adjacent to the point of attachment, R-O, the ligand will be more efficacious with adenosine receptors.

IT 50257-84-6 50257-85-7 50257-89-1  
 50257-95-9 131933-15-8 131933-17-0  
 131933-20-5 131933-26-1 131933-27-2  
 131973-26-7

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)


(theor. structure-activity relationship study of 2-alkoxyadenosines as selective agonists at the coronary artery A2-adenosine receptor)

RN 50257-84-6 CAPLUS

CN Adenosine, 2-butoxy- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

10/598,520




RN 50257-85-7 CAPLUS  
CN Adenosine, 2-(pentyloxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 50257-89-1 CAPLUS  
CN Adenosine, 2-(2-hydroxyethoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 50257-95-9 CAPLUS  
CN Adenosine, 2-(hexyloxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 131933-15-8 CAPLUS  
CN Adenosine, 2-(3-methylbutoxy)- (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 131933-17-0 CAPLUS  
CN Adenosine, 2-(cyclohexylmethoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 131933-20-5 CAPLUS  
CN Adenosine, 2-(4-cyclohexylbutoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 131933-26-1 CAPLUS  
CN Adenosine, 2-(2-ethylbutoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 131933-27-2 CAPLUS  
CN Adenosine, 2-[(3-ethylpentyl)oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520



RN 131973-26-7 CAPLUS  
CN Adenosine, 2-[(4-methylpentyl)oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)


L6 ANSWER 84 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1995:367220 CAPLUS  
DN 122:123777  
OREF 122:22939a,22942a  
TI Comparison of A4 and A2a binding sites in striatum and COS cells transfected with adenosine A2a receptors  
AU Luthin, David R.; Linden, Joel  
CS Departments Internal Medicine Cardiovascular Division, University Virginia Health Sciences Center, Charlottesville, VA, USA  
SO Journal of Pharmacology and Experimental Therapeutics (1995), 272(2), 511-18  
CODEN: JPETAB; ISSN: 0022-3565  
PB Williams & Wilkins  
DT Journal  
LA English  
AB A putative A4 adenosine receptor is characterized by a distinct structure activity profile of compds. in competition for [3H]2-phenylaminoadenosine ([3H]CV 1808) binding sites on rat brain membranes assayed at 4°. We now confirm that A4 binding sites can be demonstrated on ice-cold membranes of rat striatum and demonstrate a similar binding site on COS cells transfected with rat A2a adenosine receptors (COS/A2a). The characteristic A4 potency order is: CV 1808 > [1R-(1 $\alpha$ ,2 $\alpha$ ,3 $\beta$ ,5 $\beta$ )]-3-(2,6-diamino-N2-(3-carbethoxyphenyl)-9H-purin-9-yl)-5'-(N-ethylcarbamoyl)-1,2-cyclopentanediol (CGS 22988) » 5'-N-ethylcarboxamidoadenosine (NECA)  $\geq$  2-[4-(2-carboxyethyl)phenylethylamino]-5'-N-ethylcarboxamidoadenosine (CGS 21680); 9-chloro-2-(2-furyl)[1,2,4]-triazolo[1,5-c]-quinazolin-5-amine (CGS 15943) only partially inhibits binding at 1  $\mu$ M. If [3H]CGS 21680 is used for ice-cold assays, or if either [3H]CV 1808 or [3H]CGS 21680 are used for assays at 21°, the potency order of competing compds. changes markedly and becomes characteristic of A2a adenosine receptor binding sites; CGS 15943  $\geq$  CGS 21680 .simeq. NECA > CGS 22988  $\geq$  CV 1808. Binding of [3H]CGS 21680, but not [3H]CV 1808, is enhanced by the pore-forming antibiotic, alamethicin. Guanosine 5'-O-(3-thiotriphosphate) decreases the binding of both radioligands to striatal membranes at 21° more than to membranes on ice. We propose that differential effects of temperature on the binding characteristics of compds. with distinct physicochem. properties to various pools of a single A2a adenosine receptor can result in A4 and A2a binding profiles.  
IT 53296-10-9, CV 1808  
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL

McIntosh

10/598,520

(Biological study); PROC (Process)  
(cold effect on adenosine A2a - A4 receptor binding activity of)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)


Absolute stereochemistry.



OSC.G 14 THERE ARE 14 CAPLUS RECORDS THAT CITE THIS RECORD (14 CITINGS)

L6 ANSWER 85 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1995:349895 CAPLUS  
DN 122:154099  
OREF 122:28333a,28336a  
TI Herbicidally active sulfamoyl nucleosides. Isolation and synthesis.  
AU Kristinsson, Kaukur; Nebel, Kurt; O'Sullivan, Anthony C.; Pachlatko, J. Paul; Yamaguchi, Yasuchika  
CS Crop Protection Div., Ciba-Geigy AG, Basel, 4002, Switz.  
SO ACS Symposium Series (1995), 584(Synthesis and Chemistry of Agrochemicals IV), 206-19  
CODEN: ACSMC8; ISSN: 0097-6156  
PB American Chemical Society  
DT Journal  
LA English  
AB The isolation of the herbicidal 2-chloro-5'-O-sulfamoyladenine (I) is reported. Its relation to other herbicidal nucleosides is described. Two new and direct synthetic routes to I were established and a number of derivative were prepared. Herbicidal activity was found in analogs structurally close to I. An in vitro toxicol. screen was applied to these compds.  
IT 79936-11-1  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(reaction in herbicidal sulfamoyl nucleoside preparation)  
RN 79936-11-1 CAPLUS  
CN Adenosine, 2-cyano- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD (3 CITINGS)

L6 ANSWER 86 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1995:340312 CAPLUS  
DN 122:123739  
OREF 122:22927a,22930a  
TI Binding of the adenosine A2a receptor ligand [3H]CGS 21680 to human platelet membranes  
AU Varani, Katia; Borea, Pier Andrea; Guerra, Laura; Dionisotti, Silvio; Zocchi, Cristina; Ongini, Ennio  
CS Inst. Pharmacology, Univ. Ferrara, Ferrara, 44100, Italy  
SO Research Communications in Molecular Pathology and Pharmacology (1995), 87(1), 109-10

McIntosh



CODEN: RCMPE6; ISSN: 1078-0297

PB PJD Publications

DT Journal

LA English

AB The binding characteristics of the selective adenosine A2a agonist [3H]-CGS 21680 in human platelet membranes. Addnl., the potency of several adenosine agonists was determined in adenylate cyclase studies. Specific binding was saturable, reversible, and dependent upon protein concentration. Results indicate that in platelets [3H]-CGS 21680 labels also the nonreceptor binding site (adenotin site) for [3H]-NECA binding described in peripheral tissue.

IT 53296-10-9, CV 1808


RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(adenosine A2a receptor ligand CGS 21680 and other adenosine agonists binding and functional activity in human platelet membranes)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

L6 ANSWER 87 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1995:303926 CAPLUS

DN 122:72515

OREF 122:13611a,13614a

TI Functional characterization of the adenosine receptor mediating inhibition of intestinal secretion

AU Hancock, Debra L.; Coupar, Ian M.

CS Sch. Pharmacol., Monash Univ., Victoria, 3052, Australia

SO British Journal of Pharmacology (1995), 114(1), 152-6

CODEN: BJPCBM; ISSN: 0007-1188

PB Stockton

DT Journal


LA English

AB Previous studies have shown that the mixed A1/A2 adenosine agonist 5'-N-ethylcarboxamidoadenosine (NECA) inhibits intestinal fluid secretion which is thought to contribute its antidiarrheal effect in the rat. The aim of this study was to characterize the adenosine receptor mediating this antisecretory effect via functional studies using a range of selective agonists and antagonists and by applying the pharmacol. criteria of relative agonist and antagonist potencies. Adenosine agonists and antagonists were administered i.v. to anesthetized rats. Intestinal secretion was then stimulated by i.a. infusion of vasoactive intestinal peptide (VIP, 0.8  $\mu\text{g min}^{-1}$ ) and the net fluid transport across the wall of the jejunum was measured by a recirculation technique. The rank order agonist potency to reduce the response to VIP was: NECA > N6-cyclopentyladenosine (CPA) > R-N6-(2-phenylisopropyladenosine) (R-PIA) > S-PIA > chloroadenosine (2-CADO) > 2-phenylaminoadenosine (CV-1808). This order best complies with the rank order of agonist potency that represents activation of the recently described A2B receptor: NECA > 2-CADO > R-PIA = CPA > S-PIA > CV-1808 > GCS-21680. The most potent agonists (NECA, CPA and R-PIA) had ED50 values in the low microgram range. The antisecretory action of NECA (submaximal dose of 40  $\mu\text{g kg}^{-1}$ ) was antagonized equally (approx. 50%) by the selective adenosine antagonists 8-cyclopentyl-1,3-dipropylxanthine (DPCPX, 0.1 mg  $\text{kg}^{-1}$ ) and 8-phenyltheophylline (8-PT, 0.1 mg  $\text{kg}^{-1}$ ). This equipotent activity indicates the presence of an A2 and not an A1 receptor. It is suggested that adenosine A2B receptor agonists could be evaluated for potential use

10/598,520

as antidiarrheal drugs.  
IT 53296-10-9, 2-Phenylaminoadenosine  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(adenosine receptor subtype mediating adenosine analog inhibition of intestinal secretion and antidiarrheal activity)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 20 THERE ARE 20 CAPLUS RECORDS THAT CITE THIS RECORD (20 CITINGS)

L6 ANSWER 88 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1995:275005 CAPLUS  
DN 122:46518  
OREF 122:8734a  
TI Adenosine receptor agonists for the promotion of wound healing  
IN Cronstein, Bruce N.; Levin, Richard I.  
PA New York University, USA  
SO PCT Int. Appl., 35 pp.  
CODEN: PIXXD2  
DT Patent  
LA English  
FAN.CNT 1


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9423723	A1	19941027	WO 1994-US2011	19940218
	W: AU, CA, JP				
	RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
	AU 9465164	A	19941108	AU 1994-65164	19940218
	US 5932558	A	19990803	US 1996-712942	19960913
	US 6020321	A	20000201	US 1999-243538	19990203
PRAI	US 1993-46297	A	19930415		
	WO 1994-US2011	W	19940218		
	US 1996-712942	A1	19960913		

AB Agonists of the adenosine A2 receptor promote the migration of endothelial cells, fibroblasts and epithelial cells. Thus, methods and pharmaceutical compns. useful for treating wounds and promoting wound healing comprise agents which cause stimulation of the adenosine A2 receptor, preferably receptor agonists and adenosine uptake blockers. Preferred agonists include 2-phenylaminoadenosine, 2-p-(2-carboxyethyl)phenylamino-5'-N-ethylcarboxamidoadenosine, 5'-N-ethylcarboxamidoadenosine, 5'-N-cyclopropyladenosine, 5'-N-methylcarboxamidoadenosine and PD-125944. Preferred uptake blockers include dipyridamole, nitrobenzothioinosine, dilazep and R75231.

IT 53296-10-9, 2-Phenylaminoadenosine  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(adenosine receptor agonists for the promotion of wound healing)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.

10/598,520



OSC.G 4 THERE ARE 4 CAPLUS RECORDS THAT CITE THIS RECORD (7 CITINGS)

L6 ANSWER 89 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1995:188936 CAPLUS

DN 122:669

OREF 122:155a,158a

TI Glibenclamide reduces the coronary vasoactivity of adenosine receptor agonists

AU Niiya, Kazunori; Uchida, Shinji; Tsuji, Takao; Olsson, Ray A.

CS Dep. Intern. Med. Biochem. Mol. Biol., Univ. South Florida, Tampa, FL, USA

SO Journal of Pharmacology and Experimental Therapeutics (1994), 271(1), 14-19

CODEN: JPETAB; ISSN: 0022-3565

PB Williams & Wilkins

DT Journal

LA English

AB Expts. in guinea pig heart Langendorff preps. assessed the effect of KATP channel blockade on the coronary vasoactivity of adenosine and 17 analogs chosen to represent a variety of purine and ribose modifications.

Although glibenclamide is a functional antagonist that acts at the level of an effector rather than at a receptor, it caused parallel rightward shifts of agonist dose-response curves. The size of the shift of EC50 differed according to the kind of analog: the ranking was, generally, N6-phenethyladenosines > 2-aryl-aminoadenosines = 2-(1-alkyn-1-yl)adenosines > N6-cycloalkyladenosines = adenosine 5'-uronamides. The coronary vasoactivity ranking of agonists in the presence of supramaximal concns. of glibenclamide was 2-(1-alkyn-1-yl)adenosines = 2-aralkoxyadenosines > 2-aralkylaminoadenosines > 2-arylaminoadenosines > N6-substituted adenosines. Glibenclamide did not affect the vasoactivity of adenosine itself, perhaps because avid uptake by endothelial cells prevented penetration of the agonist to receptors deeper in the vascular wall. The results exclude a model consisting of one kind of receptor acting exclusively through a KATP channel, argue against one kind of receptor coupled to a KATP channel as well as to an addnl. effector but is consistent with two kinds of vasodilatory adenosine receptors, one of which activates a KATP channel. The identity of the adenosine receptor coupled to the KATP channel is uncertain; the other receptor has the pharmacol. profile of an A2a-adenosine receptor.

IT 53296-10-9, CV 1808 76888-18-1 102712-00-5


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)

(glibenclamide reduces coronary vasoactivity of adenosine receptor agonists)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.




McIntosh

10/598,520


RN 76888-18-1 CAPLUS  
CN Adenosine, 2-[(3-acetylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 102712-00-5 CAPLUS  
CN Adenosine, 2-[[3-(1-oxopropyl)phenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 18 THERE ARE 18 CAPLUS RECORDS THAT CITE THIS RECORD (18 CITINGS)

L6 ANSWER 90 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1994:692389 CAPLUS  
DN 121:292389  
OREF 121:53203a,53206a  
TI Failure of CGS15943A to block the hypotensive action of agonists acting at the adenosine A3 receptor  
AU Patel, M.; Sheehan, M. J.; Strong, P.  
CS Cellular Sci., Glaxo Res. Dev. Ltd., Ware, Herts, SG12 0DP, UK  
SO British Journal of Pharmacology (1994), 113(3), 741-8  
CODEN: BJPCBM; ISSN: 0007-1188  
PB Stockton  
DT Journal  
LA English  
AB Adenosine receptor agonists were evaluated for their activity at the putative adenosine A3 receptor which mediates a 'xanthine-resistant' hypotensive response in the anesthetized rat. The compds. tested were: the A1/A3 receptor agonist, N-[2-(4-aminophenyl)ethyl]adenosine (APNEA), the non-selective adenosine receptor agonist, 5'-N-ethylcarboxamidoadenosine (NECA), the adenosine A1 receptor-selective agonists, N-[(1S,trans)-2-hydroxycyclopentyl]adenosine (GR79236) and N6-cyclopentyl adenosine (CPA), the A2a receptor-selective agonists, 2-[[2-[4-(2-carboxyethyl) phenyl] ethyl] amino]-N-ethylcarboxamidoadenosine (CGS21680) and 2-phenylaminoadenosine (CV1808), and the moderately A2b selective agonist, N-[(2-methylphenyl)methyl]adenosine (metrifudil). In conformation of literature findings, APNEA (1-1000 nmol kg<sup>-1</sup>) induced hypotension and bradycardia; the hypotension was not blocked by pretreatment with the xanthine antagonist, 8-P-sulfophenyltheophylline (8-sPT; 40 mg kg<sup>-1</sup>, i.v.), whereas the bradycardia was attenuated. The non-xanthine antagonist, 9-fluoro-2-(2-furyl)-5,6-dihydro [1,2,4]triazolo[1,5-c]-quinazolin-5-imine (CGS15943A; 3 mg kg<sup>-1</sup> i.v.), also attenuated the bradycardia without affecting the hypotension. The adenosine A1 receptor-selective agonists, GR79236 and CPA, both produced dose-dependent falls in blood pressure and heart rate which were antagonized by 8-sPT (40 mg kg<sup>-1</sup>) and CGS15943A (3 mg kg<sup>-1</sup>). The adenosine A2a receptor-selective agonists, CGS21680 and CV1808, produced only a hypotensive response which was antagonized by 8-sPT (40 mg kg<sup>-1</sup>)

McIntosh

and to a much greater extent by CGS15943A (3 mg kg<sup>-1</sup>), consistent with the response being mediated solely by A2a receptors. The modestly A2b receptor-selective agonist, metrifudil, produced a dose-dependent fall in blood pressure and at higher doses a fall in heart rate. The hypotension induced by metrifudil was not antagonized by either 8-sPT (40 mg kg<sup>-1</sup>) or CGS15943A (3 mg kg<sup>-1</sup>) even though the bradycardia was abolished, suggesting that this agonist activates the putative A3 receptor. The non-selective adenosine receptor agonist, NECA, produced a hypotension and bradycardia that was attenuated by 8-sPT (40 mg kg<sup>-1</sup>), confirming previous work. The non-xanthine antagonist, CGS15943A (3 mg kg<sup>-1</sup>), also attenuated the hypotension and bradycardia. The bradycardia was blocked to a much greater extent, suggesting that NECA may therefore induce hypotension partly by activating the putative A3 receptor. In conclusion, we have confirmed that the putative A3 receptor mediating hypotension in the anesthetized rat is not blocked by 8-sPT, and further shown that it is not blocked by CGS15943A. The A2a agonists CGS21680 and CV1808 showed no discernible activity at the A3 receptor, whereas APNEA, NECA, CPA and metrifudil appear to activate this receptor. The adenosine A1 receptor agonist, GR79236, shows considerable selectivity for the A1 receptor but may activate the A3 receptor at high doses.

IT 53296-10-9, CV1808


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(adenosine receptor agonists effects on 'xanthine-resistant' hypotensive response mediated by adenosine A3 receptor)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 8 THERE ARE 8 CAPLUS RECORDS THAT CITE THIS RECORD (8 CITINGS)

L6 ANSWER 91 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1994:645905 CAPLUS

DN 121:245905

OREF 121:44639a,44642a

TI 2-[2-[4-[2-[2-[1,3-Dihydro-1,1-bis(4-hydroxyphenyl)-3-oxo-5-isobenzofuranthioureidyl]ethylaminocarbonyl]ethyl]phenyl]ethylamino]-5'-N-ethylcarboxamidoadenosine (FITC-APEC): a fluorescent ligand for A2a-adenosine receptors

AU McCabe, R. Tyler; Skolnick, Phil; Jacobson, Kenneth A.

CS Lab. Neurosci., Pharm. Discovery Corp., Elmsford, NY, 10523, USA

SO Journal of Fluorescence (1992), 2(4), 217-23

CODEN: JOFLEN; ISSN: 1053-0509

DT Journal


LA English

AB The fluorescein conjugate FITC-APEC is a novel ligand derived from a series of functionalized congeners that act as selective A2a-adenosine receptor agonists. The binding of FITC-APEC to bovine striatal A2a-adenosine receptors, measured by fluorescence techniques, was saturable and of a high affinity, with a Bmax of 2.3 pmol/mg protein and KD of 57 nM. The KD value estimated by fluorescence was consistent with the Ki (11 nM) obtained by competition studies with [3H]CGS 21680. Addnl., the Bmax value found by FITC-APEC measurement was in agreement with Bmax values obtained by radioligand binding. FITC-APEC exhibited rapid and reversible binding to bovine striatum. The potencies of chemical diverse A2a-adenosine receptor ligands, as estimated by inhibition of FITC-APEC binding, were in good agreement with their potencies determined by radioligand binding techniques (r = 0.97). FITC-APEC binding was not altered by purine derivs. that do not recognize A2a-adenosine receptors. These

findings demonstrate that the novel fluorescent ligand FITC-APEC can be used in the quant. characterization of ligand binding to A2a-adenosine receptors.

IT 53296-10-9, 2-(Phenylamino)adenosine  
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
 (adenosine analog-fluorescein conjugate binding to striatal adenosine A2a receptors inhibition by)  
 RN 53296-10-9 CAPLUS  
 CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.




OSC.G 6 THERE ARE 6 CAPLUS RECORDS THAT CITE THIS RECORD (6 CITINGS)

L6 ANSWER 92 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
 AN 1994:524874 CAPLUS  
 DN 121:124874  
 OREF 121:22297a,22300a  
 TI Inhibition of platelet aggregation by adenosine receptor agonists  
 AU Cristalli, Gloria; Vittori, Sauro; Thompson, Robert D.; Padgett, William L.; Shi, Dan; Daly, John W.; Olsson, Ray A.  
 CS Dip. Sci. Chimiche, Univ. Camerino, Camerino, I-62032, Italy  
 SO Naunyn-Schmiedeberg's Archives of Pharmacology (1994), 349(6), 644-50  
 CODEN: NSAPCC; ISSN: 0028-1298  
 DT Journal  
 LA English  
 AB 2-(Ar)alkoxyadenosines, which are agonists selective for the A2AAR in PC 12 cell and rat striatum membranes, are also agonists at the A2AR coupled to adenylate cyclase (AC) that mediates the inhibition of platelet aggregation. A panel of twelve well-characterized adenosine analogs stimulated human platelet AC and inhibited ADP-induced platelet aggregation at sub- to low-micromolar concns. with a potency ranking CGS 21680>adenosine>R-PIA. There were significant correlations between the EC50 of stimulation of platelet and PC 12 cell AC ( $r_2 = 0.66$  and  $0.67$ , resp.) or the  $K_i$  of inhibition of  $[3H]$ NECA binding to the rat striatum membranes ( $r_2 = 0.75$ ). Likewise, platelet AC stimulation correlated well with stimulation of PC 12 cell AC and with  $[3H]$ NECA binding ( $r_2 = 0.94$  and  $0.91$ , resp.). Ten 2-(ar)alkoxyadenosines stimulated platelet AC at EC50s ranging between  $0.16$  and  $2.3 \mu M$  and inhibited platelet aggregation at EC50s ranging between  $2$  and  $30 \mu M$ . There were no correlations between the EC50s of the stimulation of platelet or PC 12 AC ( $r_2 = 0.08$  and  $0.06$ , resp.) or with the  $K_i$  of the inhibition of  $[3H]$ NECA binding to the A2AAR in rat striatum ( $r_2 = 0.02$ ). The EC50s of the stimulation of platelet AC correlated with those of the stimulation of PC 12 AC ( $r_2 = 0.48$ ), and also with the  $K_i$  of  $[3H]$ NECA binding ( $r_2 = 0.71$ ). Each of the 23 adenosines completely inhibited platelet aggregation and thus, functionally, all behaved as full agonists. As stimulants of PC 12 cell AC, Group A and B analogs were equally efficacious. As stimulants of platelet AC, however, the efficacy relative to NECA ( $= 1.0$ ) of Group B analogs was significantly less than that of Group A analogs,  $0.49 \pm 0.2$  vs.  $0.72 \pm 0.05$ ,  $P < 0.01$ . The partial agonist activity of Group B analogs at the platelet A2AR but full agonist activity at the PC 12 cell A2aAR, as well as the relatively low correlations between platelet AC stimulation and other indexes of A2AAR agonist activity, suggest the platelet receptor is not a typical A2aAR. Further, the lack of a correlation between the platelet anti-aggregation and AC stimulatory activity suggests that (a) the 2-(ar)alkoxyadenosines might affect platelet aggregation by mechanisms other than AC stimulation of (b) that the stimulation of the platelet membrane AC by 2-(ar)alkoxy-adenosines does not correspond to the accumulation of cAMP in intact platelets.

10/598,520


IT 50257-84-6 50257-89-1 53296-10-9  
131865-81-1  
RL: BIOL (Biological study)  
(platelet aggregation inhibition by)  
RN 50257-84-6 CAPLUS  
CN Adenosine, 2-butoxy- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 50257-89-1 CAPLUS  
CN Adenosine, 2-(2-hydroxyethoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



RN 131865-81-1 CAPLUS  
CN Adenosine, 2-(4-phenylbutoxy)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 39 THERE ARE 39 CAPLUS RECORDS THAT CITE THIS RECORD (39 CITINGS)

McIntosh

10/598,520

L6 ANSWER 93 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1994:500054 CAPLUS

DN 121:100054

OREF 121:17759a,17762a

TI A binding site model and structure-activity relationships for the rat A3 adenosine receptor

AU van Galen, Philip J. M.; van Bergen, Andrew H.; Gallo-Rodriguez, Carola; Melman, Neli; Olah, Mark E.; Ijzerman, Ad P.; Stiles, Gary L.; Jacobson, Kenneth A.

CS Lab. Bioorganic Chem., Natl. Inst. Diabetes, Digestive and Kidney Diseases, Bethesda, MD, 20892, USA

SO Molecular Pharmacology (1994), 45(6), 1101-11  
CODEN: MOPMA3; ISSN: 0026-895X

DT Journal

LA English

AB A novel adenosine receptor, the A3 receptor, has recently been cloned. The authors have systematically investigated the hitherto largely unexplored structure-activity relationships (SARs) for binding at A3 receptors, using 125I-N6-2-(4-aminophenyl)ethyladenosine as a radioligand and membranes from Chinese hamster ovary cells stably transfected with the rat A3-cDNA. As is the case for A1 and A2a receptors, substitutions at the N6 and 5' positions of adenosine, the prototypic agonist ligand, may yield fairly potent compds. However, the highest affinity and A3 selectivity is found for N6,5'-disubstituted compds., in contrast to A2 and A2a receptors. Thus, N6-benzyladenosine-5'-N-ethylcarboxamide is highly potent (Ki, 6.8 nM) and moderately selective (13- and 14-fold vs. A1 and A2a). The N6 region of the A3 receptor also appears to tolerate hydrophilic substitutions, in sharp contrast to the other subtypes. Potencies of N6,5'-disubstituted compds. in inhibition of adenylate cyclase via A3 receptors parallel their high affinity in the binding assay. None of the typical xanthine or nonxanthine (A1/A2) antagonists tested show any appreciable affinity for rat A3 receptors. 1,3-Dialkylxanthines did not antagonize the A3 agonist-induced inhibition of adenylate cyclase. A His residue in helix 6 that is absent in A3 receptors but present in A1/A2 receptors may be causal in this respect. In a mol. model for the rat A3 receptor, this mutation, together with an increased bulkiness of residues surrounding the ligand, make antagonist binding unfavorable when compared with a previously developed A1 receptor model. Second, this A3 receptor model predicted similarities with A1 and A2 receptors in the binding requirements for the ribose moiety and that xanthine-7-ribosides would bind to rat A3 receptors. This hypothesis was supported exptl. by the moderate affinity (Ki, 6 µM) of 7-riboside of 1,3-dibutylxanthine, which appears to be a partial agonist at rat A3 receptors. The model presented here which is consistent with the detailed SAR found in this study, may serve to suggest future chemical modification, site-directed mutagenesis, and SAR studies to further define essential characteristics of the ligand-receptor interaction and to develop even more potent and selective A3 receptor ligands.

IT 53296-10-9, 2-(Phenylamino)adenosine


RL: BIOL (Biological study)

(adenosine A1 and A2a and A3 receptors affinity for, mol. structure in relation to)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 113 THERE ARE 113 CAPLUS RECORDS THAT CITE THIS RECORD (114 CITINGS)

L6 ANSWER 94 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN


McIntosh



10/598,520

AN 1994:473772 CAPLUS  
DN 121:73772  
OREF 121:13003a,13006a  
TI Modulation of intraocular pressure by adenosine agonists  
AU Crosson, Craig E.; Gray, Tracy  
CS Health Sci. Cent., Texas Tech Univ., Lubbock, TX, USA  
SO Journal of Ocular Pharmacology (1994), 10(1), 379-83  
CODEN: JOPHER; ISSN: 8756-3320  
DT Journal  
LA English  
AB To investigate the potential role of adenosine receptors in modulating intraocular pressure (IOP), the A1 agonist N6-cyclopentyladenosine (CPA), the nonselective adenosine agonist 5'-N-ethylcarboxamidoadenosine (NECA) and the A2 agonist 8-phenylaminoadenosine (CV-1808) were evaluated. Topical administration of NECA to rabbits produced a dose-related reduction in IOP. However, an initial ocular hypertension of 1-2-h duration was also observed in rabbits treated with NECA. CPA (165 µg) caused only a reduction in IOP, while CV-1808 produced only an initial ocular hypertension. As adenosine A1 receptors have been shown to be neg. coupled to adenylate cyclase in several systems, CPA was evaluated for its ability to suppress cAMP formation in the isolated iris/ciliary body. CPA produced a concentration-related suppression of the cAMP accumulation induced by 10-6M forskolin (EC50 = 3.2 nM). These results indicate that selected adenosine agonists can modulate IOP. The ocular hypotension induced by adenosine agonists is consistent with the activation of adenosine A1 receptors and may involve the modulation of cAMP levels in the iris/ciliary body.  
IT 53296-10-9, CV 1808  
RL: BIOL (Biological study)  
(eye intraocular pressure response to)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 16 THERE ARE 16 CAPLUS RECORDS THAT CITE THIS RECORD (16 CITINGS)

L6 ANSWER 95 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1994:260573 CAPLUS  
DN 120:260573  
OREF 120:45825a,45828a  
TI Functional characterization of the A2b adenosine receptor in NIH 3T3 fibroblasts  
AU Brackett, L. Ellen; Daly, John E.  
CS Lab. Bioorg. Chem., Natl. Inst. Diabetes Dig. Kidney Dis., Bethesda, MD, 20892, USA  
SO Biochemical Pharmacology (1994), 47(5), 801-14  
CODEN: BCPCA6; ISSN: 0006-2952  
DT Journal  
LA English  
AB The adenosine (ADO) receptor in NIH 3T3 fibroblasts was characterized using a series of adenosine agonists and selected xanthine and non-xanthine antagonists. The ADO receptor elicited accumulations of cAMP in intact NIH 3T3 fibroblasts and caused activation adenylate cyclase in membrane preps. The receptor had characteristics of the A2b subtype of adenosine receptor. ADO analogs had relatively high EC50 values at the receptor and were antagonized competitively by xanthines. The rank order of potency for adenosine analogs in NIH 3T3 fibroblasts for cAMP accumulation was: NECA > 2-ClADO > R-PIA » CV1808, CGS 21680. The EC50 for 2-ClADO was 4.3 µM in intact cells and 15 µM in membrane preps. All ADO analogs were more potent at the A2a receptor of pheochromocytoma PC12 membranes than at the A2b receptor of fibroblast NIH

McIntosh

3T3 membranes. Structure-activity relationships suggested that the regions of interaction with 5'- and N6-substituents of ADO were similar for both the PC12 A2a and NIH 3T3 A2b receptor. However, ADO analogs with large substituents in the 2'-position, such as 2-cyclohexylethoxyADO and CGS 21680, were highly selective for the A2a receptor. All ADO analogs tested were stimulatory to adenylate cyclase at the NIH 3T3 A2b receptor, including 5'-methylthioADO, which was a weak partial agonist. A series of xanthine antagonists were not selective for the NIH 3T3 A2b vs. the PC12 A2a receptor. In all cases, xanthines were more potent as antagonist in the intact NIH 3T3 cells than in NIH 3T3 membranes. In a series of non-xanthine antagonists, most compds. were equipotent or slightly more potent at the A2a receptor except for alloxazine, which was approx. 9-fold selective for the A2b receptor.


IT 53296-10-9, CV1808

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)  
(adenosine receptor agonist activity of, in fibroblasts and pheochromocytoma cells, structure in relation to)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 90 THERE ARE 90 CAPLUS RECORDS THAT CITE THIS RECORD (90 CITINGS)

L6 ANSWER 96 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1994:236896 CAPLUS

DN 120:236896

OREF 120:41761a,41764a

TI Discrimination of A1 versus A2 receptor subtype selectivity of adenosine receptor agonists in vivo

AU Barrett, Richard J.; Droppleman, David A.; Wright, Kathryn F.

CS Dep. Pharmacol., Whitby Res., Inc., Richmond, VA, USA

SO Journal of Pharmacology and Experimental Therapeutics (1994), 268(3), 1166-73

CODEN: JPETAB; ISSN: 0022-3565

DT Journal

LA English

AB Previous attempts to discern and quantify the selectivity of agonists for A1 vs. A2 adenosine receptors in vivo have been confounded by the activation of baroreceptor reflexes and/or simultaneous expression of responses to both A1 and A2 receptor activation. In anesthetized, vagotomized rats with isolated in situ constant-flow perfused hindquarters (HQ), bradycardic responses to i.v. agonist injections measured A1 receptor activation and HQ vasodilation elicited by i.a. agonist injections measured the stimulation of A2 receptors. Adenosine and 5'-N-ethylcarboxamidoadenosine (NECA) produced A2 receptor-mediated HQ vasodilation at doses 8- and 4-fold lower (-log ED50 values, 7.3 mol and 8.7 mol, resp.) than those required to evoke A1 receptor-mediated bradycardia (-log ED50 values, 6.4 mol and 8.1 mol, resp.). N6-cyclopentyladenosine (CPA) was approx. 8-fold selective for A1 receptors (-log ED50 values, A1, 8.5 mol; A2, 7.6 mol). 2-(Phenylamino)adenosine (CV-1808) and 2[2(4-fluorophenyl)ethoxy]adenosine (FPEA) were at least 125- and 200-fold more potent agonists at A2 receptors (-log ED50 values, 7.7 mol and 8.0 mol, resp.) than at A1 receptors (-log ED50 values, 5.6 mol and 5.7 mol, resp.). These studies demonstrated that stimulation of A1 and A2 receptors may be discriminated in vivo and that such responses are selective, reproducible, dose-dependent and quantifiable. A comparison of these in vivo measures with known in vitro data suggests that the A2a adenosine receptor mediates vasodilation in the rat HQ and that in vitro assays may predict the orders

of potency of adenosine A1 and A2 receptor agonists in vivo but they are less reliable predictors of the absolute potency and, hence, the A1/A2 receptor selectivity of agonists.

IT 53296-10-9, 2-(Phenylamino)adenosine


RL: BIOL (Biological study)

(purinergic A1 and A2 selectivity of, bradycardia and hindquarter vasodilation in discrimination of)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 4 THERE ARE 4 CAPLUS RECORDS THAT CITE THIS RECORD (4 CITINGS)

L6 ANSWER 97 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1994:125921 CAPLUS

DN 120:125921

OREF 120:22025a,22028a

TI Molecular cloning and functional expression of a sheep A3 adenosine receptor with widespread tissue distribution

AU Linden, Joel; Taylor, Heidi E.; Robeva, Anna S.; Tucker, Amy L.; Stehle, Jorg H.; Rivkees, Scott A.; Fink, J. Stephen; Reppert, Steven M.

CS Lab. Dev. Chronobiol., Massachusetts Gen. Hosp., Boston, MA, 02114, USA

SO Molecular Pharmacology (1993), 44(3), 524-32

CODEN: MOPMA3; ISSN: 0026-895X

DT Journal

LA English

AB Using the polymerase chain reaction, an A3 adenosine receptor has been cloned from the hypophyseal par tuberalis of sheep. The clone encodes a 317-amino acid protein that is 72% identical to the rat A3 adenosine receptor. In contrast to rat, where abundant A3 mRNA transcript is found primarily in testis, the sheep transcript is most abundant in lung, spleen, and pineal gland and is present in moderate levels in brain, kidney, and testis. The agonist N6-amino[125I]iodobenzyladenosine binds with high affinity (Kd .simeq. 6 nM) and specificity to recombinant A3 adenosine receptors expressed transiently in COS-1 cells or stably in CHO K1 cells. The potency order of agonists is N6-aminiodobenzyladenosine > N-ethylcarboxamidoadenosine ≥ (R)-phenylisopropyladenosine » cyclopentyladenosine. Little or no binding of purine nucleotides was detected. The potency order of antagonists is 3-(3-iodo-4-aminobenzyl)-8-(4-oxyacetate)phenyl-1-propylxanthine (I-ABOPX) (Ki = 3 nM) > 1,3-dipropyl-8-(4-acrylate)phenylxanthine (BW-A1433) > 1,3-dipropyl-8-sulfophenylxanthine = xanthine amine congener » 8-cyclopentyl-1,3-dipropylxanthine. Enprofylline does not bind. These data indicate that, in contrast to A1 adenosine receptors, A3 adenosine receptors preferentially bind ligands with aryl rings in the N6-position of adenine and in the C8-position of xanthine. Among antagonists, the A3 adenosine receptor preferentially binds 8-phenylxanthines with acidic vs. basic para-substituents (I-ABOPX > BW-A1433 > 1,3-dipropyl-8-sulfophenylxanthine = xanthine amine congener). Agonists reduce forskolin-stimulated cAMP accumulation in Chinese hamster ovary cells stably transfected with recombinant sheep A3 adenosine receptors; the reduction is blocked by BW-A1433 but not by 8-cyclopentyl-1,3-dipropylxanthine. These data suggest that (i) A3 adenosine receptors display unusual structural diversity for species homologs, (ii) in contrast to rat, sheep A3 adenosine receptors have a broad tissue distribution, and (iii) some xanthines with acidic side chains bind with high affinity to A3 adenosine receptors.

IT 53296-10-9, CV1808


RL: BIOL (Biological study)

(binding to sheep A3 adenosine receptor of)

10/598,520

RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 137 THERE ARE 137 CAPLUS RECORDS THAT CITE THIS RECORD (140 CITINGS)

L6 ANSWER 98 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1994:729 CAPLUS

DN 120:729

OREF 120:167a,170a

TI Structure-activity relationship of 2-(ar)alkoxyadenosines at the adenosine A2 receptor in coronary artery

AU Makujina, Shah R.; Olsson, Ray A.; Esinhart, James D.; Mustafa, S. Jamal

CS Sch. Med., East Carolina Univ., Greenville, NC, 27858-4354, USA

SO European Journal of Pharmacology (1993), 243(1), 35-8

CODEN: EJPHAZ; ISSN: 0014-2999

DT Journal

LA English

AB The authors examined the ability of four 2-(ar)alkoxyadenosines [2-(2-phenylethoxy)adenosine, PEA; 2-[2-(2-naphthyl)ethoxy]adenosine, NEA; 2-[2-(4-methylphenyl)ethoxy]adenosine, mPEA; and 2-(1-hexyloxy)adenosine, HOA] to relax porcine coronary artery in vitro. All four compds. produced concentration-dependent relaxations in rings contracted with 30 mM KCl. The EC<sub>25</sub> values are as follows (+ 10<sup>-9</sup> mol/L): CGS21680, (2-[p-(2-carboxyethyl)phenethylamino]-5'-N-ethylcarboxamidoadenosine) (32.7) ≈ NECA, 5'-N-ethylcarboxamidoadenosine (51.4) ≈ mPEA (74.3) ≈ NEA (160.7) > HOA (855.1) ≈ PEA (1259) ≈ 2-chloroadenosine (1871) > adenosine (9705). However, EC<sub>75</sub> values for all the compds. except adenosine and 2-chloroadenosine converged to a range of 8.16 to 22.86 μM, suggesting a biphasic response. Furthermore, the responses were found to be independent of endothelial integrity. The unselective adenosine receptor antagonist 8-p-sulfophenyltheophylline (100 μM) attenuated the relaxant response to NEA (EC<sub>25</sub> = 1172 nM), suggesting that adenosine receptors mediated relaxation. Structure-activity correlations suggest that the adenosine A2 receptor in porcine coronary artery contains a region of limited bulk tolerance juxtaposed to the region occupied by adenine C-2 and distal to that a large hydrophobic region.

IT 50257-95-9, 2-(1-Hexyloxy)adenosine


RL: BIOL (Biological study)

(coronary artery relaxation by, structure in relation to)

RN 50257-95-9 CAPLUS

CN Adenosine, 2-(hexyloxy)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 6 THERE ARE 6 CAPLUS RECORDS THAT CITE THIS RECORD (6 CITINGS)

McIntosh

10/598,520

L6 ANSWER 99 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1993:595546 CAPLUS

DN 119:195546

OREF 119:34637a,34640a

TI Adenosine agonists reduce conditioned avoidance responding in the rat

AU Martin, Gregory E.; Rossi, Donald J.; Jarvis, Michael F.

CS Dep. Pharmacol., Rhone-Poulenc Rorer Cent. Res., Collegeville, PA, 19426, USA

SO Pharmacology, Biochemistry and Behavior (1993), 45(4), 951-8

CODEN: PBBHAU; ISSN: 0091-3057

DT Journal

LA English

AB Because adenosine agonists may possess therapeutic potential as antipsychotic agents, the authors examined the activity of several prototypic agents in vivo in blocking conditioned avoidance (CAR) in the rat, a behavioral test predictive of antipsychotic efficacy in humans. Potency in blocking CAR is directly proportional to potency in alleviating schizophrenia. Hence, the adenosine A1-selective agonists [cyclopentyl adenosine (CPA) and (R)-phenylisopropyl adenosine (R-PIA)], A2-selective agonists [CV-1808 and (2-p(carboxyethyl)-(NECA)] were examined in this test. Block of CAR was first determined for standard antipsychotic agents [ED50 mg/kg, IP, and 95% confidence level (CL) in parentheses], such as haloperidol [0.23 (0.18, 0.39)], trifluoperazine [(0.9 (0.7, 1.0)], thioridazine [12.5 (10.5, 15.3)], metoclopramide [7.8 (6.4, 9.2)], and chlorpromazine [4.9 (4.2, 5.9)]. The paradigm consisted of a light- and tone-signalized footshock that could be avoided via a discrete lever press. Affinity for A1 and A2 binding sites in brain tissue from Fischer 344 rats was ascertained to be similar to that seen in other rodent strains. Each adenosine agonists blocked CAR. NECA [ED50 value (95% CL) = 0.07 (0.004, 0.12) mg/kg, IP] was the most potent agent, followed by: R-PIA [0.34 (0.23, 0.44)]; CGS 21680 [1.1 (0.8, 2.0)]; CV-1808 [1.3 (1.0, 1.8)]; and CPA [1.5 (1.3, 1.7)]. Pretreatment with caffeine (25 mg/kg, IP, -10 min) blocked the inhibition of CAR produced by adenosine agonists, suggesting the event is mediated via purinergic receptors. As a test for extrapyramidal side effect potential, each agonist was administered at dose levels corresponding to the ED5, ED50, and ED75 values for block of CAR and catalepsy was measured. Catalepsy was prominently produced by NECA and CPA, whereas CGS 21680 and R-PIA produced little. Neither potency in blocking CAR nor inducing catalepsy could be highly correlated with either relative affinity or selectivity for either A1 or A2 binding sites. The data suggest purinergic agonists might be effective antipsychotic agents but may possess side effects that might preclude their use.

IT 53296-10-9, CV1808


RL: BIOL (Biological study)

(conditioned avoidance responding reduction by, antipsychotic activity in relation to)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 26 THERE ARE 26 CAPLUS RECORDS THAT CITE THIS RECORD (26 CITINGS)

L6 ANSWER 100 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1993:532153 CAPLUS

DN 119:132153

OREF 119:23521a,23524a

TI Functional characterization of three adenosine receptor types

AU Gurden, M. F.; Coates, J.; Ellis, F.; Evans, B.; Foster, M.; Hornby, E.; Kennedy, I.; Martin, D. P.; Strong, P.; et al.

McIntosh

10/598,520

CS Pharmacol. Div., Glaxo Group Res., Ware/Hertfordshire, SG12 0DP, UK

SO British Journal of Pharmacology (1993), 109(2), 693-8

CODEN: BJPCBM; ISSN: 0007-1188

DT Journal

LA English

AB The purpose of the present study was to classify adenosine receptors into A1 and A2 subtypes in a wide range of isolated tissues and cell types (rat adipocytes and atria, guinea pig ileum and atria (A1); guinea pig aorta, dog coronary artery and human platelets and neutrophils (A2)) using the R- and S-diastereoisomers of N-phenylisopropyladenosine (PIA), N-cyclopentyladenosine (CPA), the novel compound, N-[(1S,trans)-2-hydroxycyclopentyl]adenosine (GR 79236), N-[(2-methylphenyl)methyl]adenosine (metrifudil), 2-(phenylamino)adenosine (CV 1808), and 2-[[2-[4-(2-carboxyethyl)phenyl]ethyl]amino]-N-ethylcarboxamidoadenosine (CGS 21680); N-ethylcarboxamidoadenosine (NECA) was used as a standard. Results obtained in all tissue preps. previously reported to contain A1-receptors could be described by a single rank order of agonist potency: CPA  $\geq$  GR 79236, R-PIA  $\geq$  NECA  $\gg$  S-PIA  $\geq$  metrifudil  $\geq$  CV 1808, CGS 21680. In contrast, 2 distinct rank orders of agonist potency were observed in preps. previously reported to contain A2-receptors. In dog coronary artery, human neutrophils and platelets the rank order of potency was: CV 1808, CGS 21680  $\geq$  NECA  $>$  R-PIA  $\geq$  metrifudil  $\geq$  CPA  $>$  GR 79236, S-PIA. However, in guinea pig aorta the rank order was: NECA  $>$  metrifudil  $>$  R-PIA, CPA  $>$  CV 1808, GR 79236  $\geq$  S-PIA, CGS 21680. The results indicate the existence of 3 types of adenosine receptor: A1- and 2 subtypes of A2-receptor. The receptor present in dog coronary artery, human platelets and neutrophils, probably corresponds to the A2a subtype, whilst that present in the guinea-pig aorta may be of the A2b subtype.

IT 53296-10-9


RL: BIOL (Biological study)

(adenosine receptor subtype classification using, in human and laboratory animal tissues)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 74 THERE ARE 74 CAPLUS RECORDS THAT CITE THIS RECORD (74 CITINGS)

L6 ANSWER 101 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1993:473033 CAPLUS

DN 119:73033

OREF 119:13185a,13188a

TI Preparation of 2-alkynyladenosine derivatives having high affinity for adenosine A2 receptor

IN Matsuda, Akira; Yamaguchi, Toyofumi; Miyashita, Takanori; Azebiru, Toichi; Watanabe, Yoko; Goto, Takao; Kojo, Kentaro; Narita, Senichi

PA Yamasa Shoyu Co., Ltd., Japan; Toa Eiyo, Ltd.

SO Jpn. Kokai Tokkyo Koho, 8 pp.

CODEN: JKXXAF


DT Patent

LA Japanese

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 05009197	A	19930119	JP 1991-158991	19910628
	JP 3053908	B2	20000619		
PRAI	JP 1991-158991		19910628		
OS	MARPAT 119:73033				
GI					

McIntosh



AB The title compds. [I; R = C(OH)R<sub>1</sub>R<sub>2</sub>, (CH<sub>2</sub>)<sub>n</sub>OR<sub>1</sub>, (CH<sub>2</sub>)<sub>n</sub>NR<sub>1</sub>R<sub>2</sub>, (CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>R<sub>1</sub>, COR<sub>1</sub>, (CH<sub>2</sub>)<sub>n</sub>R<sub>3</sub>; R<sub>1</sub>, R<sub>2</sub> = H, alkyl; n = 0-10; R<sub>3</sub> = alkenyl, alkynyl, aryl, N<sub>3</sub>, cyanol, having highly selective affinity to adenosine A<sub>2</sub> receptor and very low affinity to A<sub>1</sub> receptor (no data), are prepared I have antihypertensive, (peripheral) vasodilatory, and blood platelet aggregation-inhibitory activities without inhibition of heart and central nervous system activity, and are useful for the treatment of hypertension and ischemic diseases (e.g. heart and brain ischemia). Thus, 2 mmol 2-iodoadenosine was dissolved in DMF, thereto 70 mg [Ph<sub>3</sub>P]2PdCl<sub>2</sub>, 38 mg CuI, 1.4 mL Et<sub>3</sub>N, and 2-5 mmol acetylene compound RC.tplbond.CH were added, and the mixture was allowed to react at 70-120°, concentrated, dissolved in MeOH, treated with H<sub>2</sub>S, and purified by silica gel chromatog. to give I. A total of 21 I [e.g. R = Me<sub>2</sub>C(OH)C.tplbond.C, Me<sub>2</sub>NCH<sub>2</sub>C.tplbond.C, HO<sub>2</sub>C(CH<sub>2</sub>)<sub>7</sub>C.tplbond.C, 4-EtO<sub>2</sub>CC<sub>6</sub>H<sub>4</sub>(CH<sub>2</sub>)<sub>4</sub>C.tplbond.C] were prepared


IT 149009-12-1P

RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of, with selective adenosine A<sub>2</sub> receptor affinity)

RN 149009-12-1 CAPLUS

CN Adenosine, 2-(6-amino-1-hexynyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

L6 ANSWER 102 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1993:463783 CAPLUS

DN 119:63783

OREF 119:11297a,11300a

TI Characterization of adenosine A<sub>2</sub> receptors in bovine retinal pigment epithelial membranes

AU Blazynski, Christine

CS Sch. Med., Washington Univ., St. Louis, MO, 63110, USA

SO Experimental Eye Research (1993), 56(5), 595-9

CODEN: EXERA6; ISSN: 0014-4835

DT Journal

LA English

AB The pharmacol. characteristics of adenosine A<sub>2</sub> receptors are described for membranes prepared from bovine retinal pigmented epithelial (RPE). RPE cells were isolated after removal of retina, lysed by freeze-thawing, and membranes separated from cytoplasmic components. A single population of adenosine binding sites is present in RPE membranes, as determined from saturation anal. and competition binding assays. From Scatchard plots, this single class of binding sites exhibited low affinity for adenosine receptor agonists. These low affinity sites were labeled by

[3H]-N-ethylcarboxamido-adenosine (NECA) or [3H]-CGS 21680 and Kds of 423 and 5.3  $\mu\text{M}$  were determined for each radioligand, resp. NECA-mediated stimulation of adenylate cyclase demonstrated that these binding sites represent adenosine receptors. No high affinity A2a binding sites were detected in RPE membranes by either saturation studies, or by competition with adenosine A1-selective agonists which only displaced radioligand binding at high micromol. concns. The low affinity A2 receptor on RPE differs from the high affinity A2a receptor characterized in bovine retinal membranes, but may be similar or identical to the lower affinity A2b receptor detected in retinal membranes as well as other tissues.

IT 53296-10-9, CV1808


RL: BIOL (Biological study)

(adenosine A2b receptors affinity for, of retina pigment epithelial cells)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 14 THERE ARE 14 CAPLUS RECORDS THAT CITE THIS RECORD (14 CITINGS)

L6 ANSWER 103 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1993:463782 CAPLUS

DN 119:63782

OREF 119:11297a,11300a

TI Characterization of adenosine A2 receptors in bovine retinal membranes

AU Blazynski, Christine; McIntosh, Helen

CS Sch. Med., Washington Univ., St. Louis, MO, 63110, USA

SO Experimental Eye Research (1993), 56(5), 585-93

CODEN: EXERA6; ISSN: 0014-4835

DT Journal

LA English

AB Bovine retinal A2 receptors were characterized based on data obtained from both adenylate cyclase assays and radioligand binding studies. [3H]-5'-N-ethylcarboxamidoadenosine (NECA) in the presence of 10 nM cyclopentyladenosine (CPA, which selectively binds to A1 receptors) or [3H]-CGS 21680 were used to label the A2 binding sites. By using [3H]-NECA (plus CPA), two populations of binding sites, having Kds of 106 nM and 9.4  $\mu\text{M}$ , were determined [3H]-CGS 21680, a derivative of NECA which has been demonstrated to be highly selective for A2 receptors in brain synaptic membrane preps. was more potent than NECA at the higher affinity population of A2 sites, and saturation anal. revealed the presence of both a high affinity site, Kd of 18 nM, and a lower affinity site having a Kd of 4.3  $\mu\text{M}$ . The high affinity site labeled by [3H]-CGS 21680 corresponds to the A2a receptor. By using either radioligand, guanosine triphosphate-dependent shifts to a single population of binding sites were observed. Despite the differences in affinities revealed by the two radioligands for the high affinity A2 site, both [3H]-CGS 21680 and [3H]-NECA were competitively displaced by increasing concns. of a variety of adenosine receptor agonists and antagonists, and exhibited an identical rank order of potency that is consistent with that reported for high affinity A2a receptors. Receptor-mediated modulation of adenylate cyclase activities in retinal synaptic membranes was also assessed, and while NECA or N6-methyladenosine elicited decreases in forskolin-activated cyclase activity at concns. between 0.1-50 nM, this inhibition was reversed, and enzyme stimulated by higher agonist concns. CGS 21680 elicited only a stimulation of either basal and forskolin-activated adenylate cyclase activities at concns. above 50 nM. The stimulatory modulation of adenylate cyclase at these concns. is consistent with mediation by the A2a and/or A2b receptors.


IT 53296-10-9, CV1808



10/598,520

RL: BIOL (Biological study)  
(adenosine A2 receptors affinity for, of eye retina membranes)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)


Absolute stereochemistry.



OSC.G 12 THERE ARE 12 CAPLUS RECORDS THAT CITE THIS RECORD (12 CITINGS)

L6 ANSWER 104 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1993:188505 CAPLUS  
DN 118:188505  
OREF 118:32315a,32318a  
TI Effects of adenosine derivatives on human and rabbit platelet aggregation.  
Correlation of adenosine receptor affinities and antiaggregatory activity  
AU Dionisotti, Silvio; Zocchi, Cristina; Varani, Katia; Borea, Pier Andrea;  
Ongini, Ennio  
CS Res. Lab., Schering-Plough, S.p.A., Comazzo, I-20060, Italy  
SO Naunyn-Schmiedeberg's Archives of Pharmacology (1992), 346(6), 673-6  
CODEN: NSAPCC; ISSN: 0028-1298  
DT Journal  
LA English  
AB The inhibitory effects of several adenosine analogs, including the  
A2-selective agonists 2-[p-(2-carboxyethyl)phenylethylamino]-5'-N-  
ethylcarboxamidoadenosine (CGS 21680) and  
2-hexynyl-5'-N-ethylcarboxamidoadenosine (2-hexynyl-NECA), were  
investigated in vitro on human and rabbit platelet aggregation. The  
compds. inhibited ADP-induced platelet aggregation over a wide range of  
potency. The rank order of activity was similar between the 2 species  
thus showing that the rabbit is a useful animal model for studying the  
effects of adenosine derivs. on platelet aggregation. 2-Hexynyl-NECA was  
the most potent adenosine compound of those currently available, having IC50  
values of 0.10 and 0.07  $\mu$ M in human and rabbit platelets, resp.  
Conversely, the A1 agonists R(-)-N6-(2-phenylisopropyl)adenosine,  
S(+)-N6-(2-phenylisopropyl)adenosine, and 2-chloro-N6-cyclopentyladenosine  
were the least potent compds. with IC50 values in the micromolar range.  
The potency of the compds. in inhibiting platelet aggregation was  
correlated with their affinity for A2 receptors as measured using [3H]CGS  
21680 binding in rat brain striatum.  
IT 53296-10-9, CV 1808  
RL: BIOL (Biological study)  
(blood platelet aggregation inhibition by, in human and rabbit,  
adenosine A2 receptor affinity in relation to)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 24 THERE ARE 24 CAPLUS RECORDS THAT CITE THIS RECORD (24 CITINGS)

McIntosh

L6 ANSWER 105 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1993:74085 CAPLUS

DN 118:74085

OREF 118:12831a,12834a

TI [3H]2-Phenylaminoadenosine ([3H]CV 1808) labels a novel adenosine receptor in rat brain

AU Cornfield, Linda J.; Hu, Shiling; Hurt, Stephen D.; Sills, Matthew A.

CS Pharm. Div., CIBA-GEIGY Corp., Summit, NJ, USA

SO Journal of Pharmacology and Experimental Therapeutics (1992), 263(2), 552-61

CODEN: JPETAB; ISSN: 0022-3565

DT Journal

LA English

AB After the radiolabeling of CV 1808, its binding characteristics were evaluated in rat striatal, cortical and hippocampal membranes. Using 5 nM [3H]CV 1808, unlabeled CV 1808 produced shallow inhibition curves in all three brain areas, with 61-75% of the binding displaying IC50 values of 16-24 nM, whereas the remaining 28-37% of binding had lower affinity (IC50 595-1130 nM). The A2-selective agonist CGS 21680 and the nonselective adenosine agonist 5'-N-ethylcarboxamidoadenosine displayed very low affinity (IC50 > 10  $\mu$ M). The A1-selective compound N6-cyclopentyladenosine inhibited only 28-44% of specific binding, with IC50 of 272-1750 nM. In contrast, the nonselective adenosine antagonist CGS 15943A inhibited specific binding by 48-64% (at 1  $\mu$ M) with IC50 ranging 106-295 nM. Addnl., several novel adenosine analogs fully inhibited specific binding, producing multicomponent inhibition curves. Electrophysiol. studies in porcine coronary artery cells demonstrated that CV 1808, but not CGS 21680, 5'-N-ethylcarboxamidadenosine and N6-cyclopentyladenosine, activated potassium channels. Further, the CV 1808-induced activation was blocked by CGS 15943A. Thus, [3H]CV 1808 binding consists of two components in rat brain a low-affinity site with A1-like characteristics, and a novel high-affinity site, designated as the A4 receptor, where potassium channel activation appears to be a functional correlate.

IT 53296-10-9, CV 1808 53296-10-9D, CV 1808, derivs., tritium-labeled


RL: BIOL (Biological study)

(adenosine receptor subtypes labeling with, in brain)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)


Absolute stereochemistry.



RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



10/598,520

OSC.G 26 THERE ARE 26 CAPLUS RECORDS THAT CITE THIS RECORD (26 CITINGS)

L6 ANSWER 106 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1993:73788 CAPLUS

DN 118:73788

OREF 118:12763a,12766a

TI Structure-activity relationships for 2-substituted adenosines at A1 and A2 adenosine receptors

AU Daly, John W.; Padgett, William L.; Secunda, Sherrie I.; Thompson, Robert D.; Olsson, Ray A.

CS Lab. Bioorg. Chem., Natl. Inst. Diabetes Dig. Kidney Dis., Bethesda, MD, USA

SO Pharmacology (1993), 46(2), 91-100

CODEN: PHMGBN; ISSN: 0031-7012

DT Journal

LA English

AB A series of 55 2-alkyloxy-, 2-aryloxy-, and 2-aralkyloxyadenosines was screened as inhibitors of the binding of [3H]R-phenyl-isopropyladenosine to A1 adenosine receptors in rat cerebral cortical membranes, as inhibitors of the binding of [3H]N-ethylcarboxyamidoadenosine to A2 adenosine receptors in rat striatal membranes, and as agonists at A2 adenosine receptors coupled to adenylate cyclase in rat pheochromocytoma PC12 cell membranes. The activities are consonant with a hydrophobic binding site in the A2 receptors at a distance from the 2-position of the adenine ring corresponding to a spacer chain of -O-CH2-CH2-. There is little lateral steric tolerance in the region occupied by the spacer chain. Interaction with the hydrophobic binding site is greatest in the 2-alkyloxy series for 2-cyclohexylethoxy-, 2-cyclohexylpropoxy- and 2-cyclohexylbutoxyadenosine and in the 2-aralkyloxy series for 2-phenylethoxy-, 2-(4-methylphenyl)ethoxy-, 2-(4-chlorophenyl)ethoxy-, and 2-naphthylethoxyadenosine. The affinities of the 2-substituted adenosines for the rat cerebral cortical A1 receptors are not as markedly altered by structural changes, and in almost all cases are 2-100-fold less than the affinity of the 2-substituted adenosines for the rat striatal A2 receptor. There is excellent correspondence of the present data on rat A2 receptors with reported potencies of these 2-substituted adenosines as coronary vasodilators in guinea pig heart preps.

IT 50257-82-4 50257-84-6 50257-85-7  
50257-89-1 50257-95-9 131865-78-6  
131865-81-1 131865-82-2 131933-15-8  
131933-17-0 131933-20-5 131933-26-1  
131933-27-2 131973-26-7 137817-83-5  
137817-84-6 145747-85-9 145747-86-0  
145747-87-1


RL: PRP (Properties)

(purinergic receptor affinity of, structure in relation to)

RN 50257-82-4 CAPLUS

CN Adenosine, 2-phenoxy- (CA INDEX NAME)

Absolute stereochemistry.




RN 50257-84-6 CAPLUS

CN Adenosine, 2-butoxy- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

10/598,520




RN 50257-85-7 CAPLUS  
CN Adenosine, 2-(pentyloxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 50257-89-1 CAPLUS  
CN Adenosine, 2-(2-hydroxyethoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 50257-95-9 CAPLUS  
CN Adenosine, 2-(hexyloxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 131865-78-6 CAPLUS  
CN Adenosine, 2-(phenylmethoxy)- (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 131865-81-1 CAPLUS  
CN Adenosine, 2-(4-phenylbutoxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 131865-82-2 CAPLUS  
CN Adenosine, 2-[(5-phenylpentyl)oxy]- (CA INDEX NAME)

Absolute stereochemistry.



RN 131933-15-8 CAPLUS  
CN Adenosine, 2-(3-methylbutoxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 131933-17-0 CAPLUS  
CN Adenosine, 2-(cyclohexylmethoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 131933-20-5 CAPLUS  
CN Adenosine, 2-(4-cyclohexylbutoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 131933-26-1 CAPLUS  
CN Adenosine, 2-(2-ethylbutoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 131933-27-2 CAPLUS  
CN Adenosine, 2-[(3-ethylpentyl)oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 131973-26-7 CAPLUS  
CN Adenosine, 2-[(4-methylpentyl)oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 137817-83-5 CAPLUS  
CN Adenosine, 2-[(2R)-2-phenylbutoxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 137817-84-6 CAPLUS  
CN Adenosine, 2-[(2S)-2-phenylbutoxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 145747-85-9 CAPLUS  
CN Adenosine, 2-[(3,6-dimethyl-6-heptenyl)oxy]-, (S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 145747-86-0 CAPLUS  
CN Adenosine, 2-[(3,6-dimethyl-6-heptenyl)oxy]-, (R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

McIntosh



RN 145747-87-1 CAPLUS  
 CN Adenosine, 2-[(3S)-3-phenylbutoxy]- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 21 THERE ARE 21 CAPLUS RECORDS THAT CITE THIS RECORD (21 CITINGS)

L6 ANSWER 107 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1993:32762 CAPLUS

DN 118:32762

OREF 118:5811a,5814a

TI Cardiovascular selectivity of adenosine receptor agonists in anesthetized dogs

AU Gerencer, R. Z.; Finegan, B. A.; Clanachan, A. S.

CS Dep. Pharmacol., Univ. Alberta, Edmonton, AB, T6G 2H7, Can.

SO British Journal of Pharmacology (1992), 107(4), 1048-56

CODEN: BJPCBM; ISSN: 0007-1188

DT Journal

LA English

AB The relevance of adenosine (Ado) receptor classification obtained from in vitro methods to the cardiovascular actions of Ado agonists in vivo was studied. The cardiovascular effects of AMP, N6-cyclohexyladenosine (CHA, 400-fold A1-selective), 5'-N-ethylcarboxamidoadenosine (NECA, A1 ≈ A2), and 2-phenylaminoadenosine (PAA, 5-fold A2-selective) were compared in open-chest, fentanyl-pentobarbitone anesthetized dogs. Graded doses of CHA (10 to 1000 µg/kg), NECA (0.5 to 100 µg/kg), or PAA (0.1 to 20 µg/kg) were administered i.v. and changes in hemodynamics and myocardial contractility were assessed 10 min following each dose. The effects of graded infusions of AMP (200 to 1000 µg/kg·min) were also evaluated. AMP and the Ado analogs (NECA > PAA > CHA) increased the systemic vascular conductance index (SVCI) in a dose-dependent manner and reduced the mean arterial pressure (MAP). At doses causing similar increases in SVCI, the agonists caused similar reflex increases in heart rate (HR) and cardiac index (CI) and decreases in AV conduction interval (AVi), and increased coronary vascular conductance (CVC). After cardiac autonomic blockade with atropine (0.2 mg/kg) and propranolol (1 mg/kg), AMP, CHA, and PAA still increased SVCI and CVC and decreased MAP. CHA and PAA had no marked effects on HR, CI, or AVi. As in the absence of cardiac autonomic blockade, equieffective vasodilator doses of CHA and PAA had identical effects on CVC, CI, and AVi. The myocardial contractility, as assessed by Emax, was stimulated by AMP in control animals. Following cardiac autonomic blockade, PAA increased the contractility while AMP and CHA had no effects. Despite marked differences in receptor selectivity in vitro, no marked differences between the actions of these A1- and A2-selective Ado receptor agonists on the cardiovascular system in vivo were apparent. Difficulties therefore exist in the application of in vitro Ado receptor selectivity data to the prediction of the cardiovascular effects of Ado agonists in vivo.


IT 53296-10-9, 2-Phenylaminoadenosine



10/598,520

RL: PRP (Properties)  
(cardiovascular effects of, in vitro and in vivo correlation of)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)

L6 ANSWER 108 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1992:658206 CAPLUS  
DN 117:258206  
OREF 117:44527a,44530a  
TI Stable solid 2-octynyladenosine  
IN Morozumi, Manami; Kumagai, Masao; Yamaguchi, Toyofumi  
PA Yamasa Shoyu K. K., Japan  
SO Eur. Pat. Appl., 22 pp.  
CODEN: EPXXDW  
DT Patent  
LA English  
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	EP 499185	A1	19920819	EP 1992-102210	19920210
	EP 499185	B1	19950517		
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, MC, NL, PT, SE				
	JP 05194579	A	19930803	JP 1991-40894	19910212
	JP 10081697	A	19980331	JP 1997-281349	19910212
	JP 3040742	B2	20000515		
	CA 2060865	A1	19920813	CA 1992-2060865	19920207
	CA 2060865	C	20000516		
	ES 2073793	T3	19950816	ES 1992-102210	19920210
	US 5939543	A	19990817	US 1992-833718	19920211
PRAI	JP 1991-40894	A	19910212		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB Stable low-moisture 2-octynyladenosine (I) ( $\leq 3\%$  moisture) is prepared. Another stable form consists of I deposited from  $\mu\text{m}$  EtOH solution containing  $< 1\%$  water. I-H<sub>2</sub>O was vacuum-dried with P2O<sub>5</sub>, at 80°, to 0.93% water content. The product showed only 0.8% decomposition when stored for 30 days at 25° and 30% relative humidity. I is an antihypertensive (no data).


IT 137896-07-2

RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(stable solid form of, low-moisture)

RN 137896-07-2 CAPLUS

CN Adenosine, 2-octyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



McIntosh

10/598,520

OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

L6 ANSWER 109 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1992:483920 CAPLUS

DN 117:83920

OREF 117:14479a,14482a

TI Relative agonist potencies of C2-substituted analogs of adenosine: evidence for adenosine A2B receptors in the guinea pig aorta

AU Martin, Pauline L.

CS Dep. Pharmacol., Whitby Res., Inc., Richmond, VA, 23220, USA

SO European Journal of Pharmacology (1992), 216(2), 235-42

CODEN: EJPHAZ; ISSN: 0014-2999

DT Journal

LA English

AB Nine C2-substituted adenosine analogs that are potent and selective for the A2-adenosine receptor were tested for their ability to induce relaxations of the guinea pig aorta. Compds. tested were 2-phenylethoxyadenosine (PEA), 2-phenylethoxy-5'-N-ethylcarboxyamidoadenosine (PENECA), 2-cyclohexylethoxyadenosine (CEA), 2-fluorophenylethoxyadenosine (FPEA), 2-methoxyphenylethoxyadenosine (MPEA), 2-naphthylethoxyadenosine (NEA), 2-phenylaminoadenosine (CV1808), 2-phenylethylaminoadenosine (PEAA), and 2-carboxyethylphenethylamino-5'-N-ethylcarboxyamidoadenosine (CGS21680). The responses to these agents were compared with those to three standard adenosine receptor agonists, 5'-N-ethylcarboxyamidoadenosine (NECA), N6-cyclohexyladenosine (CHA) and R-N6-phenylisopropyladenosine (R-PIA). The C2-ethoxyadenosine analogs were 30-140-fold less potent than NECA and the C2-amino-substituted analogs were 250 to 1000-fold less potent than NECA at inducing relaxations of the guinea pig aorta. All of the analogs were also less potent than the A1-selective agonist R-PIA. However, only responses to NECA were competitively antagonized by the non-selective adenosine receptor antagonist 8-phenyltheophylline (8-PT), pKB = 6.83. Thus, the C2-substituted analogs produce relaxations of the guinea pig aorta through a combination of actions at A2-adenosine receptors and at xanthine resistant sites. The lack of potency of these analogs at activating the xanthine sensitive A2-receptors in the guinea pig aorta suggests that these adenosine receptors may be of the A2B-subtype.

IT 53296-10-9, CV-1808


RL: BIOL (Biological study)

(aorta relaxation by, purinergic receptor subtype in)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 34 THERE ARE 34 CAPLUS RECORDS THAT CITE THIS RECORD (34 CITINGS)

L6 ANSWER 110 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1992:440341 CAPLUS

DN 117:40341

OREF 117:6967a,6970a

TI Effect of adenosine analogs on the expression of opiate withdrawal in rats

AU Dionyssopoulos, Tim; Hope, Wendy; Coupar, Ian M.

CS Sch. Pharmacol., Victorian Coll. Pharm., Parkville, 3052, Australia

SO Pharmacology, Biochemistry and Behavior (1992), 42(2), 201-6

CODEN: PBBHAU; ISSN: 0091-3057

DT Journal

LA English


AB The adenosine A1 receptor agonist N6-[(R)-1-methyl-2-phenylethyl]adenosine (R-PIA), the A2 agonist 2-(phenylamino)adenosine (CV 1808), the nonselective A1, A2 agonist adenosine-5'-ethylcarboxamide (NECA), and the

McIntosh

$\alpha 2$ -adrenoceptor agonist clonidine were screened (each at 30, 100, and 300  $\mu\text{g/kg}$ , s.c.) for their ability to alter naloxone-precipitated withdrawal signs in morphine-dependent rats. The results indicate that there is convergent dependence involving opioid and adenosine A1 receptors on those effects expressed by withdrawal diarrhea, paw-shakes, teeth-chattering, body-shakes, and jumping. Further, dependence expressed by body-shakes involves convergence involving A1 receptors, as well as  $\alpha 2$ -adrenoceptors; while A1 receptors are involved in dependence expressed by jumping, stimulation of  $\alpha 2$ -adrenoceptors augments this sign. Adenosine analogs may be of clin. value for detoxification of opiate addicts.

IT 53296-10-9, CV 1808  
 RL: BIOL (Biological study)  
 (opiate withdrawal behaviors response to)  
 RN 53296-10-9 CAPLUS  
 CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.




OSC.G 16 THERE ARE 16 CAPLUS RECORDS THAT CITE THIS RECORD (16 CITINGS)

L6 ANSWER 111 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
 AN 1992:440225 CAPLUS  
 DN 117:40225  
 OREF 117:6935a,6938a  
 TI Characterization of human striatal A2 adenosine receptors using radioligand binding and photoaffinity labeling  
 AU Ji, Xiao Duo; Stiles, Gary L.; Van Galen, Philip J. M.; Jacobson, Kenneth A.  
 CS Lab. Bioorg. Chem., Natl. Inst. Diabet. Digest. Dis. Kidney Dis., Bethesda, MD, 20892, USA  
 SO Journal of Receptor Research (1992), 12(2), 149-69  
 CODEN: JRERDM; ISSN: 0197-5110  
 DT Journal  
 LA English  
 AB The adenosine agonist [3H]CGS21680 (2-[4-[[2-carboxethyl]phenyl]ethylamino]-5'-N-ethylcarboxamidoadenosine) bound to A2 receptors in human striatal membranes with a  $k_d$  of 17.8 nM and a  $B_{max}$  of 313 fmol/mg protein. The addition of 100  $\mu\text{M}$  GTP diminished both the affinity of agonist radioligand for A2 adenosine binding sites and the total binding, resulting in  $k_d$  and  $B_{max}$  values of 28.6 nM and 185 fmol/mg of protein. Adenosine ligands competed for [3H]CGS21680 with the expected potency order. The adenosine antagonist [3H]XAC (8-[4-[[[(2-aminoethyl)-amino]carbonyl]methyl]oxy]phenyl]-1,3-dipropylxanthine), although A1-selective in the rat, binds to human striatal A2 receptors with high affinity. 25 NM CPX (8-cyclopentyl-1,3-dipropylxanthine), an A1-selective antagonist, was added to the incubation medium and effectively eliminated 91% of [3H]XAC (1 nM) binding to human A1 receptors, yet preserved 90% of binding to A2 receptors. [3H]XAC exhibited saturable, specific binding (50% of total) to A2 sites with a  $k_d$  of 2.98 nM and a  $B_{max}$  of 0.71 pmol/mg protein (25°, non-specific binding defined with 100  $\mu\text{M}$  NECA). The potency order for antagonists against 1 nM [3H]XAC was CGS15943A > XAC = PD115,199 > PAPA-XAC > CPX > HTQZ = XCC = CP-66,713 > theophylline = caffeine, indicative of an A2-type binding site. A2a-receptors were found to be present in the human cortex, albeit at a much lower d. than in the striatum. Photoaffinity labeling using 125I-PAPA-APEC revealed a mol. weight of 45 K, but proteolytic cleavage was observed, resulting in fragments of MW 43 K and 37 K. In the absence of proteolytic inhibitors the 37 K fragment, which still bound 125I-PAPA-APEC, was predominant.  
 IT 53296-10-9, CV1808

10/598,520

RL: BIOL (Biological study)  
(CGS21680 binding to human striatal adenosine A2 receptors response to)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.




OSC.G 14 THERE ARE 14 CAPLUS RECORDS THAT CITE THIS RECORD (14 CITINGS)

L6 ANSWER 112 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1992:121533 CAPLUS  
DN 116:121533  
OREF 116:20337a,20340a  
TI Adenosine receptor-induced cAMP changes in D384 astrocytoma cells and the effect of bradykinin thereon  
AU Altiok, Nedret; Balmforth, A. J.; Fredholm, B. B.  
CS Dep. Pharmacol., Karolinska Inst., Stockholm, S-104 01, Swed.  
SO Acta Physiologica Scandinavica (1992), 144(1), 55-63  
CODEN: APSCAX; ISSN: 0001-6772  
DT Journal  
LA English  
AB In human D384 astrocytoma cells, cAMP accumulation can be conveniently studied after labeling of the ATP pool (15 fmol/cell) with [3H]adenine. In this study, adenosine had a biphasic effect on cAMP accumulation, which was scarcely altered by blocking adenosine uptake and metabolism. Low concns. of adenosine led to an inhibition of cAMP accumulation, and higher concns. led to stimulation. No effect of adenosine on cAMP was observed unless phosphodiesterase was inhibited by rolipram. The A1 receptor antagonist dipyridamole-1,3-dipropyl-8-cyclopentyl xanthine attenuated the inhibitory phase of adenosine response, and enhanced the cAMP accumulation induced by adenosine analogs. The cAMP accumulation was stimulated by NECA > ADO > CGS 21680 > CV 1808 > N6-cyclopentyladenosine ≥ N6-cyclohexyladenosine, indicating mediation by A2 receptors. The stimulatory effect of NECA was much more effectively blocked by the combined A1 and A2 receptor antagonist CGS 15943 (KB 4 nmol/L) than by the A1 antagonist DPCPX (KB 110 nmol/L). Treatment of the cells with pertussis toxin (0.2 µg/mL for 2.5 h) potentiated the cAMP response to adenosine analogs. The cAMP response to NECA was enhanced by the protein kinase C activator phorbol dibutyrate even after pertussis toxin treatment. By contrast, nanomolar concns. of bradykinin, which increases Ca2+-levels and protein kinase C activity in D384 cells, reduced NECA-induced cAMP accumulation in control and pertussis toxin-treated cells. Thus, D384 cells possess both A1 and A2 adenosine receptors influencing cAMP in opposite directions. A2 receptor-mediated cAMP accumulation can be stimulated by activating protein kinase C and inhibited by raising Ca2+. Neither the effects of protein kinase C activation nor those of bradykinin required pertussis toxin-sensitive G-proteins.  
IT 53296-10-9, CV 1808  
RL: BIOL (Biological study)  
(cAMP accumulation response to, in astrocytoma cell, mechanism for)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.


10/598,520



OSC.G 14 THERE ARE 14 CAPLUS RECORDS THAT CITE THIS RECORD (14 CITINGS)

L6 ANSWER 113 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1992:76245 CAPLUS  
DN 116:76245  
OREF 116:12755a,12758a  
TI Receptor binding at two different temperatures to discriminate agonist and antagonist behavior of adenosine A1 receptor ligands in rat brain  
AU Borea, Pier Andrea; Varani, Katia; Malaguti, Valeria; Gilli, Gastone  
CS Ist. Farmacol., Univ. Ferrara, Ferrara, 44100, Italy  
SO Journal of Pharmacy and Pharmacology (1991), 43(12), 866-8  
CODEN: JPPMAB; ISSN: 0022-3573  
DT Journal  
LA English  
AB The inhibitory binding consts.  $K_i$ , at the adenosine A1 receptor in rat brain have been measured at 0 and 25° for 25 typical ligands. The  $K_i$  ratios at the 2 temps. are greater and smaller than unity for adenosine agonists and xanthine antagonists, resp. These results suggest that 2-temperature measurements of in-vitro  $K_i$  consts. represent a simple method of discriminating between in-vivo agonistic and antagonistic behavior of A1 adenosine receptor ligands.  
IT 53296-10-9, 2-Phenylaminoadenosine  
RL: BIOL (Biological study)  
(receptor binding of, in brain, temperature effect on)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)


Absolute stereochemistry.



OSC.G 6 THERE ARE 6 CAPLUS RECORDS THAT CITE THIS RECORD (6 CITINGS)

L6 ANSWER 114 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1992:59857 CAPLUS  
DN 116:59857  
OREF 116:10376h,10377a  
TI Nucleosides and nucleotides. 103. 2-Alkynyladenosines: a novel class of selective adenosine A2 receptor agonists with potent antihypertensive effects  
AU Matsuda, Akira; Shinozaki, Misao; Yamaguchi, Toyofumi; Homma, Hiroshi; Nomoto, Rie; Miyasaka, Tadashi; Watanabe, Yohko; Abiru, Toichi  
CS Fac. Pharm. Sci., Hokkaido Univ., Sapporo, 060, Japan  
SO Journal of Medicinal Chemistry (1992), 35(2), 241-52  
CODEN: JMCMAR; ISSN: 0022-2623  
DT Journal  
LA English  
OS CASREACT 116:59857  
GI

McIntosh




AB The synthesis and receptor-binding activities at A1 and A2 adenosine receptors for a series of 2-alkynyladenosines, are described. The Pd-catalyzed cross-coupling reaction of 2-iodoadenosine (I; R = iodo) with various terminal alkynes in the presence of bis(triphenylphosphine)palladium dichloride and CuI in DMF containing NET3 gives 2-alkynyladenosines I [R = C.tplbond.CR2, R2 = Et, Pr, Bu, pentyl, hexyl, heptyl, octyl, decyl, dodecyl, tetradecyl, hexadecyl, CH2OH, CH2CH2OH, CH2OMe, CH2O(CH2)3Me]. An economical synthetic method for the preparation of 9-(2,3,5-tri-O-acetyl-1-β-D-ribofuranosyl)-6-chloro-2-iodopurine (II; R2 = iodo), which is a precursor of I (R = iodo) is also included. Several transformation reactions of 2-(1-octyn-1-yl)adenosine I [R = C.tplbond.C (CH2-Me)] and 2-(1-ethyn-1-yl)adenosine I (R = C.tplbond.CH) and a similar cross-coupling reaction of 6-chloropurine derivative II (R2 = H) and 8-bromoadenosine III with 1-octyne are also reported. Many of these 2-alkynyladenosines tested for A1 and A2 adenosine receptor binding activities in rat brain are selective for the A2 adenosine receptor. Among them, 2-(1-hexyn-1-yl)adenosine has the highest affinity for both A1 and A2 receptors with Ki values of 126.5 and 2.8 nM, resp. The structure-activity relationship of this series of compds. including 6- or 8-alkynylpurine nucleosides and 2-alkyl- and 2-alkenyladenosines is discussed in terms of potency at both receptor subtypes. Addnl., how hypotensive activity and heart rate decrease brought on by I (R = C.tplbond.CR3) and some other compds. with spontaneously hypertensive rats are proportional to the order of the potency to both A1 and A2 binding affinities, are described. Thus, 2-alkynyladenosines are interesting and promising as antihypertensive agents that should be considered for further detailed preclin. evaluation.

IT 137896-07-2P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (preparation of, as adenosine receptor agonist)  
 RN 137896-07-2 CAPLUS  
 CN Adenosine, 2-octyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

10/598,520



OSC.G 98 THERE ARE 98 CAPLUS RECORDS THAT CITE THIS RECORD (98 CITINGS)

L6 ANSWER 115 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1992:34585 CAPLUS  
DN 116:34585  
OREF 116:5737a,5740a  
TI Methods for treatment of alcohol intoxication and dependence  
IN Diamond, Ivan F.; Gordon, Adrienne S.  
PA USA  
SO Can. Pat. Appl., 24 pp.  
CODEN: CPXXEB  
DT Patent  
LA English  
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	CA 2029581	A1	19910510	CA 1990-2029581	19901108
	US 5069895	A	19911203	US 1989-434066	19891109
	EP 431758	A2	19910612	EP 1990-312252	19901108
	EP 431758	A3	19920115		

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE  
PRAI US 1989-434066 A 19891109

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB Alc.-related disorders are treated by the administration of adenosine antagonists and adenosine agonists to a host. Adenosine antagonists are used to inhibit both acute intoxication and chronic dependence by administering prior to alc. consumption. The symptoms associated with alc. withdrawal syndrome may be treated by administering adenosine agonists which reduce the physiol. dependence on alc. during the withdrawal period. Acute exposure to EtOH increased the concentration of extracellular adenosine which then activated adenosine A2 receptors to increase intracellular cAMP levels. Accumulation of extracellular adenosine was required for the development of chronic EtOH-induced heterologous desensitization of receptor-stimulated cAMP production. Extracellular adenosine accumulation was greater in lymphocytes of alcoholics than in lymphocytes of nonalcoholics. After chronic exposure to 100 mM EtOH for 24 h, rechallenge with EtOH did not increase extracellular adenosine in lymphocytes from nonalcoholics whereas it caused a 73% increase in lymphocytes from alcoholics.

IT 53296-10-9


RI: BIOL (Biological study)

(as adenosine agonist, for ethanol withdrawal syndrome treatment)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (3 CITINGS)


L6 ANSWER 116 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

McIntosh

10/598,520

AN 1992:34525 CAPLUS  
DN 116:34525  
OREF 116:5729a,5732a  
TI Comparative pharmacology of the nitrobenzylthioguanosine-sensitive and -resistant nucleoside transport mechanisms of Ehrlich ascites tumor cells  
AU Hammond, James R.  
CS Dep. Pharmacol. Toxicol., Univ. West. Ontario, London, ON, N6A 5C1, Can.  
SO Journal of Pharmacology and Experimental Therapeutics (1991), 259(2), 799-807  
CODEN: JPETAB; ISSN: 0022-3565  
DT Journal  
LA English  
AB A variety of nucleoside transport inhibitors and substrates were compared for their capacities to inhibit the zero-trans influx of [3H]uridine in Ehrlich ascites tumor cells. ATP-depleted cells accumulated [3H]uridine primarily by facilitated diffusion ( $V_{\max} = 16 \text{ pmol/s/}\mu\text{L cell water}$ ) via both nitrobenzylthioguanosine (NBTGR)-sensitive ( $IC_{50} = 0.53 \text{ nM}$ ,  $100 \mu\text{M}$  [3H]uridine) and NBTGR-resistant ( $IC_{50} = 71 \mu\text{M}$ ,  $100 \mu\text{M}$  [3H]uridine) mechanisms with uridine  $K_M$  ests. of 99 and  $284 \mu\text{M}$ , resp. Dilazep also distinguished between the transporter subtypes with  $IC_{50}$  values of 14 nM and  $1.8 \mu\text{M}$ , resp., for inhibiting  $100 \mu\text{M}$  [3H]uridine influx. Incubation of cells with 50 mM NBTGR allowed the selective study of inhibitor effects on NBTGR-resistant [3H]uridine influx. Dipyridamole, cyclopentyladenosine, 2-phenylaminoadenosine, etoposide, teniposide, diazepam, chlordiazepoxide, triazolam and the lidoflazine derivative R75231, were less potent as inhibitors of NBTGR-resistant influx, when compared with their capacities to inhibit the total mediated influx [3H]uridine. In contrast, 2-fluoroadenosine, 2-chloroadenosine, 5'-N-ethylcarboxamidoadenosine and solufazine were relatively more effective as inhibitors of the NBTGR-resistant component. Mioflazine, a compound related to both solufazine and R75231, did not distinguish between transporter subtypes. The NBTGR-resistant transporter also had a distinctive substrate specificity; guanosine, 2'-deoxyguanosine, cytidine and 2'-deoxycytidine were less effective as inhibitors of NBTGR-resistant [3H]uridine influx. These results show that the NBTGR-sensitive and -resistant nucleoside transporters of Ehrlich cells have distinctive pharmacol. profiles that extend beyond the well-characterized differential affinities for dilazep and S6-thiopurine derivs., and that relatively minor modifications in mol. structure have a significant impact on transporter selectivity. Further structure activity studies are clearly warranted, and may lead to the development of more selective inhibitors for the NBTGR-resistant nucleoside transport system.  
IT 53296-10-9, CV-1808  
RL: BIOL (Biological study)  
(nucleoside transport system response to, in Ehrlich ascites tumor cells, nitrobenzylthioguanosine sensitivity in)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 19 THERE ARE 19 CAPLUS RECORDS THAT CITE THIS RECORD (20 CITINGS)

L6 ANSWER 117 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1992:6922 CAPLUS  
DN 116:6922  
OREF 116:1370h,1371a  
TI Preparation of 2-aralkoxy- and 2-alkoxyadenosine derivatives as coronary vasodilators and antihypertensive agents  
IN Olsson, Ray A.; Thompson, Robert D.  
PA Whitby Research, Inc., USA

McIntosh



10/598,520

SO PCT Int. Appl., 31 pp.

CODEN: PIXXD2

DT Patent

LA English


FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9113082	A1	19910905	WO 1991-US1023	19910214
	W: AU, BB, BG, BR, CA, FI, HU, JP, KP, KR, LK, MC, MG, MW, NO, RO, SD, SU				
	RW: AT, BE, BF, BJ, CF, CG, CH, CM, DE, DK, ES, FR, GA, GB, GR, IT, LU, ML, MR, NL, SE, SN, TD, TG				
	US 5140015	A	19920818	US 1990-482282	19900220
	AU 9173255	A	19910918	AU 1991-73255	19910214
	AU 645784	B2	19940127		
	EP 515514	A1	19921202	EP 1991-904813	19910214
	EP 515514	B1	20000830		
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE				
	JP 05506436	T	19930922	JP 1991-505571	19910214
	JP 3160611	B2	20010425		
	AT 195946	T	20000915	AT 1991-904813	19910214
	ES 2150903	T3	20001216	ES 1991-904813	19910214
	CA 2074853	A1	19940130	CA 1992-2074853	19920729
	CA 2074853	C	20050607		
	US 36494	E	20000111	US 1993-98180	19930726
PRAI	US 1990-482282	A	19900220		
	WO 1991-US1023	A	19910214		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OS MARPAT 116:6922

GI



AB Title compds. I [R<sub>1</sub> = (substituted) C<sub>1</sub>-6 hydrocarbyl, cyclic hydrocarbyl, (substituted) Ph, (substituted) thienyl, (substituted) naphthyl, (substituted) indolyl, etc.; R<sub>2</sub> = (hydroxy) C<sub>1</sub>-4 hydrocarbyl; X = 2H or O; B = O, N; with provisos] were prepared as adenosine A<sub>2</sub> receptor agonists useful as coronary vasodilators and antihypertensives. Thus, n-BuLi in hexanes was added to a solution of 4-FlC<sub>6</sub>H<sub>4</sub>(CH<sub>2</sub>)<sub>2</sub>OH in THF at 10°. The solution was stirred 15 min at room temperature, then 2-chloro-2',3'-O-(ethoxymethylidene)adenosine was added and the mixture was refluxed 4 days. The resulting product was deprotected by HOAc hydrolysis to give 2-[2-(4-fluorophenyl)ethoxy]adenosine (II). II at 0.9 nM gave a half-maximal increase in coronary blood flow in guinea pigs vs. 49.7 nM for adenosine.

IT 131865-78-6P 131865-81-1P 131865-82-2P  
137817-83-5P 137817-84-6P

RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of, as adenosine A<sub>2</sub> receptor agonists)


RN 131865-78-6 CAPLUS

CN Adenosine, 2-(phenylmethoxy)- (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 131865-81-1 CAPLUS  
CN Adenosine, 2-(4-phenylbutoxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 131865-82-2 CAPLUS  
CN Adenosine, 2-[(5-phenylpentyl)oxy]- (CA INDEX NAME)

Absolute stereochemistry.



RN 137817-83-5 CAPLUS  
CN Adenosine, 2-[(2R)-2-phenylbutoxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 137817-84-6 CAPLUS  
CN Adenosine, 2-[(2S)-2-phenylbutoxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

McIntosh


10/598,520



OSC.G 6 THERE ARE 6 CAPLUS RECORDS THAT CITE THIS RECORD (7 CITINGS)  
RE.CNT 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 118 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1992:6886 CAPLUS  
DN 116:6886  
OREF 116:1363a,1366a  
TI An efficient synthesis of 2-(phenylamino)adenosine [CV-1808], an adenosine  
A2 receptor selective agonist  
AU Trivedi, Bharat K.  
CS Dep. Chem., Warner-Lambert/Parke-Davis Pharm. Res. Div., Ann Arbor, MI,  
48105, USA  
SO Nucleic Acid Chem. (1991), 264-8. Editor(s): Townsend, Leroy B.; Tipson,  
R. Stuart. Publisher: Wiley, New York, N. Y.  
CODEN: 39GCA6  
DT Conference  
LA English  
OS CASREACT 116:6886  
AB A five-step process for synthesizing title CV-1808 from com. available  
guanosine is reported. A key step is the conversion of guanosine  
2',3',5'-triacetate into 2-bromoinosine 2',3',5'-triacetate by using CHBr3  
and an alkyl nitrite.  
IT 53296-10-9P  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



L6 ANSWER 119 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1991:680484 CAPLUS  
DN 115:280484  
OREF 115:47683a,47686a  
TI Preparation of 8-hydroxy-2',3',-dideoxyadenosine as an antiviral  
IN Nair, Vasu; Buenger, Greg S.  
PA University of Iowa Research Foundation, USA  
SO U.S., 6 pp.  
CODEN: USXXAM  
DT Patent  
LA English  
FAN.CNT 1


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 5013829	A	19910507	US 1989-343334	19890426
PRAI	US 1989-343334		19890426		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

McIntosh

10/598,520

GI




AB Many intermediates [I; R1, R2 = cyano, H; COHN2, H; Et, H, H, OMe; etc.] for the title compound [I; R1 = H, R2 = OH] (II) stable against deamination and hydrolytic cleavage of the glycosidic bond, an antiviral especially useful for the treatment of AIDS (no data) were prepared E.g., a solution of 8-bromo-2'-deoxyadenosine in MeOH containing MeONa was refluxed for 20 to give 55% 2'-deoxy-8-methoxyadenosine, which was converted to 2',3'-dideoxy-8-methoxyadenosine via formation of 2'-deoxy-3'-O-(1-imidazolylthiocarbonyl)-5'-O-(tert-butyltrimethylsilyl)adenosine, deoxygenation, and desilylation (detailed procedures not given). The conversion into II is not illustrated.

IT 79936-11-1P  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of, as intermediate for stable antivirals)

RN 79936-11-1 CAPLUS

CN Adenosine, 2-cyano- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)

RE.CNT 10 THERE ARE 10 CITED REFERENCES AVAILABLE FOR THIS RECORD

ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 120 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1991:652834 CAPLUS

DN 115:252834

OREF 115:42921a,42924a

TI Cardiovascular actions of adenosines, but not adenosine receptors, differ in rat and guinea pig

AU Ueda, Masayuki; Thompson, Robert D.; Padgett, William L.; Secunda, Sherrie; Daly, John W.; Olsson, Ray A.

CS Dep. Intern. Med. Biochem. Mol. Biol., Univ. South Florida, Tampa, FL, 33612, USA

SO Life Sciences (1991), 49(18), 1351-8  
CODEN: LIFSAK; ISSN: 0024-3205

DT Journal

LA English


AB The structure-activity relationships of 16 analogs at the A1 and A2 adenosine receptors (A1AR, A2AR) of rat and guinea pig were compared. Radioligand binding studies revealed no marked differences in the affinities of each analog at the A2AR of brain cortex or the A2AR of brain striatum. Bioassay employing Langendorff heart preps. showed that the guinea pig is more sensitive than the rat to A1AR-mediated slowing of conduction through the atrioventricular node and, in some instances, to A2AR-mediated coronary vasodilation. This difference could reflect factors such as receptor d. or efficacy of coupling to effector systems.

McIntosh

10/598,520


IT 50257-82-4 50257-89-1 50257-95-9,  
2-Hexyloxyadenosine 53296-10-9 131865-78-6  
131933-17-0  
RL: BIOL (Biological study)  
(brain adenosine A1 and A2 receptors binding of and cardiovascular  
action of, in guinea pig and rat)  
RN 50257-82-4 CAPLUS  
CN Adenosine, 2-phenoxy- (CA INDEX NAME)

Absolute stereochemistry.




RN 50257-89-1 CAPLUS  
CN Adenosine, 2-(2-hydroxyethoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 50257-95-9 CAPLUS  
CN Adenosine, 2-(hexyloxy)- (CA INDEX NAME)

Absolute stereochemistry.



RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.




McIntosh

10/598,520


RN 131865-78-6 CAPLUS  
CN Adenosine, 2-(phenylmethoxy)- (CA INDEX NAME)

Absolute stereochemistry.



RN 131933-17-0 CAPLUS  
CN Adenosine, 2-(cyclohexylmethoxy)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



OSC.G 6 THERE ARE 6 CAPLUS RECORDS THAT CITE THIS RECORD (6 CITINGS)

L6 ANSWER 121 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1991:651461 CAPLUS  
DN 115:251461  
OREF 115:42649a,42652a  
TI Modulation of [3H]nitrobenzylthioinosine binding kinetics  
AU Hammond, James R.  
CS Dep. Pharmacol. Toxicol., Univ. West. Ontario, London, ON, N6A 5C1, Can.  
SO Nucleosides & Nucleotides (1991), 10(5), 1103-6  
CODEN: NUNUD5; ISSN: 0732-8311  
DT Journal  
LA English  
AB Inhibitors and substrates of the nucleoside transporter of Ehrlich cell membrane were tested for their effects on the kinetics of [3H]nitrobenzylthioinosine binding. Results are discussed in terms of a distinct site mediating the allosteric modulation of [3H]nitrobenzylthioinosine-binding affinity.  
IT 53296-10-9, CV-1808  
RL: ANST (Analytical study)  
(nitrobenzylthioinosine binding kinetics in mammalian cell membrane response to)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



McIntosh

10/598,520

OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

L6 ANSWER 122 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1991:624462 CAPLUS

DN 115:224462

OREF 115:38075a,38078a

TI Adenosine and ATP produce vasoconstriction in the feline pulmonary vascular bed by different mechanisms

AU Neely, Constance Fisher; Haile, Daniel M.; Cahill, Bruce E.; Kadowitz, Philip J.

CS Sch. Med., Univ. Pennsylvania, Philadelphia, PA, 19104, USA

SO Journal of Pharmacology and Experimental Therapeutics (1991), 258(3), 753-61

CODEN: JPETAB; ISSN: 0022-3565

DT Journal

LA English

AB Adenosine and ATP produce dose- and tone-dependent responses in the feline pulmonary vascular (PV) bed. The mechanisms mediating vasoconstrictor (VC) responses to adenosine and ATP in the intact-chest, spontaneously breathing cats under conditions of controlled blood flow and constant left atrial pressure were studied. The order of potency of adenosine receptor agonists to produce VC in the PV bed was the selective adenosine A1 receptor agonist R-phenylisopropyladenosine > the mixed A1, A2 receptor agonist adenosine > the selective adenosine A2 receptor agonist 2-phenylaminoadenosine. The dose-related increase in lobar arterial pressure in response to adenosine was blocked by the adenosine (P1) receptor antagonist BWA1433U, the cyclooxygenase inhibitor meclofenamate, and the TXA2 receptor antagonist SQ29548. The order of potency of ATP analogs to produce VC in the PV bed was  $\alpha,\beta$ -methylene ATP ( $\alpha,\beta$ -meATP) »  $\beta,\tau$ -methylene ATP > ATP. BWA1433U inhibited VC responses to ATP without affecting the responses to its degradation-resistant analogs  $\beta,\tau$ -methylene ATP and  $\alpha,\beta$ -meATP. In the presence of BWA1433U and a continuous intralobar infusion of the selective 5'-nucleotidase inhibitor  $\alpha,\beta$ -methyleneadenosine-5'-diphosphate, ATP VC responses were enhanced compared to those after BWA1433U.  $\alpha,\beta$ -Methyleneadenosine-5'-diphosphate had no effect on the VC response to U44069 after BWA1433U. Meclofenamate inhibited the vasoconstrictor responses to ATP but not to  $\alpha,\beta$ -meATP. Repeated injections of  $\alpha,\beta$ -meATP produced selective inhibition of the VC responses to ATP without affecting VC responses to adenosine, norepinephrine, or angiotensin II. By using this technique to desensitize P2x receptors, subsequent injections of ATP blocked the VC responses to adenosine. Adenosine may produce VC in the feline PV bed by acting on an adenosine A1-'like' receptor coupled to a phospholipase which causes the release of TXA2. ATP may produce VC following its metabolism to adenosine but also by acting on the specific ATP receptor, P2x not coupled to a phospholipase.

IT 53296-10-9, 2-Phenylaminoadenosine


RL: BIOL (Biological study)

(pulmonary vasoconstriction from, receptor mechanism of)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 16 THERE ARE 16 CAPLUS RECORDS THAT CITE THIS RECORD (16 CITINGS)

L6 ANSWER 123 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1991:536628 CAPLUS


DN 115:136628

McIntosh

10/598,520

OREF 115:23451a,23454a  
TI An efficient synthesis of 2-(phenylamino)adenosine [CV-1808]: an adenosine  
A2 receptor selective agonist.  
AU Trivedi, Bharat K.  
CS Dep. Chem., Warner-Lambert/Parke-Davis Res. Div., Ann Arbor, MI, 48105,  
USA  
SO Nucleic Acid Chem. (1991), Volume 4, 264-8. Editor(s): Townsend, Leroy  
B.; Tipson, R. Stuart. Publisher: Wiley, New York, N. Y.  
CODEN: 39GCA6  
DT Conference  
LA English  
AB A five-step process for synthesizing CV-1808 from com. available guanosine  
is reported. A key step is the conversion of guanosine  
2',3',5'-triacetate by using CHBr<sub>3</sub> and an alkyl nitrite.  
IT 53296-10-9P, 2-Phenylaminoadenosine  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

L6 ANSWER 124 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1991:487624 CAPLUS  
DN 115:87624  
OREF 115:14955a,14958a  
TI Kinetic analysis of ligand binding to the Ehrlich cell nucleoside  
transporter: pharmacological characterization of allosteric interactions  
with the [3H]nitrobenzylthioinosine binding site  
AU Hammond, James R.  
CS Dep. Pharmacol. Toxicol., Univ. West. Ontario, London, ON, N6A 5C1, Can.  
SO Molecular Pharmacology (1991), 39(6), 771-9  
CODEN: MOPMA3; ISSN: 0026-895X  
DT Journal  
LA English  
AB Kinetic anal. of the binding of [3H]nitrobenzylthioinosine ([3H]NBMPR) to  
Ehrlich ascites tumor cell plasma membranes was conducted in the presence  
and absence of a variety of nucleoside transport inhibitors and  
substrates. The association of [3H]NBMPR with Ehrlich cell membranes occurred  
in two distinct phases, possibly reflecting functional conformation  
changes in the [3H]NBMPR binding site/nucleoside transporter complex.  
Inhibitors of the equilibrium binding of [3H]NBMPR, tested at submaximal  
inhibitory concns., generally decreased the rate of association of [3H]NBMPR,  
but the magnitude of this effect varied significantly with the agent  
tested. Adenosine and diazepam had relatively minor effects on the  
association rate, whereas dipyrindamole and mioflazine slowed the rate  
dramatically. Inhibitors of nucleoside transport also decreased the rate  
of dissociation of [3H]NBMPR, with an order of potency different from their  
relative potencies as inhibitors of the equilibrium binding of [3H]NBMPR.  
Dilazep, dipyrindamole, and mioflazine were effective inhibitors of both  
[3H]NBMPR dissociation and equilibrium binding. The lidoflazine analog R75231, on  
the other hand, had no effect on the rate of dissociation of [3H]NBMPR at  
concns. below 300 µM, even though it was one of the most potent  
inhibitors of [3H]NBMPR binding tested (K<sub>i</sub> < 100 nM). In contrast, a  
series of natural substrates for the nucleoside transport system enhanced  
the rate of dissociation of [3H]NBMPR with an order of effectiveness that  
paralleled their relative affinities for the permeant site of the  
transporter. The most effective enhancers of [3H]NBMPR dissociation, however,  
were the benzodiazepines diazepam, chlordiazepoxide, and triazolam.

McIntosh



Comparable effects of adenosine and dipyridamole on [3H]NBMPR dissociation rate were obtained upon solubilization of the membranes with octylglucoside, suggesting that this phenomenon was not due to changes in membrane fluidity. These results are compatible with the existence of specific ligand recognition sites on the nucleoside transport complex of Ehrlich cells that are pharmacol. distinct from, but allosterically linked to, the high affinity binding sites for [3H]NBMPR. The marked effects on [3H]NBMPR binding kinetics that result from ligand interactions with these sites must be considered in the design and anal. of all studies involving the use of [3H]NBMPR as a high affinity probe for the nucleoside transport system.

IT 53296-10-9, CV-1808


RL: BIOL (Biological study)

(nucleoside transport system binding of nitrobenzylthioinosine response to)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 14 THERE ARE 14 CAPLUS RECORDS THAT CITE THIS RECORD (15 CITINGS)

L6 ANSWER 125 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1991:442471 CAPLUS

DN 115:42471

OREF 115:7225a,7228a

TI The antinociceptive effect of intrathecally administered adenosine analogs in mice correlates with the affinity for the A1-adenosine receptor

AU Karlsten, Rolf; Post, Claes; Hide, Izumi; Daly, John W.

CS Dep. Anesthesiol., Univ. Hosp., Uppsala, S-751 85, Swed.

SO Neuroscience Letters (1991), 121(1-2), 267-70

CODEN: NELED5; ISSN: 0304-3940

DT Journal

LA English

AB The antinociceptive effects after intrathecal injection of each of 6 N6-substituted adenosine analogs and of 2-phenylaminoadenosine were compared with the affinity for the A1- and A2-adenosine receptors. Adenosine analogs, substituted in the N6-position, had stereoselective structure-dependent antinociceptive effects in the tail flick and hot plate assays after intrathecal injection in mice. The antinociceptive activity for N6-R- and S-phenylisopropyladenosine, N6-R- and S-1-phenylethyladenosine, N6-1,1-dimethyl-2-phenylethyladenosine, and N6-cyclooctyladenosine correlated with the affinity for central A1-adenosine receptors. An adenosine analog, 2-phenylaminoadenosine, selective for A2-adenosine receptors was inactive in the 2 tests. These results strongly suggest that spinal A1-adenosine receptors are responsible for the antinociceptive effects of adenosine and its analogs after intrathecal injection.

IT 53296-10-9


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)

(analgesic action of, after intrathecal administration, affinity for A1 adenosine receptors in relation to)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 25 THERE ARE 25 CAPLUS RECORDS THAT CITE THIS RECORD (25 CITINGS)

L6 ANSWER 126 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1991:241169 CAPLUS

DN 114:241169

OREF 114:40541a,40544a

TI Relaxant effects of adenosine analogs on guinea pig trachea in vitro:  
xanthine-sensitive and xanthine-insensitive mechanisms

AU Brackett, L. E.; Daly, J. W.

CS Lab. Bioorg. Chem., Natl. Inst. Diabetes, Dig. Kidney Dis., Bethesda, MD,  
20892, USA

SO Journal of Pharmacology and Experimental Therapeutics (1991), 257(1),  
205-13

CODEN: JPETAB; ISSN: 0022-3565

DT Journal

LA English

AB Adenosine analogs were tested for their ability to relax carbachol-contracted trachea in vitro. The rank order of potency was: 5'-N-ethylcarboxamidoadenosine (NECA) > 2-chloroadenosine (2-ClADO) > 5'-chloroadenosine = N6-R-1-phenyl-2-propyladenosine (R-PIA) > N6-cyclohexyladenosine > 2-phenylaminoadenosine (CV1808) > 5'-methylthiadenosine (MTA). The rank order of potency for NECA, 2-ClADO and R-PIA is characteristic of an A2 subtype of adenosine receptor. 8-Para-sulfophenyltheophylline (8-p-ST) and 8-cyclopentyl-1,3-dipropylxanthine (DPCPX), were used to antagonize tracheal relaxation elicited by adenosine analogs. 8-p-ST antagonized the 2-ClADO, N6-cyclohexyladenosine, R-PIA and 5'-chloroadenosine responses, but had little or no effect on the CV1808 and MTA responses. 8-p-ST antagonized responses to NECA at concns. of NECA up to .apprx.30  $\mu$ M, but had no effect on responses to higher concns. of NECA. The differences in antagonist potency of 8-p-ST and the clear biphasic response of NECA are indicative of at least 2 mechanisms of adenosine analog action leading to tracheal relaxation. One mechanism is mediated through a xanthine-sensitive site, at which NECA acted in a potent manner, whereas the other mechanism or mechanisms are insensitive to blockade by xanthines and account for the effects of action of MTA and CV1808, as well for NECA at high concns. The low potency of the A1-selective antagonist DPCPX indicates that the xanthine-sensitive site is an A2 type receptor. MTA is known to be an antagonist at A2-adenosine receptors that stimulate adenylate cyclase activity, yet MTA did not antagonize the NECA-induced relaxation of trachea. Thus, the A2-type adenosine receptors in smooth muscle appear different from the A2-adenosine receptors that are linked to adenylate cyclase in other tissues.

IT 53296-10-9, CV1808

RL: BIOL (Biological study)


(trachea relaxation induction by, xanthine-sensitive and -insensitive mechanisms for)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.

10/598,520



OSC.G 18 THERE ARE 18 CAPLUS RECORDS THAT CITE THIS RECORD (18 CITINGS)

L6 ANSWER 127 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1991:240324 CAPLUS

DN 114:240324

OREF 114:40361a,40364a

TI The antihypertensive effect of 2-alkynyladenosines and their selective affinity for adenosine A2 receptors

AU Abiru, Toichi; Yamaguchi, Toyofumi; Watanabe, Yohko; Kogi, Kentaro; Aihara, Kazuyuki; Matsuda, Akira

CS Res. Dev. Div., Yamasa Shoyu Co., Ltd., Choshi, 288, Japan

SO European Journal of Pharmacology (1991), 196(1), 69-76

CODEN: EJPHAZ; ISSN: 0014-2999

DT Journal

LA English

AB The affinity for adenosine receptors and the antihypertensive effects of nine adenosine derivs., especially the alkynyl compds. 2-hexynyladenosine (2-H-Ado) and 2-octynyladenosine (2-O-Ado), was studied. The order of decreasing affinity of the agonists tested for rat brain A1 receptors was N6-cyclopentyladenosine (CPA) > N6-cyclohexyladenosine (CHA) > N6-R-phenylisopropyladenosine (R-PIA) > 2-chloroadenosine (CADO) = 5'-N-ethylcarboxamidoadenosine (NECA) > N6-S-phenylisopropyladenosine (S-PIA) > 2-H-Ado > 2-phenylaminoadenosine (CV-1808), and that for A2 receptors was 2-H-Ado > 2-O-Ado = NECA > CADO > CV-1808 > R-PIA > CPA > CHA > S-PIA. The Ki values of 2-H-Ado and 2-O-Ado for inhibiting [3H] NECA binding to A2 receptors were 4.1 and 12.1 nM, resp., and those for [3H]CHA binding to A1 receptors were 146 and 211 nM, resp.: the affinity of 2-H-Ado and 2-O-Ado for A2 receptors was about 36- and 17-fold, resp., higher than their affinity for A1 receptors. Injection of 2-H-Ado and 2-O-Ado (0.03-100 µg/kg) decreased the blood pressure of anesthetized, spontaneously hypertensive rats (SHR). A slight decrease in heart rate was observed after i.v. injection of 100 µg 2-H-Ado and 2-O-Ado/kg. A potent and long-lasting antihypertensive effect was also observed after oral administration of 2-H-Ado and 2-O-Ado to conscious SHR. These results show that 2-H-Ado and 2-O-Ado are potent and selective adenosine A2 receptor agonists; these agents lower blood pressure after oral administration but are less effective in decreasing heart rate.

IT 53296-10-9, 2-Phenylaminoadenosine


RL: BIOL (Biological study)

(antihypertensive activity and purinergic A2 receptor affinity of)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 19 THERE ARE 19 CAPLUS RECORDS THAT CITE THIS RECORD (20 CITINGS)

L6 ANSWER 128 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN


AN 1991:240169 CAPLUS

McIntosh

10/598,520


DN 114:240169  
OREF 114:40325a,40328a  
TI Adenosine and 2-phenylaminoadenosine (CV-1808) inhibit human neutrophil bactericidal function  
AU Hardart, G. E.; Sullivan, G. W.; Carper, H. T.; Mandell, G. L.  
CS Dep. Med., Univ. Virginia, Charlottesville, VA, 22908, USA  
SO Infection and Immunity (1991), 59(3), 885-9  
CODEN: INFIBR; ISSN: 0019-9567  
DT Journal  
LA English  
AB Adenosine is a natural autocoid and immunomodulator that serves an anti-inflammatory role. Stimulation of polymorphonuclear neutrophils (PMN) with soluble stimuli has been shown to inhibit the PMN oxidative burst. The authors examined the effects of adenosine and the adenosine analog 2-phenylaminoadenosine (CV-1808) on PMN bactericidal function. Adenosine (10 mM) and CV-1808 (10 to 100  $\mu$ M) inhibited PMN killing of Staphylococcus aureus. There were more surviving bacteria after 240 min of incubation of PMN with S. aureus and adenosine (10 mM) or CV-1808 (100  $\mu$ M) (254 and 739% of control, resp.) than there were in the control. In contrast, inosine (10 mM), the major degradation product of adenosine, did not affect killing. Adenosine and CV-1808 did not alter cell association of S. aureus, but S. aureus-activated PMN superoxide release was decreased by adenosine (10  $\mu$ M) and CV-1808 (10  $\mu$ M) to 67 and 32% that of the control, resp. Since adenosine inhibited PMN bactericidal function only at .apprx.10,000 times peak physiol. concns., endogenous adenosine levels would not be expected to adversely affect PMN bactericidal function. On the other hand, pharmacol. concns. of adenosine derivs. may decrease the oxidative burst and killing sufficiently to increase host susceptibility to infection.  
IT 53296-10-9, CV 1808  
RL: BIOL (Biological study)  
(polymorphonuclear neutrophils of humans bactericidal function inhibition by)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 10 THERE ARE 10 CAPLUS RECORDS THAT CITE THIS RECORD (10 CITINGS)

L6 ANSWER 129 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1991:185903 CAPLUS  
DN 114:185903  
OREF 114:31415a,31418a  
TI 2-Aralkoxyadenosines: potent and selective agonists at the coronary artery A2 adenosine receptor  
AU Ueeda, Masayuki; Thompson, Robert D.; Arroyo, Luis H.; Olsson, Ray A.  
CS Dep. Intern. Med., Univ. South Florida, Tampa, FL, 33612, USA  
SO Journal of Medicinal Chemistry (1991), 34(4), 1340-4  
CODEN: JMCMAR; ISSN: 0022-2623  
DT Journal  
LA English  
GI



AB A Langendorff guinea pig heart preparation served for the assay of agonist potency of 26 2-aralkoxyadenosines I (R = Ph, Ph(CH<sub>2</sub>)<sub>n</sub>, R<sub>1</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>CH<sub>2</sub>, R<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>, n = 2-5; R<sub>1</sub> = 2-, 3-, 4-F, 2-, 3-, 4-Cl, 2-, 3-, 4-MeO, 2-, 3-, 4-Me, R<sub>2</sub> = 2-, 3-thienyl, 3-indolyl, 1-, 2-naphthyl, 3,4-(MeO)<sub>2</sub>C<sub>6</sub>H<sub>3</sub>, 3,4,5-(MeO)<sub>3</sub>C<sub>6</sub>H<sub>2</sub>] at the A<sub>1</sub> and A<sub>2</sub> receptors of, resp., the atrioventricular node (conduction block) and coronary arteries (vasodilation). All of the analogs are weak agonists at the A<sub>1</sub> receptor, requiring concns. >9 μM to cause heart block. At the A<sub>2</sub> receptor 2-phenethoxyadenosine (I; R = PhCH<sub>2</sub>CH<sub>2</sub>) is the most potent of the 2-phenylalkyladenosines. The activity of ring-substituted (F, Cl, CH<sub>3</sub>, and OCH<sub>3</sub>) 2-phenethoxyadenosines increases ortho < meta < para. The EC<sub>50</sub>s of coronary vasodilation of 190 pM and an A<sub>1</sub>/A<sub>2</sub> selectivity ratio of 44000. Aryl groups such as thienyl, indolyl, or naphthyl also support A<sub>2</sub> agonist activity. Although the 2-oxoadenosine is 3 times more potent than 2-aminoadenosine, the activities of the Ph derivs. are markedly different; 2-phenoxyadenosine (I; R = Ph) is 23 times weaker than 2-(phenylamino)adenosine (CV-1808).


IT 50257-82-4P 131865-78-6P 131865-81-1P  
131865-82-2P

RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation and cardiac and coronary activity of)

RN 50257-82-4 CAPLUS

CN Adenosine, 2-phenoxy- (CA INDEX NAME)


Absolute stereochemistry.



RN 131865-78-6 CAPLUS

CN Adenosine, 2-(phenylmethoxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 131865-81-1 CAPLUS

CN Adenosine, 2-(4-phenylbutoxy)- (CA INDEX NAME)


10/598,520

Absolute stereochemistry.



RN 131865-82-2 CAPLUS  
CN Adenosine, 2-[(5-phenylpentyl)oxy]- (CA INDEX NAME)

Absolute stereochemistry.




OSC.G 22 THERE ARE 22 CAPLUS RECORDS THAT CITE THIS RECORD (22 CITINGS)

L6 ANSWER 130 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1991:185902 CAPLUS  
DN 114:185902  
OREF 114:31415a,31418a  
TI 2-Alkoxyadenosines: potent and selective agonists at the coronary artery  
A2 adenosine receptor  
AU Ueeda, Masayuki; Thompson, Robert D.; Arroyo, Luis H.; Olsson, Ray A.  
CS Dep. Intern. Med., Univ. South Florida, Tampa, FL, 33612, USA  
SO Journal of Medicinal Chemistry (1991), 34(4), 1334-9  
CODEN: JMCMAR; ISSN: 0022-2623  
DT Journal  
LA English  
OS CASREACT 114:185902  
AB A Langendorff guinea pig heart preparation served for the assay of agonist activity of a series of 24 2-alkoxyadenosines at the A1 and A2 adenosine receptors of, resp., the atrioventricular node (conduction block) and coronary arteries (vasodilation). Activities are low at the A1 receptor and do not show a clear relationship to the size or hydrophobicity of the C(2) substituent. All the analogs are more potent at the A2 receptor, activity varying directly with the size and hydrophobicity of the alkyl group. The most potent analog in this series, 2-(2-cyclohexylethoxy)adenosine, has an EC50 of 1 nM for coronary vasodilation and is 8700-fold selective for the A2 receptor.  
IT 50257-84-6P, 2-Butoxyadenosine 50257-85-7P  
50257-89-1P 50257-95-9P, 2-(Hexyloxy)adenosine  
131933-15-8P 131933-17-0P 131933-20-5P  
131933-26-1P 131933-27-2P 131973-26-7P  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation and adenosine receptor agonist activity of)  
RN 50257-84-6 CAPLUS  
CN Adenosine, 2-butoxy- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

10/598,520




RN 50257-85-7 CAPLUS  
CN Adenosine, 2-(pentyloxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 50257-89-1 CAPLUS  
CN Adenosine, 2-(2-hydroxyethoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 50257-95-9 CAPLUS  
CN Adenosine, 2-(hexyloxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 131933-15-8 CAPLUS  
CN Adenosine, 2-(3-methylbutoxy)- (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 131933-17-0 CAPLUS  
CN Adenosine, 2-(cyclohexylmethoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 131933-20-5 CAPLUS  
CN Adenosine, 2-(4-cyclohexylbutoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 131933-26-1 CAPLUS  
CN Adenosine, 2-(2-ethylbutoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 131933-27-2 CAPLUS  
CN Adenosine, 2-[(3-ethylpentyl)oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

McIntosh




10/598,520



RN 131973-26-7 CAPLUS  
CN Adenosine, 2-[(4-methylpentyl)oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 32 THERE ARE 32 CAPLUS RECORDS THAT CITE THIS RECORD (32 CITINGS)

L6 ANSWER 131 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1991:39747 CAPLUS

DN 114:39747

OREF 114:6883a,6886a

TI Characterization of adenosine A1 receptors in intact DDT1 MF-2 smooth muscle cells

AU Gerwins, P.; Nordstedt, C.; Fredholm, B. B.

CS Dep. Pharmacol., Karolinska Inst., Stockholm, S-104 01, Swed.

SO Molecular Pharmacology (1990), 38(5), 660-6

CODEN: MOPMA3; ISSN: 0026-895X

DT Journal

LA English


AB Adenosine receptors in the smooth muscle cell line DDT1 MF-2 were studied by radioligand binding using the A1 receptor-selective antagonist [3H]8-cyclopentyl-1,3-dipropylxanthine ([3H]DPCPX) as the ligand. Binding characteristics were similar in intact cells and in membranes (KD value of .apprx.1 nM). The maximum binding amounted to 183 fmol/106 intact cells or 344 fmol/mg of membranes. To characterize the receptor, competition expts. were performed by inhibiting [3H]DPCPX binding with several adenosine agonists and antagonists. Adenosine receptor antagonists appeared to bind to a single class of binding site, both in membranes and intact cells. The order of potency was DPCPX = CGS 15943A > 8-cyclopentyl-1,3-dimethylxanthine > 8-(p-sulphophenyl)-theophylline > 3-isobutyl-1-methylxanthine > theophylline. Competition curves with adenosine agonists in membranes were best described by a 2-site rather than a 1-site model. At equilibrium in intact cells, only a single site was detected at both 4° and 25°. However, short term incubations (1-4 min) at 25° showed biphasic binding curves in intact cells. The equilibrium KD values for intact cells were similar to the low affinity KD values in membranes (KL). The order of potency was N6-cyclopentyladenosine ≥ (-)-(R)-N6-phenylisopropyladenosine[(R)-PIA] ≥ N6-cyclohexyl adenosine > 5'-N-ethylcarboxamidoadenosine NECA > 2-chloroadenosine > adenosine (intact cells only) > 2-phenylaminoadenosine (CV 1808). Treatment of cells with pertussis toxin ADP-ribosylated GTP-binding proteins and eliminated the high-affinity agonist binding in membranes but did not affect binding to intact cells. The addition of GTP (100 μM) also shifted the competition curves from bi- to monophasic curves in membranes. Adenosine receptor agonists inhibited the formation of cAMP induced by isoprenaline (IC50 for (R)-PIA, 0.4 nM). This inhibition could be prevented with adenosine receptor antagonists. Pretreatment with pertussis toxin also reversed these effects and actually revealed functional A2 receptors, as shown by the formation of cAMP

McIntosh

induced by NECA. In conclusion, the equilibrium binding of A1 receptor agonists to intact smooth muscle cells is similar to the low affinity binding observed in membranes. In addition, it is suggested that agonists may transiently convert the A1 receptor from a resting low-affinity state to a high-affinity state coupled to a GTP-binding protein. DDT1 MF-2 cells should prove useful for studying regulation of A1 receptor signalling in intact cells.

IT 53296-10-9  
 RL: BIOL (Biological study)  
 (adenosine A1 receptor binding by, in smooth muscle cells)  
 RN 53296-10-9 CAPLUS  
 CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 22 THERE ARE 22 CAPLUS RECORDS THAT CITE THIS RECORD (23 CITINGS)

L6 ANSWER 132 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1990:604633 CAPLUS

DN 113:204633

OREF 113:34369a,34372a

TI Hemodynamic effects of adenosine agonists in the conscious spontaneously hypertensive rat

AU Webb, R. L.; McNeal, R. B., Jr.; Barclay, B. W.; Yasay, G. D.

CS Pharm. Div., Ciba-Geigy Corp., Summit, NJ, 07901, USA

SO Journal of Pharmacology and Experimental Therapeutics (1990), 254(3), 1090-9

CODEN: JPETAB; ISSN: 0022-3565

DT Journal

LA English

AB The hemodynamic mechanisms contributing to the reduction in blood pressure were studied in conscious spontaneously hypertensive rats after systemic administration of adenosine agonists. The effects produced by i.v. and intraarterial injections of 2-phenylaminoadenosine (CV-1808, adenosine A2 selective agonist), 5'-N-ethylcarboxamide adenosine (NECA, nonselective agonist), 2-chloroadenosine (2-CADO, A1 selective agonist), and cyclopentyladenosine (CPA, A1 selective agonist) were evaluated and compared to those of hydralazine. All agents produced hypotensive effects after bolus i.v. injections. Although CPA, NECA, and 2-CADO elicited dose-dependent bradycardia, CV-1808 and hydralazine increased the heart rate. These effects, with the exception of hydralazine-evoked responses, were attenuated by prior treatment with 8-(p-sulphophenyl)theophylline (2 mg/kg/min), whereas both CV-1808 and hydralazine produced regional vasodilation; increases in blood flow occurred only after CV-1808 (3-30 µg/kg). The regional hemodynamic responses to NECA were more complex; low doses (0.1-1 µg/kg) produced consistent redns. in regional vascular resistance, whereas at the highest dose renal vasoconstriction occurred. Although regional vasodilation occurred after 2-CADO, mesenteric vasoconstriction was observed subsequent to CPA administration. Whereas increases in renin release were evident in animals treated with CV-1808 and hydralazine, no changes occurred in response to the NECA-, 2-CADO- or CPA-induced hypotension. The predominant hemodynamic response after selective activation of A2 receptors is the regional vasodilation and hypotension leading to a reflex increase in heart rate and renin release. The reduction in arterial pressure seen after A1 receptor activation is associated primarily with a reduction in heart rate and an inhibition of renin release. NECA and 2-CADO are nonselective adenosine agonists capable of activating both A1 and A2 receptors in the conscious spontaneously hypertensive rat.

IT 53296-10-9, 2-Phenylaminoadenosine  
 RL: PRP (Properties)


10/598,520

(hemodynamic effects of, adenosine receptors in, in hypertension)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 21 THERE ARE 21 CAPLUS RECORDS THAT CITE THIS RECORD (21 CITINGS)

L6 ANSWER 133 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1990:545750 CAPLUS

DN 113:145750

OREF 113:24593a,24596a

TI Characterization of the adenosine receptor in porcine coronary arteries

AU King, A. D.; Milavec-Krizman, M.; Mueller-Schweinitzer, E.

CS Sandoz Pharma A.-G., Basel, CH-4002, Switz.

SO British Journal of Pharmacology (1990), 100(3), 483-6

CODEN: BJPCBM; ISSN: 0007-1188

DT Journal

LA English

AB Relaxant responses of ring preps. from porcine ventricular coronary arteries to adenosine and various stable adenosine analogs were investigated in vitro. The adenosine analogs did not produce contraction but elicited almost complete relaxation of coronary arteries precontracted with 3  $\mu$ M prostaglandin F2 $\alpha$  (PGF2 $\alpha$ ), even after removal of the endothelium. The order of potency was 5'-N-ethylcarboxamide-adenosine (NECA) > 2-(2-phenylethylamino)-5'-N-ethylcarboxamide-adenosine (2-PEA-NECA) > 2-phenylamino-adenosine (CV-1808) > N6-[R(-)-1-phenyl-2-propyl]adenosine (R-PIA) > N6-[S(+)-1-phenyl-2-propyl]adenosine (S-PIA) > N6-cyclopentyladenosine (CPA) > adenosine > ATP = ADP, which suggested the presence of adenosine A2-receptor subtypes. There was an excellent correlation between the calculated pD2 values on coronary arteries and the pKD values at adenosine A2 binding sites, whereas no correlation was obtained when the pD2 values were compared to the pKD values at adenosine A1-binding sites on membranes from porcine striata. The relaxant effects of adenosine and its analogs were competitively antagonized by 8-(p-sulfophenyl)theophylline (8-SPT), producing pA2 values similar to the resp. pKD value of the antagonist at adenosine A2 binding sites. It is suggested that the porcine coronary artery possesses adenosine A2 receptors which seem to be similar to the adenosine A2 binding site in pig striatum, whereas no evidence was obtained for the presence of adenosine A1 receptors.

IT 53296-10-9, CV-1808


RL: BIOL (Biological study)

(coronary artery relaxation induction by, purinergic A2 receptors in mediation of)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



McIntosh

OSC.G 12 THERE ARE 12 CAPLUS RECORDS THAT CITE THIS RECORD (12 CITINGS)

L6 ANSWER 134 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1990:545035 CAPLUS

DN 113:145035

OREF 113:24425a,24428a

TI Adenosine receptors and modulation of natural killer cell activity by purine nucleosides

AU Priebe, Teresa; Platsoucas, Chris D.; Nelson, J. Arly

CS M. D. Anderson Cancer Cent., Univ. Texas, Houston, TX, 77030, USA

SO Cancer Research (1990), 50(14), 4328-31

CODEN: CNREA8; ISSN: 0008-5472

DT Journal

LA English

AB Natural killer (NK) cell activity is inhibited in vivo by the adenosine analog tubercidin (Tub) and stimulated by the deoxyadenosine analog 2-fluoro-1- $\beta$ -D-arabinofuranosyladenine 5'-monophosphate (F-ara-AMP) in the spleen lymphocytes from mice. The inhibition by Tub and stimulation by F-ara-AMP of NK cell activity are readily demonstrable in murine and human lymphocytes exposed to the drugs in vitro. In mouse spleen lymphocytes, NK cell activity is also inhibited by adenosine receptor A2 agonists, whereas potent A1 receptor agonists are more effective stimulators. Inhibition produced by adenosine, deoxyadenosine, and adenosine receptor agonists, but not by Tub, is partially prevented by the adenosine receptor antagonist 1,3-dipropyl-8-phenylxanthine amine congener. Agents that stimulate NK cell activity (deoxyadenosine, A1 receptor agonists, F-ara-AMP) do not increase further the 1.5-fold enhancement produced by a 10<sup>-6</sup>M 1,3-dipropyl-8-phenylxanthine amine congener. The nucleoside transport inhibitor p-nitrobenzylthioinosine 5'-monophosphate has no effect on NK cell activity or intracellular ribonucleotide pools; however, it partially prevents Tub 5'-triphosphate formation, ATP depletion, and NK cell inhibition in mouse spleen cells treated with Tub. Nitrobenzylthioinosine 5'-monophosphate also partially prevents the F-ara-AMP stimulation of NK cell activity, but it does not influence the effects of adenosine or deoxyadenosine. The results obtained with the adenosine receptor agonists suggest roles for both A1 and A2 receptors in regulating murine NK cell activity. Tub inhibition of NK cell activity does not involve adenosine receptors; however, inhibition by the other agents may be mediated via an A2 receptor (stimulatory for adenylyl cyclase). Since p-nitrobenzylthioinosine 5'-monophosphate inhibited the stimulation of NK cell activity by F-ara-AMP, this stimulation may occur via an intracellular P site (inhibitory to adenylyl cyclase).

IT 53296-10-9, 2-Phenylaminoadenosine


RL: BIOL (Biological study)

(spleenocyte natural killer activity modulation by, adenosine receptors in)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 19 THERE ARE 19 CAPLUS RECORDS THAT CITE THIS RECORD (19 CITINGS)

L6 ANSWER 135 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1990:508747 CAPLUS

DN 113:108747

OREF 113:18193a,18196a

TI Study of the lipophilic character of xanthine and adenosine derivatives.


II. Relationships between log k', RM and log P values

AU Biagi, G. L.; Guerra, M. C.; Barbaro, A. M.; Barbieri, S.; Recanatini, M.;

10/598,520

Borea, P. A.  
CS Ist. Farmacol., Univ. Bologna, Bologna, Italy  
SO Journal of Liquid Chromatography (1990), 13(5), 913-27  
CODEN: JLCHD8; ISSN: 0148-3919  
DT Journal  
LA English  
AB The log k' values of a series of xanthine and adenosine derivs. were measured by reversed-phase HPLC. The HPLC data correlated with previously reported RM and RMC18 values. The equations describing the relationships log k'/RM and log k'/RMC18 allowed the calcn. of the log k' values of some compds. which were not tested in the HPLC system. Since the relationship log k'/log P is very close to the previously described relationships RM/log P and RMC18/log P, reversed-phase TLC and HPLC are very similar in describing the lipophilicity of the compds.  
IT 53296-10-9, 2-Phenylaminoadenosine  
RL: PRP (Properties)  
(lipophilicity of, reversed-phase HPLC in determination of)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.




OSC.G 15 THERE ARE 15 CAPLUS RECORDS THAT CITE THIS RECORD (15 CITINGS)

L6 ANSWER 136 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1990:454059 CAPLUS  
DN 113:54059  
OREF 113:9041a,9044a  
TI 2-(p-Nitrophenyl)-2'-deoxyadenosine, a new type of mutagenic nucleoside  
AU Matsuda, Akira; Ohara, Yoshiko; Kakutani, Toshifumi; Negishi, Kazuo;  
Wataya, Yusuke; Hayatsu, Hikoya; Ueda, Tohru  
CS Fac. Pharm. Sci., Hokkaido Univ., Sapporo, 060, Japan  
SO Nucleic Acids Research (1990), 18(7), 1833-8  
CODEN: NARHAD; ISSN: 0305-1048  
DT Journal  
LA English  
AB A crude preparation of 2-phenyladenosine was mutagenic in the Ames salmonella assay. In the purification of this preparation, it was revealed that 2-phenyladenosine itself was nonmutagenic but that 2-(m- and p-nitrophenyl)adenosines (5m,p) contaminating the sample were the mutagenic principles. A structure-activity relationship study was carried out, and it was found that 5p, 2-(p-nitrophenyl)adenine (7p), and 2-(p-nitrophenyl)-2'-deoxyadenosine (15p) were strongly mutagenic toward S. typhimurium TA98 and TA100 without metabolic activation, the potency being in the order 15p > 7p > 5p. The potency of 15p in TA98 was one order of magnitude greater than that of 4-nitroquinoline N-oxide. The 15p also showed mutagenicity in the mouse cell line FM3A in culture.  
IT 109875-49-2P  
RL: ADV (Adverse effect, including toxicity); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)  
(preparation and mutagenicity of, in Ames test)  
RN 109875-49-2 CAPLUS  
CN Adenosine, 2-(4-aminophenyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

McIntosh



OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

L6 ANSWER 137 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1990:435339 CAPLUS

DN 113:35339

OREF 113:5913a,5916a

TI Adenosine receptors are coupled negatively to release of tachykinin(s) from enteric nerve endings

AU Christofi, F. L.; McDonald, T. J.; Cook, M. A.

CS Dep. Pharmacol. Toxicol., Univ. West. Ontario, London, ON, N6A 5C1, Can.

SO Journal of Pharmacology and Experimental Therapeutics (1990), 253(1), 290-5

CODEN: JPETAB; ISSN: 0022-3565

DT Journal

LA English

AB Adenosine receptors capable of modulating tachykininergic transmission were characterized in functional studies using both field-stimulated and cholecystokinin octapeptide-stimulated contractile responses of atropinized guinea pig longitudinal muscle-myenteric plexus preps. These tetrodotoxin-sensitive responses, which were mediated by release of one or more tachykinins, were inhibited by adenosine analogs in a concentration-dependent manner. The rank order of potencies of the analogs as inhibitors of the responses to cholecystokinin octapeptide was: N6-cyclopentyladenosine > 5'-N-ethylcarboxamidoadenosine >> 2-phenylaminoadenosine (CV 1808). Schild anal. of the antagonism of the presynaptic inhibitory effects of 5'-N-ethylcarboxamidoadenosine and N6-cyclopentyladenosine on cholecystokinin octapeptide-stimulated responses using the A1 selective antagonists 1,3-dipropyl-8-(4-sulfophenyl)xanthine and 1,3-dipropyl-8-(cyclopentyl)xanthine yielded linear isoboles with unit slopes indicating competitive antagonism. The affinity of the antagonists for the receptor site(s) involved in inhibition of tachykininergic transmission was similar to those established previously for cholinergic transmission. The rank order of potency of adenosine analogs as inhibitors of the field-stimulated responses was such that N6-cyclopentyladenosine = 5'-ethylcarboxamidoadenosine. Reverse-phase HPLC anal. performed on lysates of isolated myenteric nerve endings demonstrated the presence of substance P and neurokinin-A. Neurokinin-B was undetectable. These studies indicate that adenosine receptor(s) on myenteric nerve endings are coupled neg. to tachykinin release and that they are probably identical to those involved in the modulation of acetylcholine release.

IT 53296-10-9, CV 1808


RL: BIOL (Biological study)

(tachykinin-induced contractions of ileum-myenteric plexus preparation inhibition by)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)


Absolute stereochemistry.



10/598,520


OSC.G 14 THERE ARE 14 CAPLUS RECORDS THAT CITE THIS RECORD (14 CITINGS)

L6 ANSWER 138 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1990:424426 CAPLUS  
DN 113:24426  
OREF 113:4255a,4258a  
TI 2-(Arylalkylamino)adenosin-5'-uronamides: a new class of highly selective  
adenosine A2 receptor ligands  
AU Hutchison, Alan J.; Williams, Michael; De Jesus, Reynalda; Yokoyama, Rina;  
Oei, Howard H.; Ghai, Geetha R.; Webb, Randy L.; Zoganas, Harry C.; Stone,  
George A.; Jarvis, Michael F.  
CS Pharm. Div., Ciba-Geigy Corp., Summit, NJ, 07901, USA  
SO Journal of Medicinal Chemistry (1990), 33(7), 1919-24  
CODEN: JMCMAR; ISSN: 0022-2623  
DT Journal  
LA English  
OS CASREACT 113:24426  
GI



AB The synthesis and receptor-binding profiles at adenosine receptor subtypes for a series of 2-arylalkylamino-adenosine-5'-uronamides is described. Halogenated 2-phenethylamino analogs such as I (R = Cl) show greater than 200-fold selectivity for the A2 receptor subtype on the basis of rat brain receptor binding. The general structure-activity relationship of this series of compds. is discussed both in terms of potency at A2 receptors as well as receptor subtype selectivity. It is possible to introduce a hydrophilic carboxyalkyl substituent to this series such as in CGS 21680A (I; R = HO2CCH2CH2) and still retain good potency and selectivity for A2 receptors. In addition, functional data in a perfused working rat heart model shows that these compds. possess full agonist properties at A2 receptors with I (R = HO2CCH2CH2) having a greater than 1500-fold separation between A2 (coronary vasodilatory) and A1 (neg. chronotropic) receptor mediated events.  
IT 53296-10-9DP, CV1808, analogs  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation and adenosine acceptor selectivity of)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 40 THERE ARE 40 CAPLUS RECORDS THAT CITE THIS RECORD (40 CITINGS)


L6 ANSWER 139 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1990:135402 CAPLUS

McIntosh

10/598,520


DN 112:135402  
OREF 112:22805a,22808a  
TI Study of the lipophilic character of xanthine and adenosine derivatives.  
I. RM and log P values  
AU Biagi, G. L.; Guerra, M. C.; Barbaro, A. M.; Barbieri, S.; Recanatini, M.;  
Borea, P. A.; Pietrogrande, M. C.  
CS Ist. Farmacol., Univ. Bologna, Bologna, 40126, Italy  
SO Journal of Chromatography (1990), 498(1), 179-90  
CODEN: JOCRAM; ISSN: 0021-9673  
DT Journal  
LA English  
AB The RM values of a series of xanthine and adenosine derivs. were measured  
using silicone reversed-phase TLC and C18 reversed-phase high-performance  
TLC systems. The 2 series of data were well correlated. Both were  
compared with exptl. log P and calculated CLOGP values. For xanthine derivs.,  
a good linear relation was shown between the RM values from the 2  
chromatog. systems and the log P or CLOGP data. For adenosine derivs.,  
the CLOGP values had to be corrected to fit the data to the same equation. The  
TLC data proved to be reliable parameters for describing the lipophilic  
properties of the test compds.  
IT 53296-10-9, 2-Phenylaminoadenosine  
RL: PRP (Properties)  
(lipophilicity of, determination of, by reversed-phase high-performance TLC and  
reversed-phase TLC, octanol-water partition coefficient in relation to)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 18 THERE ARE 18 CAPLUS RECORDS THAT CITE THIS RECORD (18 CITINGS)

L6 ANSWER 140 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1990:119289 CAPLUS  
DN 112:119289  
OREF 112:20227a,20230a  
TI Synthesis of congeners of adenosine resistant to deamination by adenosine  
deaminase  
AU Nair, Vasu; Purdy, David F.; Sells, Todd B.  
CS Dep. Chem., Univ. Iowa, Iowa City, IA, 52242, USA  
SO Journal of the Chemical Society, Chemical Communications (1989), (13),  
878-9  
CODEN: JCCCAT; ISSN: 0022-4936  
DT Journal  
LA English  
OS CASREACT 112:119289  
GI




McIntosh



10/598,520

AB The metal-mediated preparation of deaminase resistant adenosine congeners I [R = CH<sub>2</sub>:CH, HOCH<sub>2</sub>CH(OH), Et, F<sub>3</sub>C, cyano] from I (R = iodo) is described.  
IT 79936-11-1P  
RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation)  
(preparation and resistance of, to deamination by adenosine deaminase)  
RN 79936-11-1 CAPLUS  
CN Adenosine, 2-cyano- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




OSC.G 15 THERE ARE 15 CAPLUS RECORDS THAT CITE THIS RECORD (15 CITINGS)


L6 ANSWER 141 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1990:116389 CAPLUS  
DN 112:116389  
OREF 112:19659a,19662a  
TI Adenosine transporters in vascular smooth muscle and endothelium:  
multiple [3H]nitrobenzylthioinosine binding sites in human umbilical vein  
endothelium  
AU Williams, Evan F.; Harris-Hooker, Sandra; Gordon, Portia B.  
CS Dep. Pharmacol., Morehouse Sch. Med., Atlanta, GA, USA  
SO Drug Development Research (1990), 19(1), 79-90  
CODEN: DDREDK; ISSN: 0272-4391  
DT Journal  
LA English  
AB Cultured vascular smooth muscle and endothelial cells may be useful models  
for studying the cardiovascular adenosine transport system and metabolism The  
nucleoside transporter elements of cultured primate vascular smooth  
muscle, bovine aortic endothelial, and human umbilical vein endothelial  
cells were quantified by radioligand binding and by using membrane preps.  
of these cells and the nucleoside transporter probe nitrobenzylthioinosine  
([3H]NBMPR), a potent and tightly bound inhibitor of nucleoside transport.  
The binding was rapid, reversible, saturable, and site-specific.  
Scatchard anal. of the saturation data showed that [3H]NBMPR bound to high and  
low affinity binding sites in human umbilical vein endothelial cell  
membranes with apparent binding affinities (K<sub>D</sub>) of 0.093 nM and 1.92 nM,  
and binding site densities (B<sub>max</sub> values) of 13.48 and 69 fmol/mg protein,  
resp. In contrast, the binding to primate vascular smooth muscle and  
bovine aortic endothelial cell membranes occurred to an apparently high  
affinity single class of binding sites at which the K<sub>D</sub> was 1.4 nM and 0.28  
nM, resp., and which had B<sub>max</sub> values of 1,977 and 1,284 fmol/mg protein,  
resp. Scatchard anal. of the binding inhibition by dipyridamole showed a  
mixed type inhibition, while NBMPR inhibited the binding competitively.  
Several recognized nucleoside transport inhibitors and vasodilators  
inhibited the binding with an order of potency similar to that observed for  
the inhibition of [3H]NBMPR binding to guinea pig cardiac membranes.  
IT 53296-10-9, 2-Phenylaminoadenosine  
RL: PROC (Process)  
(binding of, to vascular smooth muscle and endothelium of humans and  
laboratory animals, adenosine transporter in relation to)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.

McIntosh



L6 ANSWER 142 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
 AN 1990:112580 CAPLUS  
 DN 112:112580  
 OREF 112:18911a,18914a  
 TI [3H]CGS 21680, a selective A2 adenosine receptor agonist directly labels  
 A2 receptors in rat brain  
 AU Jarvis, Michael F.; Schulz, Rainer; Hutchison, Alan J.; Do, Un Hoi; Sills,  
 Matthew A.; Williams, Michael  
 CS Pharm. Div., Ciba-Geigy Corp., Summit, NJ, 07901, USA  
 SO Journal of Pharmacology and Experimental Therapeutics (1989), 251(3),  
 888-93  
 CODEN: JPETAB; ISSN: 0022-3565  
 DT Journal  
 LA English  
 GI



I

AB Characterization of the adenosine A2 receptor has been limited due to the lack of available ligands which have high affinity and selectivity for this adenosine receptor subtype. In the present study, the binding of a highly A2-selective agonist radioligand, [3H]CGS 21680 (I) is described. [3H]CGS 21680 specific binding to rat striatal membranes was saturable, reversible, and dependent on protein concentration. Saturation studies revealed that [3H]CGS 21680 bound with high affinity ( $K_d = 15.5$  nM) and limited capacity (apparent  $B_{max} = 375$  fmol/mg protein) to a single class of recognition sites. Ests. of ligand affinity (16 nM) determined from association and dissociation kinetic expts. were in close agreement with the results from the saturation studies. [3H]CGS 21680 binding was greatest in striatal membranes with negligible specific binding obtained in rat cortical membranes. Adenosine agonists ligands competed for the binding of 5 nM [3H]CGS 21680 to striatal membranes with the following order of activity; CGS 21680 = 5'-N-ethylcarboxamidoadenosine > 2-phenylaminoadenosine (CV-1808); 5'-N-methylcarboxamidoadenosine = 2-chloroadenosine > R-phenylisopropyladenosine > N6-cyclohexyladenosine > N6-cyclopentyltheophylline > S-phenylisopropyladenosine. The nonxanthine adenosine antagonist, CGS 15943A, was the most active compound in inhibiting the binding of [3H]CGS 21680. Other adenosine antagonists inhibited binding in the following order; xanthine amine congener = (1,3-dipropyl-8-(2-amino-4-chloro)phenylxanthine > 1,3-dipropyl-8-cyclopentylxanthine > 1,3-diethyl-8-phenylxanthine > 8-phenyltheophylline > 8-cyclopentyltheophylline = xanthine carboxylic acid congener > 8-parasulfophenyltheophylline > theophylline > caffeine. The pharmacol. profile of both adenosine agonist and antagonist compds. to compete for the binding of [3H]CGS 21680 was consistent with a selective interaction at the high affinity adenosine A2 receptor. A high pos. correlation was observed between the pharmacol. profile of adenosine ligands to inhibit the binding of [3H]CGS 21680 and the selective binding of

[3H]NECA (+50 nM CPA) to high affinity A2 receptors. However, some differences between these assays were found for compds. which have moderate affinity and nonselective actions at both the A1 and A2 adenosine receptor subtypes. Unlike data obtained with nonselective adenosine ligands, the present results indicate that [3H]CGS 21680 directly labels the high affinity A2 receptor in rat brain without the need to block binding activity at the A1 receptor. The high degree of selectivity (>170-fold) and high affinity of [3H]CGS 21680 make this the current ligand of choice for the in vitro characterization of high affinity A2 receptors.

IT 53296-10-9, CV 1808


RL: BIOL (Biological study)

(CGS 21680 binding by purinergic receptors inhibition by, in brain striatum)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 267 THERE ARE 267 CAPLUS RECORDS THAT CITE THIS RECORD (267 CITINGS)

L6 ANSWER 143 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1990:91613 CAPLUS

DN 112:91613

OREF 112:15383a,15386a

TI A selective binding site for 3H-NECA that is not an adenosine A2 receptor

AU Keen, Mary; Kelly, Eamonn; Nobbs, Peter; MacDermot, John

CS Med. Sch., Univ. Birmingham, Birmingham, B15 2TJ, UK

SO Biochemical Pharmacology (1989), 38(21), 3827-33

CODEN: BCPA6; ISSN: 0006-2952

DT Journal

LA English

AB In homogenates of NG108-15 cells, adenosine analogs activate adenylate cyclase with the following order of potency: N-ethylcarboxamidoadenosine (NECA) > 2-chloroadenosine > N6-(L-phenylisopropyl)adenosine (PIA) = cyclohexyladenosine = 2-phenylaminoadenosine. Adenosine receptor antagonists inhibit NECA-stimulated adenylate cyclase activity with the order of potency 3-isobutyl-1-methyl-xanthine (IBMX) > theophylline > caffeine. These data suggest that these ligands act at an adenosine A2 receptor. There is an apparently homogeneous population of saturable 3H-NECA binding sites in homogenates of NG108-15 cells. These sites have an affinity for 3H-NECA of .apprx.1  $\mu$ M and are present at a d. of .apprx.10 pmol/mg protein. Unlabeled NECA, 2-chloroadenosine, IBMX and theophylline displace 3H-NECA binding, with an order of potency that suggests that the 3H-NECA binding site may represent an adenosine A2 receptor. However, PIA, cyclohexyladenosine and 2-phenylaminoadenosine produce no detectable displacement of 3H-NECA binding at concns. that produce a maximal stimulation of adenylate cyclase activity. Pretreatment of NG108-15 cells with either NECA or PIA produces a homologous desensitization of subsequent responses to all the adenosine analogs, with no effect on subsequent responses to a prostacyclin receptor agonist or NaF. This suggests that all the adenosine analogs examined activate an adenosine A2 receptor. Therefore, the 3H-NECA site at which PIA is inactive cannot represent this receptor.

IT 53296-10-9

RL: BIOL (Biological study)


(adenosine receptors binding of ethylcarboxamidoadenosine response to, adenylate cyclase in)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)


10/598,520

Absolute stereochemistry.



OSC.G 6 THERE ARE 6 CAPLUS RECORDS THAT CITE THIS RECORD (6 CITINGS)

L6 ANSWER 144 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1990:30236 CAPLUS  
DN 112:30236  
OREF 112:5069a,5072a  
TI Effects of adenosine A2 receptor agonists on nucleoside transport  
AU Balwierczak, Joseph L.; Krulan, Christine M.; Wang, Zhi Chao; Chen, Jen; Jeng, Arco Y.  
CS Res. Dep., Ciba-Geigy Corp., Summit, NJ, 07901, USA  
SO Journal of Pharmacology and Experimental Therapeutics (1989), 251(1), 279-87  
CODEN: JPETAB; ISSN: 0022-3565  
DT Journal  
LA English  
GI



AB A series of adenosine A2 receptor agonists I (R = CH2OH or CONHET; R1 = Ph, etc.; X = O or CH2) were examined for their ability to activate adenosine A2 receptors and inhibit nucleoside transport. A2 receptor activation was measured by the ability of these adenosine agonists to relax porcine coronary smooth muscle, where I varied in their EC50 values. Nucleoside transport was measured as the nitrobenzylthioinosine-sensitive cellular accumulation of [3H]uridine into guinea pig erythrocytes at 22°. The initial velocity of transport was dependent on substrate concentration and a substrate-velocity curve yielded a Km of 78 µM and a Vmax of 0.31 mmol/L of cell water per h. Dipyridamole, a known potent inhibitor of nucleoside transport, blocked cellular [3H]uridine accumulation with an EC50 of 29.4 nM. Whereas a number of the adenosine agonists tested showed little or no inhibition of nucleoside transport, CV 1808 inhibited transport with an EC50 of 140 nM. In addition, 2 carbocyclic derivs. of CV 1808, CGS 23321 and CGS 23302 inhibited nucleoside transport with resp. EC50 values of 366 and 168 nM. The data suggest that these compds. have a different structure-activity relationship for adenosine A2 receptors and for the site mediating nucleoside transport inhibition.


IT 53296-10-9, CV 1808  
RL: BIOL (Biological study)  
(nucleoside transport by erythrocyte response to and artery relaxation by)

RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




OSC.G 5 THERE ARE 5 CAPLUS RECORDS THAT CITE THIS RECORD (5 CITINGS)

L6 ANSWER 145 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1989:574576 CAPLUS  
DN 111:174576  
OREF 111:29091a,29094a  
TI Novel, stable congeners of the antiretroviral compound  
2',3'-dideoxyadenosine  
AU Nair, Vasu; Buenger, Greg S.  
CS Dep. Chem., Univ. Iowa, Iowa City, IA, 52242, USA  
SO Journal of the American Chemical Society (1989), 111(22), 8502-4  
CODEN: JACSAT; ISSN: 0002-7863  
DT Journal  
LA English  
OS CASREACT 111:174576  
GI



AB Novel congeners I (R = cyano, Et, SMe, iodo, CF3) and the 2',3'-dideoxy analog of I (R = cyano) of the antiretroviral compound 2',3'-dideoxyadenosine (I, R = H) have been synthesized through metal-mediated and photochem. conversions as the key steps. These compds. are inherently more stable than I (R = H) with respect to both glycosidic bond cleavage and deamination by adenosine deaminase.  
IT 79936-11-1P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(preparation, silylation, and thiocarbonylation of)  
RN 79936-11-1 CAPLUS  
CN Adenosine, 2-cyano- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




OSC.G 34 THERE ARE 34 CAPLUS RECORDS THAT CITE THIS RECORD (34 CITINGS)

McIntosh

10/598,520

L6 ANSWER 146 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1989:511808 CAPLUS  
DN 111:111808  
OREF 111:18687a,18690a  
TI Affinity chromatography of A1 adenosine receptors of rat brain membranes  
AU Nakata, Hiroyasu  
CS Lab. Clin. Sci., Natl. Inst. Ment. Health, Bethesda, MD, 20892, USA  
SO Molecular Pharmacology (1989), 35(6), 780-6  
CODEN: MOPMA3; ISSN: 0026-895X  
DT Journal  
LA English  
AB The A1 adenosine receptor of rat brain membranes has been solubilized with digitonin and purified .apprx.150-fold by affinity chromatog. The digitonin-solubilized receptor, which can be labeled with 8-cyclopentyl-1,3-[3H]dipropylxanthine ([3H]DPCPX), was adsorbed on xanthine amine congener (XAC)-linked agarose. The interaction of the solubilized receptor activity with the affinity gel was biospecific. Adenosine agents blocked adsorption of solubilized receptor activity to the XAC-agarose with the appropriate A1 adenosine selectivity. For agonists, 8-cyclopentyladenosine > (R)-phenylisopropyladenosine > CV-1808, whereas, for antagonists, 8-cyclopentyltheophylline (CPT) > XAC > isobutylmethylxanthine = theophylline. The same A1 adenosine receptor specificity was observed for elution of [3H]DPCPX binding activity from the gel. XAC-agarose adsorbed 65-80% of the solubilized [3H]DPCPX binding activity and, after the gel was washed, 30-40% of the adsorbed activity could be eluted with 100  $\mu$ M CPT, with specific binding activity of .apprx.60 pmol/mg of protein. The order of potency of adenosine agonists [8-Cyclopentyladenosine > (R)-phenylisopropyladenosine > 5'-N-ethylcarboxamidoadenosine > (S)-phenylisopropyladenosine] and antagonists (DPCPX > XAC > CPT > isobutylmethylxanthine) with the affinity-purified preparation was found to be similar to that of the solubilized adenosine A1 receptor. This affinity chromatog. procedure should prove to be valuable in the isolation and mol. characterization of A1 adenosine receptors.  
IT 53296-10-9, CV-1808  
RL: ANST (Analytical study)  
(A1 adenosine receptors affinity chromatog. of brain membranes on xanthine amine congener-agarose gel response to)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)


Absolute stereochemistry.



OSC.G 5 THERE ARE 5 CAPLUS RECORDS THAT CITE THIS RECORD (5 CITINGS)

L6 ANSWER 147 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1989:458249 CAPLUS  
DN 111:58249  
OREF 111:9899a,9902a  
TI C2,N6-Disubstituted adenosines: synthesis and structure-activity relationships  
AU Trivedi, Bharat K.; Bruns, Robert F.  
CS Parke-Davis Pharm. Res. Div., Warner-Lambert Co., Ann Arbor, MI, 48105, USA  
SO Journal of Medicinal Chemistry (1989), 32(8), 1667-73  
CODEN: JMCMAR; ISSN: 0022-2623  
DT Journal  
LA English  
OS CASREACT 111:58249  
GI

McIntosh



AB Extracellular adenosine receptors have been divided into two major subtypes, called A1 and A2. Substitution of the adenosine mol. with appropriate groups at C-2 or N-6 is known to impart selectivity for the A2 receptor over the A1 receptor. The present study investigated whether substitution at both C-2 and N-6 would have additive effects on the A2/A1 affinity ratio, thereby providing compds. with greater A2 selectivity than presently available agents. Disappointingly, additivity appeared to hold only when an A1-selective group was present at N-6. For instance, conversion of the A1-selective agonist I (R = H, R1 = cyclopentyl) to I (R = NHPh, R1 = cyclopentyl) resulted in a 70-fold shift in selectivity in favor of the A2 receptor, but the same substitution applied to the A2-selective agonist I [R = H, R1 = 3,5-(MeO)2C6H3CHPhCH2] resulted in a 100-fold loss of affinity with no change in A2-selectivity.

IT 53296-10-9


RL: PRP (Properties)

(adenosine receptor affinity of)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 18 THERE ARE 18 CAPLUS RECORDS THAT CITE THIS RECORD (18 CITINGS)

L6 ANSWER 148 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1989:451028 CAPLUS

DN 111:51028

OREF 111:8557a,8560a

TI An unusual receptor mediates adenosine-induced SA nodal bradycardia in dogs

AU Belloni, Francis L.; Belardinelli, Luiz; Halperin, Cidio; Hintze, Thomas H.

CS Dep. Physiol., New York Med. Coll., Valhalla, NY, 10595, USA

SO American Journal of Physiology (1989), 256(6, Pt. 2), H1553-H1564

CODEN: AJPHAP; ISSN: 0002-9513

DT Journal


LA English

AB To characterize the receptor mediating the neg. chronotropic effect of adenosine in dogs, expts. were performed on conscious dogs with chronically implanted cardiovascular instrumentation. Autonomic blockade was used to eliminate any reflex influences on heart rate. I.v. bolus injections of various adenosine analogs caused dose-dependent, aminophylline-blockable redns. in heart rate with a potency order for NECA:2-chloroadenosine:adenosine of 78:17:1. Dipyrindamole enhanced the

potency of adenosine to equal that of 2-chloroadenosine. Moderately selective A1-receptor agonists (N6-(L-2-phenylisopropyl)-adenosine (R-PIA) and N6-cyclohexyladenosine) and an A2-selective agonist (2-phenylaminoadenosine) had no neg. chronotropic effect in the conscious dog. Adenosine and its analogs, including R-PIA, caused coronary vasodilation at smaller doses than were required to slow the heart rate. The selective A1-adenosine receptor blocker xanthine amine congener (XAC) antagonized the neg. chronotropic action of adenosine, but did so nonselectively, as the coronary vasodilative and neg. chronotropic actions of adenosine were antagonized equally well. The spontaneous contraction rate of isolated perfused dog right atrial preps., which included the sinoatrial (SA) node, was reduced by intrasinoatrial node artery infusions of adenosine analogs with a potency ratio for NECA:adenosine:N6-cyclopentyladenosine:R-PIA of 100:15:2.3:1. Apparently, the adenosine receptor mediating the neg. chronotropic action of adenosine in the dog does not display the pharmacol. characteristics of either typical A1- or A2-adenosine receptors. Instead, either a novel adenosine receptor or an A1-receptor with unusual agonist and antagonist binding properties appears to exist in the dog's sinoatrial node.

IT 53296-10-9, 2-Phenylaminoadenosine  
 RL: BIOL (Biological study)  
 (heart rate response to, receptors and mediation of)  
 RN 53296-10-9 CAPLUS  
 CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 5 THERE ARE 5 CAPLUS RECORDS THAT CITE THIS RECORD (5 CITINGS)


L6 ANSWER 149 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
 AN 1989:421457 CAPLUS  
 DN 111:21457  
 OREF 111:3723a,3724a  
 TI NECA-induced hypomotility in mice: evidence for a predominantly central site of action  
 AU Durcan, Michael J.; Morgan, Philip F.  
 CS Lab. Clin. Stud., Natl. Inst. Alcohol Abuse Alcohol., Bethesda, MD, 20892, USA  
 SO Pharmacology, Biochemistry and Behavior (1989), 32(2), 487-90  
 CODEN: PBBHAU; ISSN: 0091-3057  
 DT Journal  
 LA English  
 AB The behavioral effects of 4 adenosine analogs (NECA, cyclohexyladenosine (CHA), cyclopentyladenosine (CPA), and CV 1808) were investigated in mice using a holeboard test, which measures both directed exploration (head-dipping) and locomotor activity. NECA, CHA, and CPA showed dose-related redns. in all the holeboard measures (NECA » CHA = CPA), but CV 1808 was inactive in all of the measures over the dose range tested. In a subsequent experiment NECA-induced hypomotility was attenuated by the adenosine receptor antagonists, theophylline (which is both centrally and peripherally active) and, though to a lesser extent, by the adenosine receptor antagonist 8-(p-sulfophenyl)theophylline (8-pSPT), which poorly penetrates the blood-brain barrier. Thus, NECA-induced hypomotility may be predominantly mediated centrally since the centrally active antagonist was the most effective in reversing the effect; however, peripheral mechanisms may also play a role since equimol. concns. of 8-pSPT elicit some reversal of NECA-induced hypomotility.  
 IT 53296-10-9, CV 1808  
 RL: BIOL (Biological study)  
 (motor behavior in presence of)  
 RN 53296-10-9 CAPLUS



10/598,520

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 9 THERE ARE 9 CAPLUS RECORDS THAT CITE THIS RECORD (9 CITINGS)

L6 ANSWER 150 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1989:417561 CAPLUS

DN 111:17561

OREF 111:2963a,2966a

TI Comparison of the behavioral effects of adenosine agonists and dopamine antagonists in mice

AU Heffner, Thomas G.; Wiley, James N.; Williams, Ann E.; Bruns, Robert F.; Coughenour, Linda L.; Downs, David A.

CS Parke-Davis Pharm. Res. Div., Warner-Lambert Co., Ann Arbor, MI, 48105, USA

SO Psychopharmacology (Berlin, Germany) (1989), 98(1), 31-7

CODEN: PSCHDL; ISSN: 0033-3158

DT Journal

LA English

AB The adenosine agonists 5'-N-ethylcarboxamideadenosine (NECA), 2-chloroadenosine (2-CLA), N6-cyclohexyladenosine (CHA), N6-cyclopentyladenosine (CPA), 2-(phenylamino)adenosine (CV-1808) and R and S isomers of N6-phenylisopropyladenosine (R-PIA and S-PIA) decreased spontaneous locomotor activity in mice and, except for CPA, did so at doses that did not impair motor coordination, a profile shared by dopamine antagonists. CV-1808, the only agent with higher affinity for A<sub>2</sub> as compared with A<sub>1</sub> adenosine receptors, displayed the largest separation between locomotor inhibitory and ataxic potency. Like dopamine antagonists, NECA and CV-1808 also decreased hyperactivity caused by d-amphetamine at doses that did not cause ataxia whereas A<sub>1</sub>-selective adenosine agonists reduced amphetamine's effects only at ataxic doses. Unlike dopamine antagonists, adenosine agonists inhibited apomorphine-induced cage climbing only at doses that caused morphine-induced cage climbing only at doses that caused ataxia. Involvement of central adenosine receptors in these effects was suggested by the significant correlation obtained between potency for locomotor inhibition after i.p. and intracerebroventricular administration. Affinity for A<sub>1</sub> but not A<sub>2</sub> adenosine receptors was significantly correlated with potency for inducing ataxia. These results suggest that the behavioral profile of adenosine agonists in mice is related to their affinity for A<sub>1</sub> and A<sub>2</sub> adenosine receptors and indicate that adenosine agonists produce certain behavioral effects that are similar to those seen with dopamine antagonists.

IT 53296-10-9, 2-(Phenylamino)adenosine

RL: BIOL (Biological study)


(behavioral response to, as adenosine agonist, antipsychotic activity in relation to)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.

10/598,520



OSC.G 43 THERE ARE 43 CAPLUS RECORDS THAT CITE THIS RECORD (43 CITINGS)

L6 ANSWER 151 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1989:108716 CAPLUS

DN 110:108716

OREF 110:17818h,17819a

TI Correlation between binding affinities for brain A1 and A2 receptors of adenosine agonists and antagonists and their effects on heart rate and coronary vascular tone

AU Oel, H. H.; Ghai, G. R.; Zoganas, H. C.; Stone, G. A.; Zimmerman, M. B.; Field, F. P.; Williams, M.

CS Pharm. Div., Ciba-Geigy Corp., Summit, NJ, 07901, USA

SO Journal of Pharmacology and Experimental Therapeutics (1988), 247(3), 882-8

CODEN: JPETAB; ISSN: 0022-3565

DT Journal

LA English

AB The activities of a series of A1 and A2 adenosine receptor agonists and antagonists were determined using radioligand binding techniques in rat brain tissues. The potencies of these agonists on heart rate and coronary vascular tone were also assessed in the perfused working rat heart preparation. The order of potency of these agonists in producing neg. chronotropic effects was similar to the rank order for their A1 receptor binding activities [6-N-cyclohexyladenosine (CHA) = 6-N-(R-phenylisopropyl)adenosine > 5'-N-ethylcarboxamideadenosine (NECA) = 2-chloroadenosine > 2-phenylaminoadenosine] with a correlation coefficient of 0.97. Their order of potency in decreasing coronary vascular tone followed the same rank order as their A2 receptor binding activities with a correlation coefficient of 0.97 (NECA > 2-chloroadenosine = 6-N-(R-phenylisopropyl)-adenosine = 2-phenylaminoadenosine > CHA). In addition, the antagonists 8-[4-[[[(2-aminoethyl)amino]carbonyl]methyl]ox]phenyl-1,3-dipropylxanthine (XAC), 1,3-dipropyl-8-(2-amino-4-chlorophenyl)xanthine (PACPX), and 8-phenyltheophylline (8-PT) blocked the neg. chronotropic effect of CHA and the vasodilatory effect of NECA in a concentration-dependent manner. The same order of potency of the antagonists was noted in blocking CHA-induced bradycardia and A1 receptor binding activities (XAC = PACPX > 8-PT). A similar correlation was observed for their effects in blocking NECA-induced vasodilation and A2 receptor binding activity (XAC > PACPX > 8-PT). The results obtained with both agonists and antagonists indicate a pos. correlation between adenosine receptor-mediated effects in the heart and adenosine receptor binding activities in brain tissues; thus, providing support for similarities of these receptors in heart and brain tissues.

IT 53296-10-9, 2-Phenylaminoadenosine

RL: BIOL (Biological study)


(receptor binding of, in brain, coronary vascular tone and heart rate in relation to)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.

10/598,520



OSC.G 5 THERE ARE 5 CAPLUS RECORDS THAT CITE THIS RECORD (5 CITINGS)

L6 ANSWER 152 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1988:623029 CAPLUS

DN 109:223029

OREF 109:36749a,36752a

TI Characterization of agonist radioligand interactions with porcine atrial A1 adenosine receptors

AU Leid, Mark; Schimerlik, Michael I.; Murray, Thomas F.

CS Coll. Pharm., Oregon State Univ., Corvallis, OR, 97331, USA

SO Molecular Pharmacology (1988), 34(3), 334-9

CODEN: MOPMA3; ISSN: 0026-895X

DT Journal

LA English

AB The agonist radioligand (-)-N6-[125I]-p-hydroxyphenylisopropyladenosine ([125I]HPIA) was used to characterize adenosine recognition sites in porcine atrial membranes. [125I]HPIA showed saturable binding to an apparently homogeneous population of sites with a maximum binding capacity of 35 fmol/mg protein and an equilibrium dissociation constant of 2.5 nM. Kinetics expts. were performed to address the mol. mechanism of [125I]HPIA binding in porcine atrial membranes. [125I]HPIA apparently interacts with the cardiac adenosine receptor in a simple bimol. reaction. A kinetically derived [125I]HPIA dissociation constant (2.4 nM) was in good agreement with that parameter measured at equilibrium. Guanyl nucleotides neg. modulated [125I]HPIA binding by increasing its rate of dissociation. This finding is consonant with the formation of a ternary complex in porcine atrial membranes, consisting of ligand, receptor, and guanyl nucleotide-binding protein. Prototypic adenosine receptor agonists and antagonists inhibited specific binding in a manner consistent with the labeling of an A1 adenosine receptor. Apparently, the adenosine receptor present in porcine atrial membranes, as labeled by [125I]HPIA, is of the A1 subtype.

IT 53296-10-9, 2-Phenylaminoadenosine


RL: BIOL (Biological study)

(hydroxyphenylisopropyladenosine binding by receptors of atrium inhibition by)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

L6 ANSWER 153 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1988:611374 CAPLUS

DN 109:211374

OREF 109:34987a,34990a


TI Studies toward synthesis of C-2 substituted adenosines: an efficient synthesis of 2-(phenylamino)adenosine [CV-1808]

AU Trivedi, Bharat K.

McIntosh


10/598,520

CS Dep. Chem., Warner/Lambert Co., Ann Arbor, MI, 48105, USA  
SO Nucleosides & Nucleotides (1988), 7(3), 393-402  
CODEN: NUNUD5; ISSN: 0732-8311  
DT Journal  
LA English  
OS CASREACT 109:211374  
GI



AB 2-(Phenylamino)adenosine (I) was prepared from guanosine triacetate (II) by sequential 2-bromination with amyl nitrite and  $\text{CHBr}_3$ , 2-phenylamination with  $\text{PhNH}_2$ , 6-chlorination with  $\text{POCl}_3$  in MeCN in the presence of  $\text{PhNMe}_2$  and  $\text{Et}_4\text{NCl}$ , and finally treatment with  $\text{NH}_3/\text{MeOH}$ . Also prepared were (R)-N-(1-methyl-2-phenylethyl)-2-(phenylamino)adenosine, 2-(phenylthio)adenosine, and (R)-N-(1-methyl-2-phenylethyl)-2-(phenylthio)adenosine.  
IT 53296-10-9P, 2-(Phenylamino)adenosine  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(improved synthesis of)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 6 THERE ARE 6 CAPLUS RECORDS THAT CITE THIS RECORD (6 CITINGS)


L6 ANSWER 154 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1988:523020 CAPLUS  
DN 109:123020  
OREF 109:20355a,20358a  
TI Behavior induced by putative nociceptive neurotransmitters is inhibited by adenosine or adenosine analogs coadministered intrathecally  
AU DeLander, Gary E.; Wahl, Jeffrey J.  
CS Coll. Pharm., Oregon State Univ., Corvallis, OR, 97331, USA  
SO Journal of Pharmacology and Experimental Therapeutics (1988), 246(2), 565-70  
CODEN: JPETAB; ISSN: 0022-3565  
DT Journal  
LA English  
AB The role of adenosine in antinociception was studied by examining the effect of adenosine agonists on behavior induced by 2 putative spinal nociceptive neurotransmitters, substance P and N-methyl-D-aspartate. Coadministration of each of several adenosine agonists with substance P or N-methyl-D-aspartate intrathecally significantly decreased the intensity of behaviors induced by putative nociceptive neurotransmitters in mice. Adenosine agonist-mediated inhibition was antagonized by theophylline supporting adenosine agonist interactions with cell membrane surface

McIntosh

adenosine receptors. Rank order potencies were determined for several adenosine analogs with varying selectivity for A1 and A2 adenosine receptor subtypes. However, rank order potencies did not correlate well with rank order potencies reported previously for adenosine receptor subtypes in biochem. assays. Evidently, adenosine inhibits behavior induced by nociceptive neurotransmitters interacting with spinal substance P or N-methyl-D-aspartate receptors. Furthermore, observations provide addnl. support for endogenous antinociceptive pathways that utilize adenosine at spinal sites.


IT 53296-10-9, 2-Phenylaminoadenosine  
 RL: BIOL (Biological study)  
 (nociception inhibition by)  
 RN 53296-10-9 CAPLUS  
 CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 32 THERE ARE 32 CAPLUS RECORDS THAT CITE THIS RECORD (32 CITINGS)

L6 ANSWER 155 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
 AN 1988:438171 CAPLUS  
 DN 109:38171  
 OREF 109:6475a,6478a  
 TI N6-[2-(3,5-dimethoxyphenyl)-2-(2-methylphenyl)ethyl]adenosine and its uronamide derivatives. Novel adenosine agonists with both high affinity and high selectivity for the adenosine A2 receptor  
 AU Bridges, Alexander J.; Bruns, Robert F.; Ortwine, Daniel F.; Priebe, Steven R.; Szotek, Deedee L.; Trivedi, Bharat K.  
 CS Parke-Davis Pharm. Res. Div., Warner-Lambert Co., Ann Arbor, MI, 48105, USA  
 SO Journal of Medicinal Chemistry (1988), 31(7), 1282-5  
 CODEN: JMCMAR; ISSN: 0022-2623  
 DT Journal  
 LA English  
 OS CASREACT 109:38171  
 GI



AB Several N6-(diarylethyl)adenosines, e.g., title compound I, were prepared by treating the corresponding 2,2-diarylethylamines with 6-chloropurine

10/598,520

riboside. Also prepared were uronamide derivs. II (R = Et, Me, cyclopropyl). Receptor binding affinities of the compds. prepared and of several other N6-substituted adenosines are given and discussed.


IT 53296-10-9

RL: RCT (Reactant); RACT (Reactant or reagent)  
(adenosine agonist, receptor binding affinity of)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 26 THERE ARE 26 CAPLUS RECORDS THAT CITE THIS RECORD (27 CITINGS)

L6 ANSWER 156 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1988:105930 CAPLUS

DN 108:105930

OREF 108:17195a,17198a

TI Definition of subclasses of adenosine receptors associated with adenylate cyclase: interaction of adenosine analogs with inhibitory A1 receptors and stimulatory A2 receptors

AU Ukena, Dieter; Olsson, Ray A.; Daly, John W.


CS Natl. Inst. Arthritis, Diabetes, Dig. Kidney Dis., Natl. Inst. Health, Bethesda, MD, 20892, USA

SO Canadian Journal of Physiology and Pharmacology (1987), 65(3), 365-76  
CODEN: CJPPA3; ISSN: 0008-4212

DT Journal

LA English

GI




AB The structure-activity relationships of 63 adenosine analogs as agonists for the A1 adenosine receptors that mediate inhibition of adenylate cyclase activity in rat fat cells and for the A2 adenosine receptors that mediate stimulation of adenylate cyclase in rat pheochromocytoma PC12 cells and human platelets were determined. The lack of correspondence between the structure-activity relationships of these analogs at the A1 and A2 receptors appear definitive in terms of establishing the existence of A1 and A2 subclasses of adenosine receptors. However, significant differences in the agonist profiles at A2 receptors of platelet and PC12 indicate a certain degree of structural heterogeneity within the members of the A2 adenosine receptor subclass. Whether such differences are due to different species or different cell types is not known. A set of adenosine analogs, such as N6-cyclohexyl- (I), N6-R-, and N6-S-1-phenyl-2-propyladenosine, 5'-N-ethylcarboxamidoadenosine and its N6-cyclohexyl derivative, 2-chloroadenosine, and 2-phenylaminoadenosine,

McIntosh

10/598,520

appear to represent a series of analogs useful for pharmacol.  
characterization of A1 and A2 classes of adenosine receptors.  
IT 53296-10-9, 2-Phenylaminoadenosine  
RL: PRP (Properties)  
(interaction of, with A1 and A2 adenosine receptors, of humans and laboratory  
animals, structure in relation to)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)


Absolute stereochemistry.



OSC.G 13 THERE ARE 13 CAPLUS RECORDS THAT CITE THIS RECORD (13 CITINGS)

L6 ANSWER 157 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1988:16690 CAPLUS  
DN 108:16690  
OREF 108:2729a,2732a  
TI Correlation of adenosine receptor affinities and cardiovascular activity  
AU Hamilton, H. W.; Taylor, M. D.; Steffen, R. P.; Haleen, S. J.; Bruns, R.  
F.  
CS Dep. Chem., Warner-Lambert/Parke-Davis Pharm. Res., Ann Arbor, MI, 48105,  
USA  
SO Life Sciences (1987), 41(20), 2295-302  
CODEN: LIFSAR; ISSN: 0024-3205  
DT Journal  
LA English  
AB Binding affinities of 28 adenosine analogs at A1 adenosine receptors [rat  
whole brain membranes, [3H]N6-cyclohexyladenosine (CHA)], and at A2  
adenosine receptors [rat striatal membranes,  
5'-N-ethylcarboxamidoadenosine (NECA) were compared to their EC25 (25%  
change from control) values for decreasing heart rate and increasing  
coronary flow in the isolated rat heart. Heart rate (an A1 response)  
correlated with A1 binding affinity but not with A2 binding affinity;  
conversely, coronary flow (an A2 response) correlated with A2 binding  
affinity but not with A1 binding affinity. Apparently, the brain A1 and  
A2 receptors, studied by binding methods, bear close similarities to their  
resp. counterparts in the heart, studied by means of functional responses.  
IT 53296-10-9  
RL: BIOL (Biological study)  
(adenosine receptor affinity for, in brain, cardiovascular activity in  
relation to)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 8 THERE ARE 8 CAPLUS RECORDS THAT CITE THIS RECORD (8 CITINGS)


L6 ANSWER 158 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

McIntosh

10/598,520

AN 1987:612266 CAPLUS  
DN 107:212266  
OREF 107:33919a,33922a  
TI Identification of A1 and A2 adenosine receptors in the rat spinal cord  
AU Choca, Jose Ignacio; Proudfit, Herbert K.; Green, Richard D.  
CS Coll. Med., Univ. Illinois, Chicago, IL, 60680, USA  
SO Journal of Pharmacology and Experimental Therapeutics (1987), 242(3),  
905-10  
CODEN: JPETAB; ISSN: 0022-3565  
DT Journal  
LA English  
AB The adenosine receptors in membranes prepared from rat ventral and dorsal  
lumbar spinal cord were characterized by comparing the binding  
characteristics of [3H]5'-N-ethylcarboxamide adenosine ([3H]NECA), an  
agonist with nearly equal affinities at the A1 and A2 adenosine receptor  
subtypes, with those of [3H]N6-[(R)-1-methyl-2-phenylethyl]adenosine  
([3H]R-PIA), an A1-selective agonist. Saturation isotherms of the ventral and  
dorsal spinal cord yielded dissociation constant values 1.9-2.3 nM for [3H]R-PIA  
and 18.1-19.5 nM for [3H]NECA. The maximum binding capacity (Bmax) for  
[3H]NECA was approx. twice the Bmax for [3H]R-PIA in ventral and dorsal  
halves (267 vs. 128 fmol/mg protein and 402 vs. 206 fmol/mg protein,  
resp.). Displacement of specific [3H]NECA binding by the A2-selective  
agonist, 2-(phenylamino)adenosine, the relatively nonselective antagonist,  
theophylline and 6 A1-selective agonists, R-PIA, S-PIA,  
N6-(cyclohexyl)adenosine, N6-(cyclopentyl)adenosine,  
N6-(m-aminophenyl)adenosine, and N6-(m-iodophenyl)adenosine, revealed 2  
[3H]NECA binding components with the characteristics of A1 and A2  
receptors. All curves best fit a 2-site model when analyzed by the  
computer program LIGAND. R-PIA, N6-(cyclohexyl)adenosine, and  
N6-(cyclopentyl)adenosine were the most potent displacers at the 1st site  
( $K_i = 0.6-1.4$  nM). All A1-selective agonists were poor displacers of  
[3H]NECA at the 2nd site ( $K_i = 0.6-18.6$   $\mu$ M). The A2-selective agonist,  
2-(phenylamino)adenosine, was as potent as R-PIA in displacing [3H]NECA  
from this site with a  $K_i$  value 0.57  $\mu$ M. Finally, the A1 and A2  
adenosine receptor-mediated inhibition and stimulation of adenylate  
cyclase were demonstrated directly in synaptic membranes prepared from the  
spinal cord.  
IT 53296-10-9, 2-(Phenylamino)adenosine  
RL: PROC (Process)  
(receptor binding of, in spinal cord)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 59 THERE ARE 59 CAPLUS RECORDS THAT CITE THIS RECORD (59 CITINGS)

L6 ANSWER 159 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1987:508844 CAPLUS  
DN 107:108844  
OREF 107:17515a,17518a  
TI The effects of parenteral injections of adenosine and its analogs on blood  
pressure and heart rate in the rat  
AU Barraco, Robin A.; Marcantonio, David R.; Phillis, John W.; Campbell, W.  
Richard  
CS Sch. Med., Wayne State Univ., Detroit, MI, 48201, USA  
SO General Pharmacology (1987), 18(4), 405-16  
CODEN: GEPHDP; ISSN: 0306-3623  
DT Journal  
LA English  
AB The dose-response effects of i.v. adenosine and its analogs on

McIntosh



cardiovascular parameters were examined in rats.

5'-N-Ethylcarboxamido-adenosine (NECA) was by far the most potent analog in reducing mean arterial blood (PA) pressure, whereas N6-(3-pentyl)-adenosine exerted the most potent bradycardic action. The N6-substituted (S-)-diastereoisomers were substantially less potent in reducing PA and heart rate than NECA and the N6-substituted (R)-diastereoisomers. The results of the study are consistent with the notion that the bradycardiac action of adenosine is principally mediated via A1 receptors, whereas the vasodilator action of adenosine is mediated via A2 receptors. It is also apparent that adenosine is rapidly removed from the circulation and inactivated. In contrast, the cardiovascular effects of the adenosine analogs persist, to varying degrees, much longer than those of adenosine itself.

IT 53296-10-9, 2-Phenylamino-adenosine


RL: BIOL (Biological study)

(blood pressure and heart rate response to, structure in relation to)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 4 THERE ARE 4 CAPLUS RECORDS THAT CITE THIS RECORD (4 CITINGS)

L6 ANSWER 160 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1987:497042 CAPLUS

DN 107:97042

OREF 107:15844h,15845a

TI Synthesis of a mutagenic nucleoside, 2'-deoxy-2-(p-nitrophenyl)adenosine

AU Matsuda, Akira; Ueda, Tohru; Ohara, Yoshiko; Satake, Hiroyasu; Negishi, Kazuo; Wataya, Yusuke; Hayatsu, Hikoya

CS Fac. Pharm. Sci., Hokkaido Univ., Sapporo, 060, Japan

SO Nucleic Acids Symposium Series (1986), 17(Symp. Nucleic Acids Chem., 14th, 1986), 141-3


CODEN: NACSD8; ISSN: 0261-3166

DT Journal

LA English

OS CASREACT 107:97042

GI



AB The reaction of 2-amino-6-chloropurine riboside with isoamyl nitrite in C6H6 in the presence of Cu2O, followed by treatment with NH3/MeOH, gave 2-phenyladenosine I (R = H, R1 = OH) (II). The crude sample of II was found to be mutagenic to bacteria (Salmonella typhimurium TA 98 and TA 100, without metabolic activation). When this material was subjected to

10/598,520

high pressure liquid chromatog., the mutagenic activity was found only in contaminating minor components, whose structures were assigned as 2-(m- and p-nitrophenyl)adenosines (I; R = m-, p-NO<sub>2</sub>; R<sub>1</sub> = OH). In order to study structure-activity relationships, several nucleoside and base analogs were prepared. Among them, 2'-deoxy-2-(p-nitrophenyl)adenosine (I; R = p-O<sub>2</sub>N, R<sub>1</sub> = H) was the most potent mutagen as tested either with TA 98 or TA 100.

IT 109875-49-2P


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)

(preparation and mutagenic activity of)

RN 109875-49-2 CAPLUS

CN Adenosine, 2-(4-aminophenyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L6 ANSWER 161 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1987:78255 CAPLUS

DN 106:78255

OREF 106:12705a,12708a

TI Species differences in structure-activity relationships of adenosine agonists and xanthine antagonists at brain A<sub>1</sub> adenosine receptors

AU Ukena, Dieter; Jacobson, Kenneth A.; Padgett, William L.; Ayala, Cristina; Shamim, Mah T.; Kirk, Kenneth L.; Olsson, Ray O.; Daly, John W.

CS Natl. Inst. Diabetes, and Dig. Kidney Dis., Natl. Inst. Health, Bethesda, MD, 20892, USA

SO FEBS Letters (1986), 209(1), 122-8

CODEN: FEBLAL; ISSN: 0014-5793

DT Journal

LA English

AB A series of 28 adenosine analogs and 17 xanthines were assessed as inhibitors of N<sub>6</sub>-[<sup>3</sup>H]phenylisopropyladenosine binding to A<sub>1</sub> adenosine receptors in membranes from rat, calf, and guinea pig brain. Potencies of N<sub>6</sub>-alkyl- and N<sub>6</sub>-cycloalkyladenosines are similar in the different species. However, the presence of an aryl or heteroaryl moiety in the N<sub>6</sub> substituent results in marked species differences with certain such analogs being about 30-fold more potent at receptors in calf than in guinea pig brain. Potencies at receptors in rat brain are intermediate. Conversely, 2-chloroadenosine [146-77-0] and 5'-N-ethylcarboxamidoadenosine [35920-39-9] are .apprx.10-fold less potent at receptors in calf brain than in guinea pig brain. Potencies of xanthines, such as theophylline [58-55-9], caffeine [58-08-2] and 1,3-dipropylxanthine [31542-62-8] are similar in the different species. However, the presence of an 8-Ph or 8-cycloalkyl substituent results in marked species differences. For example, a xanthine amine conjugate of 1,3-dipropyl-8-phenylxanthine is 9-fold more potent at receptors in calf than in rat brain and 110-fold more potent in calf than in guinea pig brain. Such differences indicate that brain A<sub>1</sub> adenosine receptors are not identical in recognition sites for either agonists or antagonists in different mammalian species.

IT 53296-10-9

RL: BIOL (Biological study)

(adenosine A<sub>1</sub> receptor binding by, structure and species in relation to)


RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.

McIntosh


10/598,520



OSC.G 22 THERE ARE 22 CAPLUS RECORDS THAT CITE THIS RECORD (22 CITINGS)

L6 ANSWER 162 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1987:16911 CAPLUS  
DN 106:16911  
OREF 106:2905a,2908a  
TI The effects of adenosine agonists on human neutrophil function  
AU Schrier, Denis J.; Imre, Kathleen M.  
CS Warner-Lambert/Parke-Davis Pharm. Res., Ann Arbor, MI, 48105, USA  
SO Journal of Immunology (1986), 137(10), 3284-9  
CODEN: JOIMA3; ISSN: 0022-1767  
DT Journal  
LA English  
AB Adenosine is a potent physiol. substance with a variety of biol. activities. Many of the effects of adenosine appear to be mediated by 2 populations of cell-surface adenosine receptors (A1 and A2). The effects were examined of several adenosine receptor agonists on human neutrophils stimulated with the chemotactic peptide N-formyl-Met-Leu-Phe (FMLP). Both superoxide generation and degranulation (as assessed by lysozyme release) were inhibited. Inhibition correlated most strongly with A2 receptor affinity for both parameters and was reversible by the adenosine receptor antagonist 8-phenyltheophylline. Because toxic O metabolites and degradative enzymes are implicated in a variety of inflammatory disorders, adenosine agonists may be useful probes to help expand knowledge of the role of these mediators in human disease.  
IT 53296-10-9, 2-(Phenylamino)-adenosine  
RL: BIOL (Biological study)  
(neutrophil function response to, purinergic receptors in, of human)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 33 THERE ARE 33 CAPLUS RECORDS THAT CITE THIS RECORD (33 CITINGS)

L6 ANSWER 163 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1987:13071 CAPLUS  
DN 106:13071  
OREF 106:2157a,2160a  
TI Adenosine analogs mediating depressant effects on synaptic transmission in rat hippocampus: structure-activity relationships for the N6 subregion  
AU Dunwiddie, Thomas V.; Worth, Thomas S.; Olsson, Ray A.  
CS Veterans Adm. Med. Res. Serv., Denver, CO, 80220, USA  
SO Naunyn-Schmiedeberg's Archives of Pharmacology (1986), 334(1), 77-85  
CODEN: NSAPCC; ISSN: 0028-1298  
DT Journal  
LA English  
AB The potencies of a number of N6-substituted adenosine analogs in depressing excitatory synaptic transmission were investigated in slices of rat

McIntosh

hippocampus, an electrophysiol. response mediated by receptors of the A1 subtype. These potencies correlated well with previously reported affinities of these analogs for A1 receptor sites in brain, but not with coronary vasodilation in the dog heart, an A2 receptor-mediated response. Analogs with alkyl or aryl substituents at the N6 position were generally more potent than adenosine [41552-82-3], although analogs with a tertiary C attached directly to the N6-N were usually only weakly active. Although it has been suggested that there may be a subregion of the A1 receptor with some specificity for aryl groups, these expts. did not suggest that this was the case. Analogs with chiral centers attached to the N6-N usually displayed stereoselectivity, with R-isomers more potent than the S-isomers. The mechanism underlying this selectivity appeared to be both a facilitating effect of alkyl substituents in the Pr C1 position of N6-1-phenyl-2(R)-propyladenosine (R-PIA) [38594-96-6], and a hindering effect of substituents in the position normally occupied by the H attached to Pr C2 of R-PIA. Although there are some similarities in terms of requirements for activity at A1 and A2 receptors, differences between the N6 subregions of these receptors are sufficient to permit the development of selective analogs for these 2 receptor sites.

IT 53296-10-9


RL: BIOL (Biological study)

(synaptic neurotransmission in hippocampus inhibition by, structure in relation to)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 6 THERE ARE 6 CAPLUS RECORDS THAT CITE THIS RECORD (6 CITINGS)

L6 ANSWER 164 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1986:526814 CAPLUS

DN 105:126814

OREF 105:20297a,20300a

TI Synthetic studies of 2-substituted adenosines. III. Coronary vasodilatory activity of 2-arylaminoadenosines

AU Marumoto, Ryuji; Shima, Shunsuke; Omura, Kiyoshi; Tanabe, Masao; Fujiwara, Syuji; Furukawa, Yoshiyasu

CS Cent. Res. Div., Takeda Chem. Ind., Ltd., Osaka, 532, Japan


SO Takeda Kenkyushoho (1985), 44(3/4), 220-30

CODEN: TAKHAA; ISSN: 0371-5167

DT Journal

LA English

GI



AB Sixty-one derivs. of 2-phenylaminoadenosine (CV-1808) (I) were synthesized and their coronary vasodilatory activities were tested after intracoronary administration in anesthetized dogs. Introduction of a variety of substituents into the 4-position of the Ph group led to a considerable decrease in the activity; substitution at the 3-position did not alter the potency, whereas substitution at the 3- or 4-position with a carbamoyl or acyl group increased the activity approx. 10 times. Replacement of the Ph group in I derivs. by a 3-pyridyl ring also resulted in an increase in the activity. Structure-activity relations are discussed.

IT 53296-10-9DP, derivs. 53296-10-9P  
 53296-19-8P 53296-20-1P 53296-21-2P  
 70590-18-0P 70590-20-4P 70590-22-6P  
 70590-23-7P 70590-25-9P 70590-26-0P  
 70590-27-1P 70590-28-2P 70590-29-3P  
 70590-30-6P 71231-76-0P 71231-77-1P  
 71231-78-2P 71231-79-3P 71231-80-6P  
 71231-81-7P 71231-82-8P 71231-83-9P  
 71231-84-0P 71231-85-1P 71231-86-2P  
 74615-32-0P 74615-33-1P 74615-36-4P  
 74615-37-5P 74615-38-6P 74615-39-7P  
 74615-40-0P 74615-41-1P 74615-42-2P  
 75106-29-5P 75106-30-8P 75106-32-0P  
 75106-33-1P 76888-18-1P 102711-68-2P  
 102711-69-3P 102711-70-6P 102711-71-7P  
 102711-72-8P 102711-87-5P 102711-88-6P  
 102711-89-7P 102711-90-0P 102711-91-1P  
 102711-92-2P 102711-93-3P 102711-94-4P  
 102711-95-5P 102711-96-6P 102711-97-7P  
 102711-98-8P 102711-99-9P 102712-00-5P  
 102712-01-6P 102712-02-7P


RL: SPN (Synthetic preparation); PREP (Preparation)

(preparation and vasodilator activity of, structure in relation to)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)


Absolute stereochemistry.



RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.




RN 53296-19-8 CAPLUS

CN Adenosine, 2-[(4-methoxyphenyl)amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

10/598,520




RN 53296-20-1 CAPLUS  
CN Adenosine, 2-[(4-methylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 53296-21-2 CAPLUS  
CN Adenosine, 2-[(4-chlorophenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 70590-18-0 CAPLUS  
CN Adenosine, 2-[(3-chlorophenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 70590-20-4 CAPLUS  
CN Adenosine, 2-[(3,4-dimethylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 70590-22-6 CAPLUS  
CN Adenosine, 2-[(3-methoxyphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 70590-23-7 CAPLUS  
CN Adenosine, 2-[(3-methylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 70590-25-9 CAPLUS  
CN Adenosine, 2-[(3-fluorophenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 70590-26-0 CAPLUS  
CN Adenosine, 2-[(3-bromophenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

McIntosh


10/598,520



RN 70590-27-1 CAPLUS

CN Adenosine, 2-[(3-chloro-4-methylphenyl)amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 70590-28-2 CAPLUS

CN Adenosine, 2-[(3,4-dimethoxyphenyl)amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 70590-29-3 CAPLUS

CN Adenosine, 2-[(3-chloro-4-methoxyphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 70590-30-6 CAPLUS


CN Adenosine, 2-[(3,4-dichlorophenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

McIntosh




10/598,520




RN 71231-76-0 CAPLUS  
CN Adenosine, 2-[(2-methylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 71231-77-1 CAPLUS  
CN Adenosine, 2-[(2-methoxyphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 71231-78-2 CAPLUS  
CN Adenosine, 2-[(4-bromophenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 71231-79-3 CAPLUS  
CN Adenosine, 2-[(4-fluorophenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 71231-80-6 CAPLUS  
CN Adenosine, 2-[(2-fluorophenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 71231-81-7 CAPLUS  
CN Adenosine, 2-[(4-ethylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 71231-82-8 CAPLUS  
CN Adenosine, 2-[(2,5-dimethylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 71231-83-9 CAPLUS  
CN Adenosine, 2-[(3,5-dimethylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

McIntosh


10/598,520



RN 71231-84-0 CAPLUS

CN Adenosine, 2-[(2,4-dimethylphenyl)amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 71231-85-1 CAPLUS

CN Adenosine, 2-[(4-chloro-2-methoxyphenyl)amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 71231-86-2 CAPLUS

CN Adenosine, 2-[(2,3-dichlorophenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 74615-32-0 CAPLUS

CN Adenosine, 2-[[4-(aminocarbonyl)phenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 74615-33-1 CAPLUS  
CN Adenosine, 2-[[4-[(propylamino)carbonyl]phenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 74615-36-4 CAPLUS  
CN Adenosine, 2-[[3-(aminocarbonyl)phenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 74615-37-5 CAPLUS  
CN Adenosine, 2-[[4-(aminocarbonyl)-3-chlorophenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 74615-38-6 CAPLUS  
CN Adenosine, 2-[[4-[(methylamino)carbonyl]phenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 74615-39-7 CAPLUS  
CN Adenosine, 2-[[4-[(ethylamino)carbonyl]phenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 74615-40-0 CAPLUS  
CN Adenosine, 2-[[4-[(cyclohexylamino)carbonyl]phenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 74615-41-1 CAPLUS  
CN Adenosine, 2-[[4-[(phenylamino)carbonyl]phenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




McIntosh

10/598,520

RN 74615-42-2 CAPLUS

CN Adenosine, 2-[[4-[(dimethylamino)carbonyl]phenyl]amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 75106-29-5 CAPLUS

CN Adenosine, 2-[(4-acetylphenyl)amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 75106-30-8 CAPLUS

CN Adenosine, 2-[(3-acetyl-4-ethoxyphenyl)amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 75106-32-0 CAPLUS

CN Adenosine, 2-[(4-acetyl-3-methoxyphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 75106-33-1 CAPLUS

CN Adenosine, 2-[(4-acetyl-3-chlorophenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

McIntosh


10/598,520



RN 76888-18-1 CAPLUS

CN Adenosine, 2-[(3-acetylphenyl)amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 102711-68-2 CAPLUS

CN Adenosine, 2-[(4-acetyl-2,5-dimethylphenyl)amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 102711-69-3 CAPLUS

CN Adenosine, 2-[(4-acetyl-3,5-dimethylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 102711-70-6 CAPLUS

McIntosh

10/598,520

CN Adenosine, 2-[(5-acetyl-2,4-dimethylphenyl)amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 102711-71-7 CAPLUS

CN Adenosine, 2-[(5,6,7,8-tetrahydro-5-oxo-2-naphthalenyl)amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 102711-72-8 CAPLUS

CN Adenosine, 2-[(1,3-dihydro-3-oxo-5-isobenzofuranyl)amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 102711-87-5 CAPLUS

CN Adenosine, 2-[(3,5-dichlorophenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



McIntosh




10/598,520

RN 102711-88-6 CAPLUS

CN Adenosine, 2-[(4-acetyl-3-methylphenyl)amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 102711-89-7 CAPLUS

CN Adenosine, 2-[(5-acetyl-2-methylphenyl)amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 102711-90-0 CAPLUS

CN Adenosine, 2-[(5-acetyl-2-chlorophenyl)amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 102711-91-1 CAPLUS

CN Adenosine, 2-[(4-cyanophenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 102711-92-2 CAPLUS

CN Adenosine, 2-[[4-(dimethylamino)phenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 102711-93-3 CAPLUS  
CN Adenosine, 2-[[4-(1-piperidinylcarbonyl)phenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 102711-94-4 CAPLUS  
CN Adenosine, 2-[[4-(4-morpholinylcarbonyl)phenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 102711-95-5 CAPLUS  
CN Benzoic acid, 4-[(6-amino-9-β-D-ribofuranosyl-9H-purin-2-yl)amino]-, ethyl ester (CA INDEX NAME)

Absolute stereochemistry.




McIntosh

10/598,520

RN 102711-96-6 CAPLUS

CN Adenosine, 2-[[4-(aminosulfonyl)phenyl]amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 102711-97-7 CAPLUS

CN Adenosine, 2-[[3-(trifluoromethyl)phenyl]amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 102711-98-8 CAPLUS

CN Adenosine, 2-[(3-nitrophenyl)amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 102711-99-9 CAPLUS

CN Adenosine, 2-[(3-aminophenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 102712-00-5 CAPLUS

CN Adenosine, 2-[[3-(1-oxopropyl)phenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 102712-01-6 CAPLUS  
CN Adenosine, 2-[[3-(1-oxobutyl)phenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 102712-02-7 CAPLUS  
CN Adenosine, 2-[(2,3-dimethylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)


L6 ANSWER 165 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1986:491834 CAPLUS  
DN 105:91834  
OREF 105:14729a,14732a  
TI Towards selective adenosine antagonists  
AU Bruns, R. F.; Lu, G. H.; Pugsley, T. A.  
CS Dep. Pharmacol., Warner-Lambert/Parke-Davis Pharm. Res., Ann Arbor, MI, 48105, USA  
SO Adenosine: Recept. Modulation Cell Funct., Proc. Int. Workshop Adenosine Xanthine Deriv. (1985), Meeting Date 1984, 51-8. Editor(s): Stefanovich, V.; Rudolphi, K.; Schubert, P. Publisher: IRL, Oxford, UK.  
CODEN: 55CNAD  
DT Conference  
LA English  
AB Affinities of adenosine antagonists for the A1 and A2 subtypes of adenosine receptors were determined: A1 affinities from 3H-labeled N6-cyclohexyladenosine [36396-99-3] binding to membranes from rat whole brain, and A2 affinities from 3H-labeled 1-(6-amino-9H-purin-9-yl)-1-deoxy-N-ethyl-β-D-ribofuronamide [35920-39-9] binding to rat striatal membranes in the presence of 50 nM N6-cyclopentyladenosine [41552-82-3]. The compds. were also tested for water solubility and for inhibition of the 3 forms of cytosolic phosphodiesterase from guinea pig heart. Most of the common xanthines were 3-10-fold selective for A1 receptors. 8-Cyclopentyltheophylline [35873-49-5] had 100-fold selectivity for A1-receptors and reasonable

McIntosh


water solubility Alloxazine [490-59-5] had 2-fold selectivity for A2 receptors but was not very soluble relative to its adenosine receptor median inhibitory concentration (IC50). PD 113,297 [96445-35-1], a xanthine derivative containing a tertiary amine, was a potent adenosine antagonist (A1 IC50 8 nM, A2 IC50, 100 nM) with good water solubility. The above antagonists produced negligible phosphodiesterase inhibition even at concns. which completely occupied adenosine receptors.

IT 53296-10-9  
 RL: PRP (Properties)  
 (adenosine receptor subtype affinity of)  
 RN 53296-10-9 CAPLUS  
 CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



L6 ANSWER 166 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
 AN 1986:419132 CAPLUS  
 DN 105:19132  
 OREF 105:3097a,3100a  
 TI Characterization of the A2 adenosine receptor labeled by [3H]NECA in rat striatal membranes  
 AU Bruns, Robert F.; Lu, Gina H.; Pugsley, Thomas A.  
 CS Dep. Pharmacol., Warner-Lambert/Parke-Davis Pharm. Res., Ann Arbor, MI, 48105, USA  
 SO Molecular Pharmacology (1986), 29(4), 331-46  
 CODEN: MOPMA3; ISSN: 0026-895X  
 DT Journal  
 LA English  
 GI



AB To study the putative A2 component of 3H-labeled NECA (I) [35920-39-9] binding, several compds. were examined for the ability to selectively eliminate the A1 component of binding in rat striatal membranes; N6-cyclopentyladenosine [41552-82-3] gave the most satisfactory results. Binding of [3H]NECA in the presence of 50 nM N6-cyclopentyladenosine was characterized. The rank order of potency for inhibition of [3H]NECA binding was NECA » 2-chloroadenosine [146-77-0] > N6-[(R)-1-methyl-2-phenylethyl]adenosine (R-PIA) [38594-96-6] > N6-cyclohexyladenosine [36396-99-3] > S-PIA [38594-97-7], indicating that binding was to an A2 adenosine receptor. When affinities of compds. in [3H]NECA binding to A2 receptors were compared to their affinities in [3H]N6-cyclohexyladenosine binding to A1 receptors, N6-cyclopentyladenosine was the most A1-sensitive agonist (A1 inhibition constant (Ki), 0.59 nM; A2Ki, 460 nM; Ki ratio, 780), whereas the selective

10/598,520

coronary vasodilator 2-(phenylamino)adenosine [53296-10-9] was the most A2-selective agonist (A1, 560 nM; A2, 120 nM; ratio, 0.21). The antagonist 8-cyclopentyltheophylline had considerable A1 selectivity (A1, 11 nM; A2, 1400 nM; ratio, 130), whereas alloxazine had slight A2 selectivity (A1, 5200 nM; A2, 2700; ratio, 0.52). [3H]NECA binding to A2 receptors was highest in striatum but was detectable at much lower levels in each of 7 other brain areas. The regional distribution of [3H]NECA binding and the affinities of adenosine agonists and antagonists for inhibition of binding indicate that the site labeled by [3H]NECA belongs to the high-affinity, or A2a, subclass of A2 receptor.

IT 53296-10-9


RL: BIOL (Biological study)

(purinergic A1 and A2 receptors binding of, in brain membranes, structure in relation to)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 274 THERE ARE 274 CAPLUS RECORDS THAT CITE THIS RECORD (274 CITINGS)

L6 ANSWER 167 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1986:142664 CAPLUS

DN 104:142664

OREF 104:22415a,22418a

TI Behavioral characteristics of centrally administered adenosine analogs

AU Phillis, J. W.; Barraco, R. A.; DeLong, R. E.; Washington, D. O.

CS Sch. Med., Wayne State Univ., Detroit, MI, 48201, USA

SO Pharmacology, Biochemistry and Behavior (1986), 24(2), 263-70

CODEN: PBBHAU; ISSN: 0091-3057

DT Journal

LA English

AB A series of adenosine analogs and related compds. were injected into the lateral cerebral ventricle (ICVT) and their effects on spontaneous locomotor activity of mice recorded. All analogs produced dose-related decreases in locomotor activity 5'-N6-ethyl-carboxamidoadenosine (NECA) [35920-39-9] was the most potent compound tested, with a number of N6-substituted analogs also being effective depressants of activity. Caffeine, administered either ICVT or i.p., antagonized the depressant effects of the adenosine analogs. IBMX, administered ICVT, depressed locomotor activity. However, after caffeine, IBMX elicited behavioral stimulation. Agents which inhibit the transport of adenosine [58-61-7] (dipyridamole, dilazep, papaverine) depressed locomotor activity, as did erythro-9-(2-hydroxy-3-nonyl)adenine (EHNA), an inhibitor of adenosine deaminase. The effects of dilazep, papaverine, and EHNA, but not of dipyridamole, were antagonized by caffeine. Endogenous adenosine is apparently involved in the regulation of central nervous system excitability.

IT 53296-10-9

RL: BIOL (Biological study)


(behavior response to intracerebroventricular administration of)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.

10/598,520



OSC.G 10 THERE ARE 10 CAPLUS RECORDS THAT CITE THIS RECORD (10 CITINGS)

L6 ANSWER 168 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1984:465868 CAPLUS

DN 101:65868

OREF 101:10039a,10042a

TI Further studies on the inhibition of adenosine uptake into rat brain synaptosomes by adenosine derivatives and methylxanthines

AU Wu, P. H.; Barraco, R. A.; Phillis, J. W.

CS Sch. Med., Wayne State Univ., Detroit, MI, 48201, USA

SO General Pharmacology (1984), 15(3), 251-4

CODEN: GEPHDP; ISSN: 0306-3623

DT Journal

LA English

AB Various compds. were tested for their abilities to inhibit the rapid uptake of adenosine [58-61-7] by rat cerebral cortical synaptosomes. Several pharmacol. potent derivs. of adenosine were weak inhibitors of uptake, with 20% inhibitory concns. (IC20) >10-5M. Derivs. in this category were adenosine-5'-N-ethylcarboxamide [74992-42-0], adenosine-5'-cyclopropylcarboxamide [50908-62-8], N6-cyclohexyladenosine [36396-99-3], L-N6-phenylisopropyladenosine [38594-97-7], 1-methylisoguanosine [70639-65-5], 2-phenylaminoadenosine [53296-10-9], and 5-iodotubericidin [91284-08-1]. Several methylxanthines were very weak inhibitors of adenosine uptake. These included pentoxifylline [6493-05-6], hexyltheophylline [1028-36-0], butyltheobromine [1143-30-2], and isoamyltheobromine [1024-65-3]. HL 725 [78416-81-6], a pyrimidoisoquinoline with potent phosphodiesterase-inhibitory activity, inhibited adenosine uptake with an IC20 of 2.0 + 10-6M. PK 11195 [85532-75-8], a putative ligand for the peripheral benzodiazepine binding site, did not alter uptake at 10-4M.

IT 53296-10-9


RL: BIOL (Biological study)

(adenosine uptake by brain synaptosome inhibition by)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 7 THERE ARE 7 CAPLUS RECORDS THAT CITE THIS RECORD (7 CITINGS)

L6 ANSWER 169 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1984:448402 CAPLUS

DN 101:48402

OREF 101:7403a,7406a

TI Inhibition of coronary circulatory failure and thromboxane A2 release during coronary occlusion and reperfusion by 2-phenylaminoadenosine (CV-1808) in anesthetized dogs


AU Tanabe, M.; Terashita, Z.; Nishikawa, K.; Hirata, M.

CS Cent. Res. Div., Takeda Chem. Ind. Ltd., Osaka, Japan

McIntosh

10/598,520


SO Journal of Cardiovascular Pharmacology (1984), 6(3), 442-8  
CODEN: JPCPDT; ISSN: 0160-2446  
DT Journal  
LA English  
GI



AB The effects of a potent coronary vasodilator, CV-1808 (I) [53296-10-9], on coronary circulatory failure and thromboxane (TX) A2 [66719-58-2] release were studied during coronary occlusion (for 60 min) and subsequent reperfusion (for 60 min) in anesthetized dogs. During coronary reperfusion, the reactive hyperemic response was attenuated, and coronary conductance decreased gradually with time, suggesting coronary circulatory failure. TXA2 release was markedly increased, as demonstrated by contraction of rabbit aortic strips perfused with coronary venous blood draining the ischemic myocardium, and by increased release of radioimmunol. assayable TXB2. CV-1808 (0.25 µg/kg/min i.v. infusion throughout the exptl. period, starting 10 min before coronary occlusion) inhibited coronary circulatory failure and TXA2 release. TXA2 synthetase [60832-04-4] of horse platelet microsomes was not significantly inhibited (-11.6%) by 10-4M CV-1808. The compound (10-5 and 10-4M) inhibited collagen-induced TXB2 [54397-85-2] formation in a dose-dependent manner (-23.0 and -74.0%, resp.), but not arachidonic acid-induced TXB2 formation by dog platelets, suggesting that CV-1808 inhibited phospholipases. Myocardial infarct size determined 60 min after reperfusion was significantly reduced by CV-1808. Thus, CV-1808 appeared to be effective for salvaging ischemic myocardium. The effect might be related to improvement of coronary circulation and inhibition of release of vasoactive substances, including TXA2, from the ischemic myocardium.

IT 53296-10-9  
RL: BIOL (Biological study)  
(heart ischemia response to)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD (3 CITINGS)


L6 ANSWER 170 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1983:172895 CAPLUS  
DN 98:172895  
OREF 98:26089a,26092a  
TI Potentiation of the negative chronotropic and inotropic effects of adenosine by 2-phenylaminoadenosine  
AU Chiba, Shigetoshi; Watanabe, Hidehiko  
CS Sch. Med., Shinshu Univ., Matsumoto, 390, Japan

McIntosh




10/598,520

SO Clinical and Experimental Pharmacology and Physiology (1983), 10(1), 1-5  
CODEN: CEXPB9; ISSN: 0305-1870  
DT Journal  
LA English  
GI




AB The effects of 2-phenylaminoadenosine (I) [53296-10-9] on sinoatrial nodal pacemaker activity and atrial contractility were studied in isolated, blood-perfused dog atrial prepns. The compds. were administered via the cannulated sinus node artery of the isolated atrium. I caused neg. chronotropic and inotropic effects. The compound was 100 times less potent than adenosine [58-61-7]. I potentiated the effect of adenosine on atrial muscle, but not that of acetylcholine.  
IT 53296-10-9  
RL: BIOL (Biological study)  
(heart response to adenosine in relation to)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



L6 ANSWER 171 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1983:172389 CAPLUS  
DN 98:172389  
OREF 98:25961a,25964a  
TI Quantitation of 6-amino-2-phenylamino-9- $\beta$ -D-ribofuranosyl-9H-purine (CV-1808) and its metabolite, 2-(4-hydroxyphenyl)aminoadenosine, in human serum and urine by high-performance liquid chromatography using a fluorimetric detector  
AU Hayashi, Yoshitatsu; Miyake, Sohachiro; Kuwayama, Motoaki; Hattori, Masatoshi; Usui, Yoshiro  
CS Cent. Res. Div., Takeda Chem. Ind., Ltd., Osaka, 532, Japan  
SO Chemical & Pharmaceutical Bulletin (1982), 30(11), 4107-13  
CODEN: CPBTAL; ISSN: 0009-2363  
DT Journal  
LA English  
GI

10/598,520



I, R=H


II, R=OH

AB A high performance liquid chromatog. method using a fluorimetric detector for determination of the quantities of CV-1808 (I) [53296-10-9] and its metabolite 2-(4-hydroxyphenyl)aminoadenosine (II) [81613-39-0] in human serum and urine is presented. I and II, after chromatog. extraction from urine or serum with a Sep-Pak C18 cartridge, are allowed to react with propionic anhydride in the presence of triethylamine and the quantities of the resulting propionyl derivs. of I and II (I-P and II-P) are determined by high performance liquid chromatog. on a  $\mu$ Porasil column. The detection limits of I and II are 5.0 and 10.0 ng/mL in urine and 1.0 and 2.0 ng/mL in serum, resp. For a more sensitive determination of the amount of I in serum, a concentrated eluate of I-P from the  $\mu$ Porasil column is rechromatographed on a minicolumn (10 cm + 2 mm I.D.) packed with Lichrosorb SI-60 (5  $\mu$ m). With this method, a detection limit of 0.1 ng/mL for I in serum is obtained.

IT 53296-10-9 81613-39-0  
RL: ANT (Analyte); ANST (Analytical study)  
(determination of, in blood and urine of humans by high-performance liquid chromatog.)


RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



RN 81613-39-0 CAPLUS  
CN Adenosine, 2-[(4-hydroxyphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

L6 ANSWER 172 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1983:65261 CAPLUS

DN 98:65261


OREF 98:9845a,9848a

TI Interaction of 2-phenylamimoadenosine (CV 1808) with adenosine systems in

McIntosh

10/598,520


rat tissues  
AU Taylor, David A.; Williams, Michael  
CS Dep. Pharmacol., Merck Inst. Ther. Res., West Point, PA, 19486, USA  
SO European Journal of Pharmacology (1982), 85(3-4), 335-8  
CODEN: EJPHAZ; ISSN: 0014-2999  
DT Journal  
LA English  
GI



I

AB 2-chloroadenosine (2-CADO) [146-77-0], and CV 1808 (I) [53296-10-9] were compared in a central nervous system purinergic receptor binding assay and the inhibition of neurogenic contractions of the vas deferens. Both 2-CADO and CV 1808 were more potent than adenosine in both preps. CV 1808 was 10 times more active than dipyridamole in enhancing the response of the vas deferens to exogenous adenosine. Thus, CV 1808 may owe its potent coronary vasodilator activity to both a direct action on adenosine receptors and the ability to augment adenosine responses.  
IT 53296-10-9  
RL: BIOL (Biological study)  
(as adenosine agonist, vasodilator reactivity in relation to)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)


Absolute stereochemistry.



OSC.G 4 THERE ARE 4 CAPLUS RECORDS THAT CITE THIS RECORD (4 CITINGS)

L6 ANSWER 173 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1982:173891 CAPLUS  
DN 96:173891  
OREF 96:28487a,28490a  
TI Disposition and metabolism of 2-phenylaminoadenosine (CV-1808), a new coronary vasodilator, in rats and dogs  
AU Yoshida, Kiyoshi; Kondo, Takao; Kobayashi, Takuo; Tanayama, Shigeharu  
CS Cent. Res. Div., Takeda Chem. Ind., Osaka, Japan  
SO Takeda Kenkyushoho (1981), 40(3/4), 153-62  
CODEN: TAKHAA; ISSN: 0371-5167  
DT Journal  
LA English  
GI


McIntosh



AB The metabolic fate of <sup>14</sup>C-labeled CV-1808 (I) [53296-10-9] was studied in rats and dogs after oral administration. CV-1808 was absorbed by rats to give a maximum plasma level at 2 h postadministration and an apparent half-life of 2.3 h. In dogs, the plasma level peaked at 1 h and then declined with a half life of 4.9 h. After oral administration of labeled CV-1808 to rats, radioactivity was widely distributed in tissues with relatively higher concns. found in the gastrointestinal tract, liver, kidney, adrenal gland, lung, and plasma. In both rats and dogs, elimination of the compound was complete within 24-48 h with higher activities found in feces than in urine. The metabolites identified were 8-hydroxy-2-phenylaminoadenine [81613-42-5], 2-phenylaminoadenine [81613-41-4], 8-hydroxy-2-(p-hydroxyphenyl)aminoadenine [81613-40-3], and 2-(p-hydroxyphenyl)aminoadenosine [81613-39-0].


IT 81613-39-0  
 RL: BIOL (Biological study)  
 (as phenylaminoadenosine metabolite)  
 RN 81613-39-0 CAPLUS  
 CN Adenosine, 2-[(4-hydroxyphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 53296-10-9  
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
 (metabolism of)  
 RN 53296-10-9 CAPLUS  
 CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.




L6 ANSWER 174 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
 AN 1982:85941 CAPLUS  
 DN 96:85941  
 OREF 96:14127a,14130a

10/598,520

TI N2-(Alkanoylphenyl)-2,6-diaminonebularine  
PA Takeda Chemical Industries, Ltd., Japan  
SO Jpn. Kokai Tokkyo Koho, 8 pp.  
CODEN: JKXXAF  
DT Patent  
LA Japanese  
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 56131597	A	19811015	JP 1981-30882	19810303
PRAI	IL 1980-59602	A	19800312		
OS	CASREACT 96:85941				
GI					




AB The title compds. (I, R = alkanoyl) were prepared by cyclocondensation of II [R1-R3 = (protected) hydroxy] with RC6H4NHCR4:NH [R4 = (substituted) amino]. Thus, heating a mixture of II (R1 = R2 = R3 = HO) 10, m-H2NC6H4COMe 30, and m-MeCOC6H4NHC(:NH)NH2 14 g at 130° for 3 h gave 9.7 g I (R = m-MeCO). I (R = m-, p-MeCO, 3-MeCO-4-EtO) at 0.1 µg/dog showed 269.5-295.0% increase in coronary blood flow in 30 s.

IT 75106-29-5P 75106-30-8P 76888-18-1P  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of, as coronary vasodilator)

RN 75106-29-5 CAPLUS  
CN Adenosine, 2-[(4-acetylphenyl)amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 75106-30-8 CAPLUS  
CN Adenosine, 2-[(3-acetyl-4-ethoxyphenyl)amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

10/598,520



RN 76888-18-1 CAPLUS  
CN Adenosine, 2-[(3-acetylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




L6 ANSWER 175 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1982:600 CAPLUS  
DN 96:600  
OREF 96:107a,110a  
TI Effect of 2-phenylaminoadenosine (CV-1808) on ischemic ST-segment elevation in anesthetized dogs  
AU Matsumoto, Naohiko; Kawazoe, Katsuyoshi; Tanabe, Masao; Imamoto, Tetsuji; Fujiwara, Shuji; Hirata, Minoru  
CS Med. Res. Lab., Takeda Chem. Ind., Ltd., Osaka, Japan  
SO Journal of Cardiovascular Pharmacology (1981), 3(6), 1184-92  
CODEN: JPCPDT; ISSN: 0160-2446  
DT Journal  
LA English  
AB The effect of CV-1808 (2-phenylaminoadenosine) [53296-10-9] on myocardial ischemia was studied in anesthetized dogs. During i.v. infusion of CV-1808 (0.25 and 0.5  $\mu\text{g/kg/min}$  for 10 min) the ST-segment elevation in the epicardial ECG induced by a 5-min occlusion of a coronary arterial branch was occasionally enhanced in association with cardiac acceleration. In a dose of 0.5  $\mu\text{g/kg/min}$ , the agent inhibited the ST elevation 30 and 60 min after administration. The same dose did not change myocardial blood flow in the ischemic area despite significant systemic hypotension. In hearts with continuous coronary occlusion, CV-1808 (0.3 and 1.0  $\mu\text{g/kg}$ , i.v. bolus) increased the retrograde blood flow from the ischemic area immediately after administration, suggesting a collateral vasodilating action. Nifedipine (0.5 and 2.5  $\mu\text{g/kg/min}$ , i.v. for 10 min) and nitroglycerin (0.5 and 5.0  $\mu\text{g/kg/min}$ , i.v. for 10 min) had no influence on the ischemic ST-segment elevation, whereas a significant inhibition was seen with propranolol (0.5 mg/kg. i.v.). A moderate hypotension was induced by CV-1808, nifedipine, and nitroglycerin, whereas a significant reduction in cardiac function was seen after dosing with propranolol.  
IT 53296-10-9  
RL: BIOL (Biological study)  
(heart circulation and elec. activity response to, in ischemia)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)


Absolute stereochemistry.

McIntosh

10/598,520




L6 ANSWER 176 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1981:620263 CAPLUS  
DN 95:220263  
OREF 95:36765a,36768a  
TI Synthesis of 2-formyladenosine using diethoxyacetonitrile as a synthon  
AU Murakami, Teiichi; Otsuka, Masami; Kobayashi, Susumu; Ohno, Masaji  
CS Fac. Pharm. Sci., Univ. Tokyo, Tokyo, 113, Japan  
SO Heterocycles (1981), 16(8), 1315-19  
CODEN: HTCYAM; ISSN: 0385-5414  
DT Journal  
LA English  
GI



AB Imidazole I was heated with (EtO)2CHCN in BuOH-pyridine at 120° for 10 min in the presence of BuONa to give 90% nucleoside II [R = CH(COEt)2] which on hydrolysis with H2O-AcOH gave 96% II (R = CHO). II (R = CHO) was further converted into II (R = CH:NOH) and II (R = cyano).  
2-Formyladenine was analogously prepared  
IT 79936-11-1P  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)  
RN 79936-11-1 CAPLUS  
CN Adenosine, 2-cyano- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




OSC.G 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)

L6 ANSWER 177 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1981:604338 CAPLUS  
DN 95:204338  
OREF 95:34161a,34164a

McIntosh


10/598,520

TI Synthesis of 2-phenylaminoadenosine from imidazole nucleosides  
AU Omura, Kiyoshi; Marumoto, Ryuji; Furukawa, Yoshiyasu  
CS Cent. Res. Lab., Takeda Chem. Ind. Ltd., Osaka, 532, Japan  
SO Chemical & Pharmaceutical Bulletin (1981), 29(7), 1870-5  
CODEN: CPBTAL; ISSN: 0009-2363  
DT Journal  
LA English  
GI



AB The reaction of imidazole I with PhNCS gave 7-imino-5-phenylamino-3-( $\beta$ -D-ribofuranosyl)imidazo[4,5-d][1,3]-thiazine, which, on alkaline treatment, rearranged to 6-mercapto-2-phenylamino-9-( $\beta$ -D-ribofuranosyl)purine (II). On methylation, II gave the 6-methylmercapto derivative, which was converted to title adenosine (III) by treatment with  $\text{NH}_3$ . I reacted with PhNHCN in methanolic ammonia, giving III and 2-aminoadenosine as a by-product. Et 5-amino-1-( $\beta$ -D-ribofuranosyl)-4-carboximidate was directly obtained by treatment of 5-amino-1-(2,3,5-tri-O-propionyl- $\beta$ -D-ribofuranosyl)imidazole-4-carboxamide with Meerwein's reagent followed by deacylation, and this gave III by reaction with PhNHCN.  
IT 53296-10-9P  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD (3 CITINGS)

L6 ANSWER 178 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1981:425541 CAPLUS  
DN 95:25541  
OREF 95:4471a,4474a  
TI N2-Substituted 2,6-diaminonetrilarines  
PA Takeda Chemical Industries, Ltd., Japan  
SO Jpn. Tokkyo Koho, 8 pp.  
CODEN: JAXXAD  
DT Patent  
LA Japanese  
FAN.CNT 2


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
------------	------	------	-----------------	------

McIntosh



10/598,520


PI	JP 55049596	B	19801212	JP 1973-114542	19731011
	JP 50064296	A	19750531		
	DE 2359536	A1	19740612	DE 1973-2359536	19731129
	DE 2359536	C2	19840802		
	US 3936439	A	19760203	US 1973-420380	19731130
	FR 2209567	A1	19740705	FR 1973-43252	19731204
	CH 587864	A5	19770513	CH 1973-17069	19731205
	CH 601342	A5	19780714	CH 1976-12948	19731205
	NL 7316749	A	19740611	NL 1973-16749	19731206
	BE 808377	A1	19740607	BE 1973-138648	19731207
	GB 1418120	A	19751217	GB 1973-56781	19731207
	HU 167859	B	19751225	HU 1973-TA1284	19731207
	DK 134490	B	19761115	DK 1973-6631	19731207
	CA 1012534	A1	19770621	CA 1973-187678	19731207
PRAI	JP 1972-123602	A	19721208		
GI	JP 1973-114542	A	19731011		



AB Reaction of the ribofuranosides I (R = reactive group, R1 = alkyl, alkoxy, halo; R2-R4 = protected OH) with NH3 gave the N2-substituted 2,6-diaminonebularines I (R = NH2, R2-R4 = OH). Thus, 17.5 g 6-chloro-2-anilino-2',3',5'-tri-O-acetylnebularine was treated with NH3-MeOH at 120° to give 2-anilinoadenosine.

IT 53296-10-9P 53296-19-8P 53296-20-1P  
53296-21-2P  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)  
RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.




RN 53296-19-8 CAPLUS  
CN Adenosine, 2-[(4-methoxyphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 53296-20-1 CAPLUS  
CN Adenosine, 2-[(4-methylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 53296-21-2 CAPLUS  
CN Adenosine, 2-[(4-chlorophenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 4 THERE ARE 4 CAPLUS RECORDS THAT CITE THIS RECORD (4 CITINGS)

L6 ANSWER 179 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1981:425538 CAPLUS

DN 95:25538

OREF 95:4471a,4474a

TI N2-Substituted-2,6-diaminonebularines

PA Takeda Chemical Industries, Ltd., Japan

SO Jpn. Kokai Tokkyo Koho, 7 pp.


CODEN: JKXXAF

DT Patent

LA Japanese

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 55136299	A	19801023	JP 1979-43257	19790409
PRAI	JP 1979-43257	A	19790409		
GI					



AB Eight nebularines I [(R, R1) = (H, H), (H, Me), (Me, H), (H, OEt), etc.] were prepared by treating II (R2 = CONH2, R3 = H) with (EtCO)2O and then with Et3O.BF4 to give II [R2 = C(:NH)OEt, R3 = EtCO], followed by reaction with the appropriate RR1C6H3NHCN, (PhNH)2C:NH, triphenylmelamine, or phenylguanidine carbonate with or without previous deprotection. Thus, 258 g II (R2 = CONH2, R3 = H) in pyridine and 400 mL (EtCO)2O were stirred 16 h at room temperature to give 355 g II (R2 = CONH2, R3 = EtCO), which (10 g) in CH2Cl2 was added dropwise to 7.2 g Et3O.BF4 in CH2Cl2 with stirring and ice cooling and the mixture was left 20 h in ice to give 14 g II [R2 = C(:NH)OEt, R3 = EtCO], which (10 g) was heated with 12 g PhNHCN in 20% MeOH-NH4OH 5 h at 180° in a sealed vessel to give 1.1 g I (R = R1 = H).


IT 53296-10-9P 53296-20-1P 70590-18-0P  
70590-23-7P 70590-28-2P 74615-40-0P  
76888-17-0P

RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)


Absolute stereochemistry.



RN 53296-20-1 CAPLUS

CN Adenosine, 2-[(4-methylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 70590-18-0 CAPLUS

CN Adenosine, 2-[(3-chlorophenyl)amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

10/598,520




RN 70590-23-7 CAPLUS  
CN Adenosine, 2-[(3-methylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 70590-28-2 CAPLUS  
CN Adenosine, 2-[(3,4-dimethoxyphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 74615-40-0 CAPLUS  
CN Adenosine, 2-[[4-[(cyclohexylamino)carbonyl]phenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 76888-17-0 CAPLUS  
CN Adenosine, 2-[(4-ethoxyphenyl)amino]- (9CI) (CA INDEX NAME)

McIntosh

10/598,520

Absolute stereochemistry.




L6 ANSWER 180 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1981:121883 CAPLUS  
DN 94:121883  
OREF 94:19951a,19954a  
TI 2,6-Diaminonebularines  
IN Sawa, Yoichi; Kawakami, Yoshiyuki; Marumoto, Ryuji  
PA Takeda Chemical Industries, Ltd., Japan  
SO Eur. Pat. Appl., 27 pp.  
CODEN: EPXXDW  
DT Patent  
LA English  
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	EP 17465	A1	19801015	EP 1980-301024	19800401
	R: AT, BE, CH, DE, FR, GB, IT, NL, SE				
	JP 55130998	A	19801011	JP 1979-39562	19790402
	NO 8000868	A	19801003	NO 1980-868	19800325
	DK 8001345	A	19801003	DK 1980-1345	19800328
	US 4293690	A	19811006	US 1980-136072	19800328
	AU 8056982	A	19801009	AU 1980-56982	19800331
	FI 8001033	A	19801003	FI 1980-1033	19800401
PRAI	JP 1979-39562	A	19790402		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OS CASREACT 94:121883; MARPAT 94:121883

GI



AB Nebularines I (R = Ph, substituted phenyl) were prepared by cyclization of nucleosides II (R1, R2, R3 = protected or unprotected OH) with RNHC(:NH)R4 (same R; R4 = NH2, substituted amino, alkylthio) followed by deprotection where necessary. Thus, II (R1 = R2 = R3 = OH) was heated with 1,3-diphenylguanidine in PhNH2 at 150-5° to give 68.8% I (R = Ph).

IT 53296-10-9P 53296-20-1P 70590-22-6P  
70590-23-7P 70590-27-1P 74615-32-0P  
74615-36-4P 74615-40-0P 75106-30-8P  
76888-17-0P 76888-18-1P

RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)


RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

McIntosh


10/598,520

Absolute stereochemistry.




RN 53296-20-1 CAPLUS  
CN Adenosine, 2-[(4-methylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 70590-22-6 CAPLUS  
CN Adenosine, 2-[(3-methoxyphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 70590-23-7 CAPLUS  
CN Adenosine, 2-[(3-methylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 70590-27-1 CAPLUS  
CN Adenosine, 2-[(3-chloro-4-methylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

McIntosh


10/598,520



RN 74615-32-0 CAPLUS

CN Adenosine, 2-[[4-(aminocarbonyl)phenyl]amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 74615-36-4 CAPLUS

CN Adenosine, 2-[[3-(aminocarbonyl)phenyl]amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 74615-40-0 CAPLUS

CN Adenosine, 2-[[4-[(cyclohexylamino)carbonyl]phenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 75106-30-8 CAPLUS

CN Adenosine, 2-[(3-acetyl-4-ethoxyphenyl)amino]- (9CI) (CA INDEX NAME)

McIntosh


10/598,520

Absolute stereochemistry.




RN 76888-17-0 CAPLUS  
CN Adenosine, 2-[(4-ethoxyphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 76888-18-1 CAPLUS  
CN Adenosine, 2-[(3-acetylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 4 THERE ARE 4 CAPLUS RECORDS THAT CITE THIS RECORD (4 CITINGS)


L6 ANSWER 181 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1980:568559 CAPLUS  
DN 93:168559  
OREF 93:26863a,26866a  
TI 2,6-Diaminonebularines  
IN Marumoto, Ryuji; Tanabe, Masao; Furukawa, Yoshiyasu  
PA Takeda Chemical Industries, Ltd., Japan  
SO Ger. Offen., 33 pp.  
CODEN: GWXXBX  
DT Patent  
LA German  
FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	DE 2941592	A1	19800424	DE 1979-2941592	19791013
	JP 55053299	A	19800418	JP 1978-127109	19781016
	JP 62003159	B	19870123		
	JP 56012400	A	19810206	JP 1979-87074	19790709
	JP 63003876	B	19880126		
PRAI	JP 1978-127109	A	19781016		
	JP 1979-87074	A	19790709		

GI

McIntosh





AB Diaminonebularines I (R = carbamoyl, acyl; R1 = H, halogen, alkoxy) were prepared Thus 4-H<sub>2</sub>NC<sub>6</sub>H<sub>4</sub>CONH<sub>2</sub>.HCl was treated with KSCN to give 4-H<sub>2</sub>NCOC<sub>6</sub>H<sub>4</sub>NHCSNH<sub>2</sub> which was treated with Pb(OAc)<sub>4</sub> to give 4-H<sub>2</sub>NCOC<sub>6</sub>H<sub>4</sub>NHCN. Treatment of the latter compound with 5-amino-1-β-D-ribofuranosyl-4-cyanoimidazole gave I (R = 4-H<sub>2</sub>NCO, R1 = H) which at 0.1 μg increased the coronary blood flow in dogs by 199% 30 s after administration.


IT 74615-32-0P 74615-33-1P 74615-38-6P  
 74615-39-7P 74615-40-0P 74615-42-2P  
 75106-22-8P 75106-25-1P 75106-26-2P  
 75106-29-5P 75106-30-8P 75106-31-9P

RL: SPN (Synthetic preparation); PREP (Preparation)  
 (preparation and coronary vasodilator activity of)

RN 74615-32-0 CAPLUS

CN Adenosine, 2-[[4-(aminocarbonyl)phenyl]amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 74615-33-1 CAPLUS

CN Adenosine, 2-[[4-[(propylamino)carbonyl]phenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 74615-38-6 CAPLUS

CN Adenosine, 2-[[4-[(methylamino)carbonyl]phenyl]amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

10/598,520




RN 74615-39-7 CAPLUS  
CN Adenosine, 2-[[4-[(ethylamino)carbonyl]phenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 74615-40-0 CAPLUS  
CN Adenosine, 2-[[4-[(cyclohexylamino)carbonyl]phenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 74615-42-2 CAPLUS  
CN Adenosine, 2-[[4-[(dimethylamino)carbonyl]phenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




McIntosh

10/598,520

RN 75106-22-8 CAPLUS

CN Adenosine, 2-[[2-(aminocarbonyl)phenyl]amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 75106-25-1 CAPLUS

CN Adenosine, 2-[[2-(1-oxopropyl)phenyl]amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 75106-26-2 CAPLUS

CN Adenosine, 2-[(2-acetylphenyl)amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 75106-29-5 CAPLUS

CN Adenosine, 2-[(4-acetylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 75106-30-8 CAPLUS

CN Adenosine, 2-[(3-acetyl-4-ethoxyphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 75106-31-9 CAPLUS  
CN Adenosine, 2-[[4-(1-oxobutyl)phenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 74615-37-5P 74615-41-1P 75106-23-9P  
75106-24-0P 75106-32-0P 75106-33-1P  
75106-34-2P  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)  
RN 74615-37-5 CAPLUS  
CN Adenosine, 2-[[4-(aminocarbonyl)-3-chlorophenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 74615-41-1 CAPLUS  
CN Adenosine, 2-[[4-[(phenylamino)carbonyl]phenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 75106-23-9 CAPLUS  
CN Adenosine, 2-[[2-[(phenylamino)carbonyl]phenyl]amino]- (9CI) (CA INDEX NAME)

McIntosh

10/598,520


Absolute stereochemistry.



RN 75106-24-0 CAPLUS

CN Adenosine, 2-[[4-[(ethylamino)carbonyl]phenyl]amino]-, monohydrochloride (9CI) (CA INDEX NAME)

Absolute stereochemistry.




● HCl

RN 75106-32-0 CAPLUS

CN Adenosine, 2-[(4-acetyl-3-methoxyphenyl)amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 75106-33-1 CAPLUS

CN Adenosine, 2-[(4-acetyl-3-chlorophenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 75106-34-2 CAPLUS

McIntosh

10/598,520

CN Adenosine, 2-[(4-acetylphenyl)amino]-, monohydrochloride (9CI) (CA INDEX NAME)

Absolute stereochemistry.



● HCl

L6 ANSWER 182 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1980:525711 CAPLUS

DN 93:125711

OREF 93:19905a,19908a

TI Coronary and cardiohemodynamic effects of 2-phenylaminoadenosine (CV-1808) in anesthetized dogs and cats

AU Kawazoe, K.; Matsumoto, N.; Tanabe, M.; Fujiwara, S.; Yanagimoto, M.; Hirata, M.; Kikuchi, K.

CS Cent. Res. Div., Takeda Chem. Ind. Ltd., Osaka, 532, Japan


SO Arzneimittelforschung (1980), 30(7), 1083-7

CODEN: ARZNAD; ISSN: 0004-4172

DT Journal

LA English

GI



I

AB The coronary vasodilating effect of intracoronary or i.v. CV 1808 (I) [53296-10-9] in dogs was greater than that of nifedipine, nitroglycerin, or dipyridamole. I increased blood flow to the superior mesenteric artery to a lesser extent than blood flow to the coronary vascular bed, and blood flow to the femoral artery was decreased. I.v. I caused a dose-dependent increase in left ventricular dp/dt, which was inhibited by pretreatment with propranolol. I was well absorbed from the intestinal tract.

IT 53296-10-9

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)  
(coronary vasodilating activity of)


RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.

McIntosh

10/598,520



OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

L6 ANSWER 183 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1980:495598 CAPLUS

DN 93:95598

OREF 93:15349a,15352a

TI N2-Substituted phenyl-2,6-diaminonebularines

PA Takeda Chemical Industries, Ltd., Japan

SO Jpn. Kokai Tokkyo Koho, 6 pp.


CODEN: JKXXAF

DT Patent

LA Japanese

FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 55053299	A	19800418	JP 1978-127109	19781016
	JP 62003159	B	19870123		
	AU 7951189	A	19800424	AU 1979-51189	19790926
	CH 642668	A5	19840430	CH 1979-9083	19791009
	DK 7904303	A	19800417	DK 1979-4303	19791011
	SE 7908480	A	19800417	SE 1979-8480	19791012
	US 4258033	A	19810324	US 1979-85057	19791012
	DE 2941592	A1	19800424	DE 1979-2941592	19791013
	NL 7907611	A	19800418	NL 1979-7611	19791015
	CA 1112641	A1	19811117	CA 1979-337577	19791015
	BE 879436	A1	19800416	BE 1979-197665	19791016
	FR 2439207	A1	19800516	FR 1979-25642	19791016
	FR 2439207	B1	19820611		
	GB 2034704	A	19800611	GB 1979-35932	19791016
	GB 2034704	B	19830330		
PRAI	JP 1978-127109	A	19781016		
	JP 1979-87074	A	19790709		
OS	MARPAT 93:95598				
GI					



AB Twelve title nebularines I [one of R and R1 is CONR2R3 (R2 = H, alkyl; R3 = H, alkyl, cyclohexyl, Ph) and the other is H or halo], having coronary vasodilating activity (data given in dogs), were prepared Thus, a mixture of 10 g 5-amino-1-β-D-ribofuranosyl-4-cyanoimidazole, 12 g 4-H2NCOC6H4NHCN, and 150 mL 20% MeOH-NH3 was autoclaved 5 h at 80° to give 2 g I (R = H, R1 = H2NCO).

IT 74615-32-0P 74615-33-1P 74615-34-2P  
74615-35-3P 74615-36-4P 74615-37-5P

McIntosh

10/598,520

74615-38-6P 74615-39-7P 74615-40-0P


74615-41-1P 74615-42-2P

RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)

RN 74615-32-0 CAPLUS

CN Adenosine, 2-[[4-(aminocarbonyl)phenyl]amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 74615-33-1 CAPLUS

CN Adenosine, 2-[[4-[(propylamino)carbonyl]phenyl]amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 74615-34-2 CAPLUS

CN Adenosine, 2-[[4-[(ethylamino)carbonyl]phenyl]amino]-, hydrochloride (9CI)  
(CA INDEX NAME)

Absolute stereochemistry.



● x HCl

RN 74615-35-3 CAPLUS


CN Adenosine, 2-[[3-[(phenylamino)carbonyl]phenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

McIntosh




10/598,520




RN 74615-36-4 CAPLUS  
CN Adenosine, 2-[[3-(aminocarbonyl)phenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 74615-37-5 CAPLUS  
CN Adenosine, 2-[[4-(aminocarbonyl)-3-chlorophenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 74615-38-6 CAPLUS  
CN Adenosine, 2-[[4-[(methylamino)carbonyl]phenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 74615-39-7 CAPLUS  
CN Adenosine, 2-[[4-[(ethylamino)carbonyl]phenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 74615-40-0 CAPLUS  
CN Adenosine, 2-[[4-[(cyclohexylamino)carbonyl]phenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 74615-41-1 CAPLUS  
CN Adenosine, 2-[[4-[(phenylamino)carbonyl]phenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 74615-42-2 CAPLUS  
CN Adenosine, 2-[[4-[(dimethylamino)carbonyl]phenyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




McIntosh

10/598,520

OSC.G 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)

L6 ANSWER 184 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1980:495591 CAPLUS  
DN 93:95591  
OREF 93:15345a,15348a  
TI 2-Substituted adenosine derivatives  
IN Ueda, Tooru; Matsuda, Akira; Nomoto, Juji  
PA Yamasa Shoyu Co., Ltd., Japan  
SO Jpn. Kokai Tokkyo Koho, 6 pp.  
CODEN: JKXXAF  
DT Patent  
LA Japanese  
FAN.CNT 1


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 55036419	A	19800314	JP 1978-109655	19780908
	JP 63003875	B	19880126		
PRAI	JP 1978-109655	A	19780908		
GI					



AB The title compds. I (R = HN:CR1 where R1 = alkoxy, OH, NH2) were prepared by treating OH-protected I (R = CN) with alkoxides. Thus, 418 mg 2',3',5'-O-triacetyl-2-cyanoadenosine reacted with MeONa-MeOH at room temperature for 17 h followed by treatment with Dowex 50 to give 288 mg I [R = HN:C(OMe)], whose hydrolysis (HCl) gave I (R = CO2Me), which was saponified to give I (R = CO2Na).

IT 70255-72-0P  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)  
RN 70255-72-0 CAPLUS  
CN Adenosine, 2-(aminocarbonyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)

L6 ANSWER 185 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1979:508193 CAPLUS  
DN 91:108193  
OREF 91:17475a,17478a  
TI 2,6-Diaminonebularines  
IN Marumoto, Ryuji; Shima, Shunsuke; Furukawa, Yoshiyasu  
PA Takeda Chemical Industries, Ltd., Japan

McIntosh

10/598,520

SO Ger. Offen., 27 pp.


CODEN: GWXXBX

DT Patent

LA German

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	DE 2845435	A1	19790426	DE 1978-2845435	19781019
	JP 54061194	A	19790517	JP 1977-127147	19771021
	GB 2007664	A	19790523	GB 1978-39583	19781006
	GB 2007664	B	19820526		
	AU 7840511	A	19800417	AU 1978-40511	19781009
	AU 521358	B2	19820401		
	ZA 7805762	A	19790926	ZA 1978-5762	19781012
	CA 1102794	A1	19810609	CA 1978-313340	19781013
	FI 7803181	A	19790422	FI 1978-3181	19781018
	SE 7810854	A	19790422	SE 1978-10854	19781018
	DK 7804655	A	19790422	DK 1978-4655	19781019
	BE 871422	A1	19790420	BE 1978-191251	19781020
	NL 7810519	A	19790424	NL 1978-10519	19781020
	NO 7803559	A	19790424	NO 1978-3559	19781020
	FR 2406640	A1	19790518	FR 1978-29945	19781020
	FR 2406640	B1	19820528		
	AT 7807552	A	19810115	AT 1978-7552	19781020
	AT 363619	B	19810825		
	US 4255565	A	19810310	US 1978-953255	19781020
PRAI	JP 1977-127147	A	19771021		
OS	MARPAT 91:108193				
GI					



AB Diaminonebularines I (R = optionally substituted Ph, cyclohexyl) were prepared by treating the aminoimidazolecarbonitrile II or its protected derivs. with RN:C:NR<sub>4</sub> (R<sub>4</sub> = H, R). Thus, 5-amino-1-β-D-ribofuranosyl-4-imidazolecarboxamide was acylated, dehydrated, and deacylated to give II, which was treated with PhNH<sub>2</sub> to give I (R = Ph). PhNH<sub>2</sub> was prepared by treating PhNH<sub>2</sub> with KSCN and H<sub>2</sub>S elimination from PhNHCSNH<sub>2</sub> with KOH and Pb(OAc)<sub>4</sub>.

IT 53296-10-9P 53296-20-1P 53296-21-2P  
70590-18-0P 70590-20-4P 70590-22-6P  
70590-23-7P 70590-25-9P 70590-26-0P  
70590-27-1P 70590-28-2P 70590-29-3P  
70590-30-6P 70590-31-7P 71231-75-9P  
71231-76-0P 71231-77-1P 71231-78-2P  
71231-79-3P 71231-80-6P 71231-81-7P  
71231-82-8P 71231-83-9P 71231-84-0P  
71231-85-1P 71231-86-2P

RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)


RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 53296-20-1 CAPLUS  
CN Adenosine, 2-[(4-methylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 53296-21-2 CAPLUS  
CN Adenosine, 2-[(4-chlorophenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 70590-18-0 CAPLUS  
CN Adenosine, 2-[(3-chlorophenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 70590-20-4 CAPLUS  
CN Adenosine, 2-[(3,4-dimethylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 70590-22-6 CAPLUS  
CN Adenosine, 2-[(3-methoxyphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 70590-23-7 CAPLUS  
CN Adenosine, 2-[(3-methylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 70590-25-9 CAPLUS  
CN Adenosine, 2-[(3-fluorophenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 70590-26-0 CAPLUS  
CN Adenosine, 2-[(3-bromophenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

McIntosh


10/598,520



RN 70590-27-1 CAPLUS

CN Adenosine, 2-[(3-chloro-4-methylphenyl)amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 70590-28-2 CAPLUS

CN Adenosine, 2-[(3,4-dimethoxyphenyl)amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 70590-29-3 CAPLUS

CN Adenosine, 2-[(3-chloro-4-methoxyphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 70590-30-6 CAPLUS

CN Adenosine, 2-[(3,4-dichlorophenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 70590-31-7 CAPLUS  
CN Adenosine, 2-[(3-butylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 71231-75-9 CAPLUS  
CN Adenosine, 2-(phenylamino)-, monohydrochloride (9CI) (CA INDEX NAME)


Absolute stereochemistry.



● HCl

RN 71231-76-0 CAPLUS  
CN Adenosine, 2-[(2-methylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 71231-77-1 CAPLUS  
CN Adenosine, 2-[(2-methoxyphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

McIntosh




10/598,520




RN 71231-78-2 CAPLUS  
CN Adenosine, 2-[(4-bromophenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 71231-79-3 CAPLUS  
CN Adenosine, 2-[(4-fluorophenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 71231-80-6 CAPLUS  
CN Adenosine, 2-[(2-fluorophenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 71231-81-7 CAPLUS  
CN Adenosine, 2-[(4-ethylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

McIntosh


10/598,520



RN 71231-82-8 CAPLUS

CN Adenosine, 2-[(2,5-dimethylphenyl)amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 71231-83-9 CAPLUS

CN Adenosine, 2-[(3,5-dimethylphenyl)amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 71231-84-0 CAPLUS

CN Adenosine, 2-[(2,4-dimethylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 71231-85-1 CAPLUS

CN Adenosine, 2-[(4-chloro-2-methoxyphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520



RN 71231-86-2 CAPLUS  
CN Adenosine, 2-[(2,3-dichlorophenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




OSC.G 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)

L6 ANSWER 186 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1979:491903 CAPLUS  
DN 91:91903  
OREF 91:14866h,14867a  
TI Synthesis and enzymic activity of adenosine 3',5'-cyclic phosphate analogs  
AU Marumoto, Ryuji; Yoshioka, Yoshio; Naka, Takehiko; Shima, Shunsuke; Miyashita, Osamu; Maki, Yoshitaka; Suzuki, Tsuyoshi; Honjo, Mikio  
CS Cent. Res. Div., Takeda Chem. Ind., Ltd., Osaka, 532, Japan  
SO Chemical & Pharmaceutical Bulletin (1979), 27(4), 990-1003  
CODEN: CPBTAL; ISSN: 0009-2363  
DT Journal  
LA English  
AB 2-Mono- and 2,6-disubstituted cAMP (cAMP = adenosine 3',5'-cyclic phosphate) were derived from the corresponding adenosine analogs via the 5'-phosphates. 2-Substituted cAMP and aristeromycin 3',6'-cyclic phosphate were converted to the 8-bromo derivs. 8-(3,5-Dimethylpyrazol-1-yl)-cAMP was derived from 8-hydrazinoadenosine. 8-(2-Hydroxypropyl-2)-cAMP was prepared by  $\gamma$ -ray irradiation of cAMP in isopropanol. N6-Butyl-2-phenyl-cAMP was derived from 2-phenylinosine. 2-Or 8-substituted cAMP was converted to its dibutyryl derivative The activities of these new analogs were assessed with cAMP-dependent protein kinases (PK) and cAMP phosphodiesterases (PDE). 8-Bromo-2-chloro-, 2-chloro-, 2-bromo-, 2-phenylthio-, 8-carbamoyl- and 8-carboxy-cAMP were better activators of PK than cAMP, while 2-substituted analogs were significant substrates and inhibitors of PDE. 2-Phenyl-, benzyl-, phenoxy-, chloro- and bromo-cAMP or aristeromycin 3',6'-cyclic phosphate had an inhibitory effect on the binding of cAMP to PK equal to or greater than that of cAMP.  
IT 50257-82-4 50257-84-6  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(phosphorylation of, with pyrophosphoryl chloride)  
RN 50257-82-4 CAPLUS  
CN Adenosine, 2-phenoxy- (CA INDEX NAME)


Absolute stereochemistry.

10/598,520



RN 50257-84-6 CAPLUS  
CN Adenosine, 2-butoxy- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



OSC.G 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)

L6 ANSWER 187 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1979:474850 CAPLUS  
DN 91:74850  
OREF 91:12117a,12120a  
TI N2-Substituted phenyl-2,6-diaminonebularine  
IN Marumoto, Ryuji; Tanabe, Masao; Furukawa, Yoshiyasu  
PA Takeda Chemical Industries, Ltd., Japan  
SO Ger. Offen., 24 pp.  
CODEN: GWXXBX  
DT Patent  
LA German  
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	DE 2845496	A1	19790426	DE 1978-2845496	19781019
	JP 54061195	A	19790517	JP 1977-127148	19771021
	GB 2007213	A	19790516	GB 1978-39582	19781006
	GB 2007213	B	19820526		
	AU 7840512	A	19800417	AU 1978-40512	19781009
	AU 521102	B2	19820318		
	SE 7810905	A	19790422	SE 1978-10905	19781019
	NL 7810520	A	19790424	NL 1978-10520	19781020
	FR 2406641	A1	19790518	FR 1978-29946	19781020
	FR 2406641	B1	19820611		
	US 4225591	A	19800930	US 1978-953254	19781020
PRAI	JP 1977-127148	A	19771021		
OS	MARPAT 91:74850				
GI					




McIntosh

10/598,520

AB The title compds. I (R = R1 = H, halogen, lower alkyl or alkoxy) and their salts were prepared for use as coronary vasodilators (test data tabulated). Thus, 5-amino-1- $\beta$ -D-ribofuranosyl-4-cyanoimidazole reacted with 3-ClC<sub>6</sub>H<sub>4</sub>NHCN to give I (R = H, R1 = Cl).  
IT 70590-19-1P 70590-20-4P 70590-22-6P  
70590-23-7P 70590-25-9P 70590-27-1P  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation and coronary vasodilator activity of)  
RN 70590-19-1 CAPLUS  
CN Adenosine, 2-[(3-chlorophenyl)amino]-, monohydrochloride (9CI) (CA INDEX NAME)


Absolute stereochemistry.



● HCl


RN 70590-20-4 CAPLUS  
CN Adenosine, 2-[(3,4-dimethylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 70590-22-6 CAPLUS  
CN Adenosine, 2-[(3-methoxyphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 70590-23-7 CAPLUS  
CN Adenosine, 2-[(3-methylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 70590-25-9 CAPLUS  
CN Adenosine, 2-[(3-fluorophenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 70590-27-1 CAPLUS  
CN Adenosine, 2-[(3-chloro-4-methylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 70590-18-0P 70590-24-8P 70590-26-0P  
70590-28-2P 70590-29-3P 70590-30-6P  
70590-31-7P  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)  
RN 70590-18-0 CAPLUS  
CN Adenosine, 2-[(3-chlorophenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 70590-24-8 CAPLUS  
CN Adenosine, 2-[(3-methylphenyl)amino]-, monohydrochloride (9CI) (CA INDEX NAME)

Absolute stereochemistry.

McIntosh


10/598,520



● HCl


RN 70590-26-0 CAPLUS  
CN Adenosine, 2-[(3-bromophenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 70590-28-2 CAPLUS  
CN Adenosine, 2-[(3,4-dimethoxyphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 70590-29-3 CAPLUS  
CN Adenosine, 2-[(3-chloro-4-methoxyphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 70590-30-6 CAPLUS  
CN Adenosine, 2-[(3,4-dichlorophenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520



RN 70590-31-7 CAPLUS  
CN Adenosine, 2-[(3-butylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

L6 ANSWER 188 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1979:439769 CAPLUS

DN 91:39769

OREF 91:6497a,6500a

TI Nucleosides and nucleotides. XXVII. Synthesis of 2- and 8-cyanoadenosines and their derivatives

AU Matsuda, Akira; Nomoto, Yuji; Ueda, Tohru

CS Fac. Pharm. Sci., Hokkaido Univ., Sapporo, 060, Japan

SO Chemical & Pharmaceutical Bulletin (1979), 27(1), 183-92

CODEN: CPBTAL; ISSN: 0009-2363

DT Journal

LA English

OS CASREACT 91:39769

AB A facile displacement of a methylsulfonyl group in adenosines with cyanide ion is described. Treatment of protected 8-(methylsulfonyl)adenosines with NaCN in DMF gave the 8-cyanoadenosine. The conversion of the cyano group to the Me imidate, methoxycarbonyl, carbamoyl, and carboxylic acid was achieved. Similar reaction was carried out with 2-(methylsulfonyl)adenosine to give the 2-cyanoadenosine and their derivs. The NMR and CD spectra of these 2- and 8-substituted adenosines are given. The 8-substituted adenosines possess syn-conformations while the 2-substituted derivs. prefer anti-conformations, as confirmed by the CD and NMR spectra.


IT 70255-72-0P

RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation) (preparation and spectra of)

RN 70255-72-0 CAPLUS

CN Adenosine, 2-(aminocarbonyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



McIntosh




10/598,520

OSC.G 20 THERE ARE 20 CAPLUS RECORDS THAT CITE THIS RECORD (21 CITINGS)

L6 ANSWER 189 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1975:514827 CAPLUS  
DN 83:114827  
OREF 83:18059a,18062a  
TI 2-Alkoxyadenosines  
IN Honjo, Mikio; Marumoto, Ryuji; Yoshioka, Yoshio  
PA Takeda Chemical Industries, Ltd., Japan  
SO Jpn. Kokai Tokkyo Koho, 4 pp.  
CODEN: JKXXAF  
DT Patent  
LA Japanese  
FAN.CNT 1


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 50053393	A	19750512	JP 1973-105286	19730918
PRAI	JP 1973-105286	A	19730918		
AB	2-Haloadenosines, where the 2'- and 3'-OH groups are protected, are treated with an aliphatic alc. and base to give 2',3'-protected 2-alkoxyadenosines. 2-Alkoxyadenosines are prepared by hydrolysis. The protected products have coronary vasodilatory, hypotensive, and diuretic activities (no data). Thus, a mixture of 6.6 g 2-chloroadenosine, 55 ml HC(OEt)3, 10 ml DMF, and 0.8 g p-toluenesulfonic acid was stirred at 30° for 0.5 hr, poured into aqueous NaHCO3, and extracted with CHCl3 to give 2',3'-O-ethoxymethylidene derivative which was heated with 3 g NaOH in 62 ml BuOH at 90° for 1 hr to give 2',3'-O-ethoxymethylidene-2-butoxyadenosine. Hydrolysis in 40% aqueous AcOH at 35° for 2 days gave 2-butoxyadenosine. Similarly prepared was 2-pentyloxyadenosine.				
IT	50257-84-6P 50257-85-7P				
RL	SPN (Synthetic preparation); PREP (Preparation) (preparation of)				
RN	50257-84-6 CAPLUS				
CN	Adenosine, 2-butoxy- (9CI) (CA INDEX NAME)				

Absolute stereochemistry.



RN 50257-85-7 CAPLUS  
CN Adenosine, 2-(pentyloxy)- (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)


L6 ANSWER 190 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1975:479518 CAPLUS  
DN 83:79518  
OREF 83:12499a,12502a

McIntosh

10/598,520


TI Synthesis and coronary vasodilating activity of 2-substituted adenosines  
AU Marumoto, Ryuji; Yoshioka, Yoshio; Miyashita, Osamu; Shima, Shunsuke;  
Imai, Kinichi; Kawazoe, Katsuyoshi; Honjo, Mikio  
CS Cent. Res. Div., Takeda Chem. Ind., Osaka, Japan  
SO Chemical & Pharmaceutical Bulletin (1975), 23(4), 759-74  
CODEN: CPBTAL; ISSN: 0009-2363  
DT Journal  
LA English  
AB 2-Haloadenosines were prepared by acetylation of 2-haloinosines followed by  
chlorination and amination. 2-Alkoxyadenosines were prepared by protection  
of 2'- and 3'-OH groups of 2-chloroadenosine (I) or 2-chloroinosine,  
followed by substitution of the C atom with alkoxy group. The reaction of  
5-amino-4-cyano-1- $\beta$ -D-ribofuranosylimidazole with CS<sub>2</sub> afforded  
2,6-di-mercapto-9- $\beta$ -D-ribofuranosylpurine, which was converted to  
2-mercaptoadenosine and its S-substituted derivs. 2-Phenylaminoadenosine  
(II) was prepared from 2-phenylamino-2',3',5'-tri-O-acetylinosine, which was  
prepared by acetylation of 2-phenylaminoinosine with AcCl in HOAc.  
O-substituted 2-hydroxyadenosines, S-substituted 2-mercaptadenosines,  
N2-substituted 2-aminoadenosines, 2-alkyl- and -aryl-adenosines were  
prepared among which several compds. had coronary vasodilating potency. II  
showed not only a strong potency, but also a longer duration of the effect  
than that of I.  
IT 50257-82-4 50257-85-7 50257-89-1  
50257-90-4 50257-91-5 50257-97-1  
53296-19-8 53296-20-1 56720-62-8  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(coronary vasodilating activity of)  
RN 50257-82-4 CAPLUS  
CN Adenosine, 2-phenoxy- (CA INDEX NAME)

Absolute stereochemistry.



RN 50257-85-7 CAPLUS  
CN Adenosine, 2-(pentyloxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 50257-89-1 CAPLUS  
CN Adenosine, 2-(2-hydroxyethoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 50257-90-4 CAPLUS  
CN Adenosine, 2-(3-methylphenoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 50257-91-5 CAPLUS  
CN Adenosine, 2-(4-methoxyphenoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 50257-97-1 CAPLUS  
CN Adenosine, 2-(2-chlorophenoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 53296-19-8 CAPLUS  
CN Adenosine, 2-[(4-methoxyphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 53296-20-1 CAPLUS  
CN Adenosine, 2-[(4-methylphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 56720-62-8 CAPLUS  
CN Adenosine, 2-(3-methoxybutoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 50257-84-6P 53296-10-9P  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation and coronary vasodilating activity of)  
RN 50257-84-6 CAPLUS  
CN Adenosine, 2-butoxy- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 53296-10-9 CAPLUS  
CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)

Absolute stereochemistry.

McIntosh



OSC.G 20 THERE ARE 20 CAPLUS RECORDS THAT CITE THIS RECORD (20 CITINGS)

L6 ANSWER 191 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1975:156651 CAPLUS

DN 82:156651

OREF 82:25025a,25028a

TI 2-Substituted adenosines

IN Miyashita, Osamu; Yoshioka, Yoshio; Honjo, Mikio

PA Takeda Chemical Industries, Ltd.

SO Jpn. Kokai Tokkyo Koho, 8 pp.

CODEN: JKXXAF


DT Patent

LA Japanese

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 49124096	A	19741127	JP 1973-37234	19730331
PRAI	JP 1973-37234	A	19730331		
GI	For diagram(s), see printed CA Issue.				
AB	2-Substituted adenosines [I; R = lower alkyl, R5O(CH2)n (R5 = H, lower alkyl, Ph acyl; n = 2-6), phenyl] were prepared by treating 2,6-disubstituted nebularine (II; R2 = H, acyl; R3 = active groups convertible into an NH2 group by reaction with NH3) with NH3. I had coronary vasodilating (in dogs) and hypotensive actions. Thus, 2.8 g 2-(β-methoxyethoxy)-6-chloro-2',3',5'-tri-O-acetylnebularine [prepared from 2-(β-methoxyethoxy)-inosine (III) via 2-(β-methoxyethoxy)-2',3',5'-tri-O-acetylinosine] was autoclaved with 20 ml NH3-MeOH 5 hr at 100° to give 0.9 g 2-(β-methoxyethoxy)adenosine. Among 13 more I prepared were 2-butoxy-, 2-(β-ethoxyethoxy)-, 2-(β-hydroxyethoxy)-, and 2-(β-phenoxyethoxy)adenosines.				
IT	50257-90-4P	50257-91-5P	50257-93-7P		
	50257-95-9P	50447-11-5P			
RL	SPN (Synthetic preparation); PREP (Preparation) (preparation of)				
RN	50257-90-4	CAPLUS			
CN	Adenosine, 2-(3-methylphenoxy)- (9CI) (CA INDEX NAME)				

Absolute stereochemistry.




RN 50257-91-5 CAPLUS

CN Adenosine, 2-(4-methoxyphenoxy)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

10/598,520




RN 50257-93-7 CAPLUS  
CN Adenosine, 2-(4-hydroxybutoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 50257-95-9 CAPLUS  
CN Adenosine, 2-(hexyloxy)- (CA INDEX NAME)

Absolute stereochemistry.



RN 50447-11-5 CAPLUS  
CN Adenosine, 2-[(6-hydroxyhexyl)oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L6 ANSWER 192 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1974:491898 CAPLUS  
DN 81:91898  
OREF 81:14577a,14580a  
TI 2,6-Diaminonebularin derivatives  
IN Marumoto, Ryuji; Yoshioka, Yoshio; Honjo, Mikio; Kawazoe, Katsuyoshi  
PA Takeda Chemical Industries, Ltd.  
SO Ger. Offen., 21 pp.  
CODEN: GWXXBX

McIntosh

10/598,520

DT Patent  
LA German  
FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	DE 2359536	A1	19740612	DE 1973-2359536	19731129
	DE 2359536	C2	19840802		
	JP 49080096	A	19740802	JP 1972-123602	19721208
	JP 55049594	B	19801212		
	JP 55049596	B	19801212	JP 1973-114542	19731011
	JP 50064296	A	19750531		
PRAI	JP 1972-123602	A	19721208		
	JP 1973-114542	A	19731011		

GI For diagram(s), see printed CA Issue.


AB Diaminonebularines I (R = Ph, cyclohexyl, p-MeOC<sub>6</sub>H<sub>4</sub>, p-MeC<sub>6</sub>H<sub>4</sub>, p-ClC<sub>6</sub>H<sub>4</sub>, p-methylcyclohexyl) were prepared by treating a 2-halo-adenosine with RNH<sub>2</sub> or by treating a 2-halo-inosine with RNH<sub>2</sub> and NH<sub>3</sub>. I (R = Ph) had 6.75 times the coronary vasodilator activity of adenosine and at 15  $\gamma$ /ml caused 38% inhibition of blood platelet aggregation.

IT 53296-10-9P 53296-19-8P  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(preparation and pharmacol. activity of)

RN 53296-10-9 CAPLUS

CN Adenosine, 2-(phenylamino)- (CA INDEX NAME)


Absolute stereochemistry.



RN 53296-19-8 CAPLUS

CN Adenosine, 2-[(4-methoxyphenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 53296-11-0P 53296-20-1P 53296-21-2P


RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)

RN 53296-11-0 CAPLUS

CN Adenosine, 2-(phenylamino)-, hydrochloride (9CI) (CA INDEX NAME)

Absolute stereochemistry.

10/598,520




● x HCl

RN 53296-20-1 CAPLUS

CN Adenosine, 2-[(4-methylphenyl)amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 53296-21-2 CAPLUS

CN Adenosine, 2-[(4-chlorophenyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L6 ANSWER 193 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1974:116229 CAPLUS

DN 80:116229

OREF 80:18675a,18678a

TI Synthesis and anti-deoxyribonucleic acid virus activity of certain 9- $\beta$ -D-arabinofuranosyl-2-substituted adenine derivatives

AU Miyai, Kenji; Allen, Lois B.; Huffman, John H.; Sidwell, Robert W.; Tolman, Richard L.

CS Nucleic Acid Res. Inst., ICN Pharm. Inc., Irvine, CA, USA

SO Journal of Medicinal Chemistry (1974), 117(2), 242-4

CODEN: JMCMAR; ISSN: 0022-2623

DT Journal

LA English

AB A series of title analogs of 9- $\beta$ -D-arabinofuranosyladenine (I) [5536-17-4], prepared by nucleophilic displacement of chloride from 9-(2,3,5-tri-O-benzyl- $\beta$ -D-arabinofuranosyl)-2-chloroadenine [10212-38-1] followed by hydrogenolytic debenzylation, showed little in vitro antiviral activity. 9- $\beta$ -D-arabinofuranosyl-2-chloroadenine (II) [10147-12-3] had anti-DNA virus activity comparable to I, and increased survivor nos. in mice with virus-induced encephalitis.

IT 51688-48-3P

McIntosh




10/598,520

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(preparation and antiviral activity of)

RN 51688-48-3 CAPLUS

CN 9H-Purin-6-amine, 9- $\beta$ -D-arabinofuranosyl-2-(phenylmethoxy)- (CA INDEX NAME)

Absolute stereochemistry.



L6 ANSWER 194 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1974:48334 CAPLUS

DN 80:48334

OREF 80:7885a,7888a

TI 2-Substituted adenosine 3',5'-cyclomonophosphates

IN Honjo, Mikio; Marumoto, Ryuji; Yoshioka, Yoshio; Takatsuki, Shima

PA Takeda Chemical Industries, Ltd.

SO Ger. Offen., 26 pp.

CODEN: GWXXBX

DT Patent

LA German

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	DE 2324130	A1	19731129	DE 1973-2324130	19730512
	JP 49011896	A	19740201	JP 1972-48372	19720515
	FR 2184813	A1	19731228	FR 1973-17376	19730514
	AU 7355675	A	19741114	AU 1973-55675	19730514
	CA 1000696	A1	19761130	CA 1973-171274	19730514
	HU 169628	B	19761228	HU 1973-TA1253	19730514
	NL 7306790	A	19731119	NL 1973-6790	19730515
	AT 7304243	A	19751215	AT 1973-4243	19730515
	AT 331998	B	19760910		
	GB 1433507	A	19760428	GB 1973-23035	19730515
PRAI	JP 1972-48372	A	19720515		

GI For diagram(s), see printed CA Issue.

AB Twenty-two cyclophosphates I (R = e.g. PhCH<sub>2</sub>S, CH<sub>2</sub>:CHCH<sub>2</sub>, MeOCH<sub>2</sub>CH<sub>2</sub>S, BuO, Ph, 2-furyl, or 4-pyridyl), useful as antiallergic or central nervous system-stimulating drugs, as diuretics, or bronchodilators, were prepared in part as ammonium salts by intramol. cyclization of the adenosine 5'-monophosphates or their activated (e.g. nitrophenyl) esters with dicyclohexylcarbodiimide or a base, e.g. Me<sub>3</sub>COK, resp.


IT 50257-84-6

RL: RCT (Reactant); RACT (Reactant or reagent)  
(phosphorylation of)

RN 50257-84-6 CAPLUS

CN Adenosine, 2-butoxy- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



McIntosh

10/598,520

L6 ANSWER 195 OF 195 CAPLUS COPYRIGHT 2010 ACS on STN  
AN 1973:466747 CAPLUS  
DN 79:66747  
OREF 79:10787a,10790a  
TI Coronary dilating 2-alkoxyadenosines  
IN Yoshioka, Yoshio; Marumoto, Ryuji; Honjo, Mikio; Kwawzoe, Katsuyoshi  
PA Takeda Chemical Industries, Ltd.  
SO Ger. Offen., 22 pp.  
CODEN: GWXXBX  
DT Patent  
LA German  
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	DE 2258378	A1	19730614	DE 1972-2258378	19721129
	JP 48061498	A	19730828	JP 1971-97431	19711201
	JP 48076894	A	19731016	JP 1972-8885	19720124
	AU 7249412	A	19740530	AU 1972-49412	19721129
	BE 792155	A1	19730530	BE 1972-124819	19721130
	NL 7216299	A	19730605	NL 1972-16299	19721130
	FR 2162128	A1	19730713	FR 1972-42673	19721130
PRAI	JP 1971-97431	A	19711201		
	JP 1972-8885	A	19720124		

GI For diagram(s), see printed CA Issue.

AB Twenty adenosines I (R = e.g., MeOCH<sub>2</sub>CH<sub>2</sub>, BuOCH<sub>2</sub>CH<sub>2</sub>, Ph, Et, Pr, Bu, C<sub>5</sub>H<sub>11</sub>, CH<sub>2</sub>:CHCH<sub>2</sub>, 3-MeC<sub>6</sub>H<sub>4</sub>, Me<sub>2</sub>CH) were prepared by reaction of 2-chloro- or 2-bromoadenosine with ROH in the presence of NaOR, KOR, KOH, NaOH, or Ca(OH)<sub>2</sub>. I had coronary dilating activities in dogs.


IT 50257-82-4P 50257-84-6P 50257-85-7P  
50257-89-1P 50257-90-4P 50257-91-5P  
50257-93-7P 50257-95-9P 50257-97-1P  
50447-11-5P

RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)

RN 50257-82-4 CAPLUS

CN Adenosine, 2-phenoxy- (CA INDEX NAME)


Absolute stereochemistry.



RN 50257-84-6 CAPLUS

CN Adenosine, 2-butoxy- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 50257-85-7 CAPLUS

CN Adenosine, 2-(pentyloxy)- (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 50257-89-1 CAPLUS  
CN Adenosine, 2-(2-hydroxyethoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 50257-90-4 CAPLUS  
CN Adenosine, 2-(3-methylphenoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 50257-91-5 CAPLUS  
CN Adenosine, 2-(4-methoxyphenoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 50257-93-7 CAPLUS  
CN Adenosine, 2-(4-hydroxybutoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


McIntosh

10/598,520




RN 50257-95-9 CAPLUS  
CN Adenosine, 2-(hexyloxy)- (CA INDEX NAME)

Absolute stereochemistry.




RN 50257-97-1 CAPLUS  
CN Adenosine, 2-(2-chlorophenoxy)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 50447-11-5 CAPLUS  
CN Adenosine, 2-[(6-hydroxyhexyl)oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



OSC.G 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD (3 CITINGS)

=> d his

(FILE 'HOME' ENTERED AT 13:31:46 ON 17 NOV 2010)

FILE 'REGISTRY' ENTERED AT 13:31:59 ON 17 NOV 2010

L1 STRUCTURE UPLOADED

L2 15 S L1

McIntosh

10/598,520

L3               STRUCTURE UPLOADED  
L4               8 S L3  
L5               148 S L3 FULL

FILE 'CAPLUS' ENTERED AT 13:40:10 ON 17 NOV 2010  
L6               195 S L5

=> file reg

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	1135.45	1333.58
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE ENTRY	TOTAL SESSION
CA SUBSCRIBER PRICE	-165.75	-165.75

FILE 'REGISTRY' ENTERED AT 13:43:01 ON 17 NOV 2010  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2010 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file  
provided by InfoChem.

STRUCTURE FILE UPDATES: 16 NOV 2010 HIGHEST RN 1253329-10-0  
DICTIONARY FILE UPDATES: 16 NOV 2010 HIGHEST RN 1253329-10-0

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH June 26, 2010.

Please note that search-term pricing does apply when  
conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and  
predicted properties as well as tags indicating availability of  
experimental property data in the original document. For information  
on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/stndoc/properties.html>